

TANGENT SPACES AND GROMOV-HAUSDORFF LIMITS OF SUBANALYTIC SPACES

ANDREAS BERNIG AND ALEXANDER LYTCHAK

ABSTRACT. It is shown that the Gromov-Hausdorff limit of a subanalytic 1-parameter family of compact connected sets (endowed with the inner metric) exists. If the family is semialgebraic, then the limit space can be identified with a semialgebraic set over some real closed field. Different notions of tangent cones (pointed Gromov-Hausdorff limits, blow-ups and Alexandrov cones) for a closed connected subanalytic set are studied and shown to be naturally isometric. It is shown that geodesics have well-defined Euclidean directions at each point.

1. INTRODUCTION

The length metric on subanalytic spaces is far from being understood. For instance, Hardt's conjecture from 1983 ([11]), claiming that the metric is a subanalytic function, remains open. Only after the introduction of L-regular decompositions by Kurdyka ([12]) and Parusiński ([17]), there has been some progress in understanding the metric structure of subanalytic sets (e.g. [10], [4], [13], [16], [8], [18], [1], [2]). However, it seems that many natural questions are still out of reach.

In this note, we provide results concerning the local structure of subanalytic spaces. Our local results rely on the following global theorem.

THEOREM 1.1. *Let $X \subset \mathbb{R} \times \mathbb{R}^n$ be a compact subanalytic 1-parameter family of subsets of \mathbb{R}^n . Suppose that each fiber $X_t := X \cap (\{t\} \times \mathbb{R}^n)$ is connected. Then the Gromov-Hausdorff limit $\lim_{t \rightarrow 0^+} (X_t, d_{X_t})$ exists.*

Considering for a single subanalytic space X the family $\frac{1}{t}X$ we obtain that the tangent cone $\lim_{t \rightarrow 0} (X, \frac{1}{t}d_X, x)$ of X with respect to the inner metric d_X exists at each point $x \in X$. However, the proof of Theorem 1.1 provides not only an abstract convergence of isometry classes of spaces but gives us an explicit metric space in the limit. This allows us to speak about the tangent space and about differentiability of maps. The next theorem is our main result concerning the infinitesimal geometry of subanalytic sets.

1991 *Mathematics Subject Classification.* 53C22, 32B20.

Key words and phrases. Tangent spaces, Gromov-Hausdorff convergence, subanalytic sets, geodesics.

The first author was supported by Deutsche Forschungsgemeinschaft grant BE 2484/2-1.

THEOREM 1.2. *Let $X \subset \mathbb{R}^n$ be a closed, connected subanalytic set, endowed with the inner metric d_X . Then the Gromov tangent space $T_x X$ is naturally isometric to the Alexandrov cone $C_x X$, i.e. each blow up $X_x^{(t_i)}$ is naturally isometric to $C_x X$. Moreover, these tangent spaces are Euclidean cones. In particular, the angle between two geodesics starting at x is well-defined. Each subanalytic Lipschitz map $f : (X, x) \rightarrow (Y, y)$ is differentiable at x in the metric sense.*

In general, Gromov-Hausdorff convergence is too weak and it does not allow to compare the topology of the limit space with the topology of the elements of the convergent sequence. For example collapsing can occur, i.e. the dimension of the tangent space $T_x X$ may be smaller than the dimension of each neighborhood of x in X . However some structure survives in $T_x X$. Recall that the density of a k -dimensional subanalytic set X at a point $x \in X$ is given by $\theta(X, x) := \lim_{r \rightarrow 0} \frac{\mathcal{H}^k(B_r(x))}{b_k r^k}$ where b_k is the volume of the k -dimensional unit ball. The existence of the limit is proven in [14]. We prove

PROPOSITION 1.3. a) *The unit sphere in $T_x X$ is connected if and only if this is the case for small spheres $S_r(x)$ in X .*
b) *The density of X at x equals the ratio of the Hausdorff measure of the unit ball in $T_x X$ and the volume b_k of the k -dimensional Euclidean ball.*

Each subanalytic subset $X \subset \mathbb{R}^n$ defines at each point $x \in X$ the so called subanalytic tangent cone $T_x^{sub} X \subset \mathbb{R}^n$ (which is the same as the tangent cone of the metric space (X, d_e) , where d_e is the Euclidean metric on X). In general, T_x^{sub} contains less information than the metric tangent cone $T_x X$, but they are closely related:

THEOREM 1.4. *Let X be as above. Then the identity $\iota : (X, d_X) \rightarrow (X, d_e)$ is differentiable at x . The differential $D_x \iota : T_x X \rightarrow T_x^{sub} X$ is a 1-Lipschitz homogeneous map that preserves the distances to the origin and lengths of arbitrary curves. Moreover each point $v \in T_x^{sub} X$ has at most $m = m(X)$ preimages.*

The known results about the inner metric structure of subanalytic sets are too weak to exclude strange behavior (like oscillations) of geodesics. One of the keys for the proof of the above results is a regularity result for geodesics. We show that each geodesic in (X, d_X) has a well defined Euclidean direction at each point.

Remark 1.1. Theorem 1.1 and Theorem 1.4 are valid for every o-minimal structure (see [19] for o-minimal structures). . Theorem 1.2 and Proposition 1.3 are valid for every polynomially bounded o-minimal structure.

In the last section, we will study the semialgebraic case. It turns out that every closed, rationally bounded semialgebraic set S in \mathbb{R}^n , where R

is any real closed field, defines a compact inner metric space (\bar{S}, \bar{d}_S) . The Gromov-Hausdorff limit of a semialgebraic 1-parameter family of such spaces is again semialgebraic, the limit space is the fiber over the point 0^+ in the real spectrum of $R[t]$.

THEOREM 1.5. *Let R be a real closed field. Let $X \subset R \times R^n$ be a semialgebraic 1-parameter family of subsets of R^n . Suppose that $X \subset B_a(0)$ for some natural number a and that each fiber $X_t := X \cap (\{t\} \times R^n)$ is semialgebraically connected. Let $S := X_{0^+}$ be the fiber over $0^+ \in \text{Spec}_r R[t]$. Then $\lim_{t \rightarrow 0^+} (\bar{X}_t, \bar{d}_{X_t}) = (\bar{S}, \bar{d}_S)$.*

Remark that S is again semialgebraic, but over the real closed field $k(0^+)$, which is the field of algebraic Puiseux series over R . The analogous convergence result for the induced metrics on the fibers X_t instead of the inner metrics was known to be true for some while, see [7].

The main technical ingredient in our proofs is a decomposition, due to Kurdyka and Orro ([13]), of a given subanalytic set into subanalytic pieces such that inner and Euclidean metric differ only by a factor near 1. For further information concerning the inner metric on subanalytic or more general stratified spaces, we refer the reader to [18] and [16]. A detailed study of geodesic metric spaces can be found in [6] and [9].

2. TANGENT CONES OF METRIC AND SUBANALYTIC SPACES

2.1. Notations. For a metric space (X, d) we will denote by d_X the inner metric on X (which can be infinite if there is no rectifiable path between two points). The identity $id : (X, d_X) \rightarrow (X, d)$ is 1-Lipschitz and preserves lengths of curves. A geodesic in a metric space (X, d) is an isometric embedding $\gamma : [a, b] \rightarrow X$ of an interval. Subsets of metric spaces will always be considered with the induced metric, if not otherwise stated.

By $B_r(x)$ we will denote the closed metric ball of radius r around x and by rX the metric space (X, rd) .

2.2. Metric cones. Compare [15] for more on metric cones. A metric cone is a pointed metric space (X, d, x) together with a (pointwise) continuous family $\delta_t, t \in \mathbb{R}^+$ of maps (dilations) $\delta_t : (X, x) \rightarrow (X, x)$, such that $d(\delta_t(y), \delta_t(z)) = td(y, z)$ and $\delta_t \circ \delta_s = \delta_{st}$. A map between metric cones is homogeneous if it commutes with the dilations. A metric cone $(T, 0)$ is called radial if for each $x \in T$ with $d(x, 0) = 1$ the map $t \rightarrow \delta_t(x)$ is a ray, i.e. if $d(\delta_t(x), \delta_s(x)) = |t - s|$. If $(T, 0)$ is a radial cone we can consider $S = \{x \in T \mid d(x, 0) = 1\}$ the unit sphere in T and the Euclidean cone CS (compare [9]) over S . Then the natural map $F : CS \rightarrow T$ that sends tx to $\delta_t(x)$ is homogeneous and bilipschitz.

2.3. Ultralimits and blow-ups. See [6] and [15] for more details. We will use a fixed non-principal ultrafilter ω on the set of natural numbers. For pointed metric spaces (X_i, x_i) we will denote their ultralimit by $\lim_\omega (X_i, x_i)$.

Remark that if the isometry classes of proper spaces (X_i, x_i) build a relatively compact set with respect to the pointed Gromov-Hausdorff topology, then the limit $\lim_\omega (X_i, x_i)$ is a proper space and its isometry class is a pointed Gromov-Hausdorff limit of a subsequence of (X_i, x_i) .

For a fixed metric space (X, x) and a zero sequence $(t_i) \rightarrow 0$ we consider the ultralimit $\lim_\omega (\frac{1}{t_i} X, x)$, denote it by $X_x^{(t_i)}$ and call it the blow-up of X at x at the scale (t_i) . The base point of the blow-ups will be denoted by 0.

If X is proper, and if the pointed Gromov-Hausdorff limit $\lim_{t \rightarrow 0} (\frac{1}{t} X, x)$ exists, then all the blow-ups $X_x^{(t_i)}$ are in the isometry class of $\lim_{t \rightarrow 0} (\frac{1}{t} X, x)$.

If $f : (X, x) \rightarrow (Y, y)$ is an L -Lipschitz map, then for each sequence $(t_i) \rightarrow 0$ there is an induced L -Lipschitz blow-up $f_x^{(t_i)} : (X_x^{(t_i)}, 0) \rightarrow (Y_y^{(t_i)}, 0)$.

2.4. Tangent cones and differentials. We refer the reader to [15] for a detailed study of differential properties of general metric spaces.

Let X be a metric space, $x \in X$. We say that a metric cone $(T, 0)$ is the tangent cone $T_x X$ at x , if for each zero sequence (t_i) an isometry $\tau^{(t_i)} : (T, 0) \rightarrow (X_x^{(t_i)}, 0)$ is fixed, such that for each $s > 0$ and each point $p \in T$ the point $\tau^{(st_i)}(\delta_s(p)) \in X_x^{(st_i)}$ coincides with $\tau^{(t_i)}(p) \in X_x^{(t_i)}$ if the sets $X_x^{(t_i)}$ and $X_x^{(st_i)}$ are identified in the natural way.

Remark 2.1. The definition implies that all the blow-ups of X at x are isometric and a fixed metric space $(T, 0)$ in the isometry class of the blow-ups is fixed. The commutation relations required in the definition are always satisfied, if the isometries $\tau^{(t_i)}$ are given in some "natural" way.

If for metric spaces (X, x) and (Y, y) the tangent cones $T_x X$ and $T_y Y$ exist, we say that a Lipschitz map $f : (X, x) \rightarrow (Y, y)$ is differentiable at x if for each zero sequence (t_i) the blow-up $f_x^{(t_i)}$ considered as a map from $T_x X$ to $T_y Y$ does not depend on the sequence (t_i) . This unique blow-up is in this case a homogeneous Lipschitz map. It will be denoted by $D_x f$ and called the differential of f at x . In particular a Lipschitz curve $\gamma : [0, a) \rightarrow X$ starting at x is differentiable at 0 iff the point $v = (\gamma(t_i)) \in X_x^{(t_i)} = T_x X$ is independent of the zero sequence t_i . In this case the differential $D_0 \gamma : T_0([0, a)) = [0, \infty) \rightarrow T_x X$ is given by $D_0 \gamma(t) = \delta_t(v)$. We will identify v with $D_0 \gamma$.

2.5. Alexandrov cone. By Γ_x we denote the set of all geodesics starting at x . On $\Lambda_x := \Gamma_x \times [0, \infty)$ we consider the pseudo metric given by $d((\gamma_1, s_1), (\gamma_2, s_2)) := \limsup_{t \rightarrow 0} \frac{d(\gamma_1(s_1 t), \gamma_2(s_2 t))}{t}$ and call its completion the Alexandrov cone $C_x (= C_x X)$. The points $(\gamma, 0) \in \Gamma_x \times [0, \infty)$ are identified to the origin 0 in C_x and for $t \in \mathbb{R}^+$ the maps $\delta_t : C_x \rightarrow C_x$ are t -dilations, that define the structure of a radial metric cone on C_x .

For each zero sequence $(t_i) \rightarrow 0$ the natural 1-Lipschitz map $\exp_x^{(t_i)} : \Gamma_x \times [0, \infty) \rightarrow X_x^{(t_i)}$ defined by $\exp_x^{(t_i)}((\gamma, s)) := (\gamma(st_i))$ uniquely extends to a 1-Lipschitz map $\exp_x^{(t_i)} : C_x \rightarrow X_x^{(t_i)}$.

Let x be a point in X . Then all the exponential maps $\exp_x^{(t_i)}$ are isometric embeddings iff the limes superior in the definition of the metric on C_x is a limes. The upper angle and the lower angle between each pair of geodesics starting at x coincide iff in addition C_x is a Euclidean cone.

2.6. Subanalytic tangent cone. Let $T_x^{sub}X$ denote the subanalytic tangent cone of X at x , i.e. the set

$$T_x^{sub}X := \{v \in \mathbb{R}^n : \forall \epsilon > 0 \exists y \in X \exists \lambda \in [0, \infty) : \|y-x\| < \epsilon, \|\lambda(y-x)-v\| < \epsilon\}$$

By the curve selection lemma, this is the same as the set of initial vectors of continuous subanalytic curves starting at x and contained in X . Note that $T_x^{sub}X$ is a subanalytic cone ([14]).

Remark 2.2. The cone $T_x^{sub} \subset \mathbb{R}^n$ equals the (metric) tangent cone to the metric space $(X, d_e) \subset \mathbb{R}^n$ at x . A Lipschitz curve $\gamma : [0, \epsilon) \rightarrow X$ starting at x is differentiable as a map to (X, d_e) , iff γ is differentiable at 0 as a curve in \mathbb{R}^n . A subanalytic map $f : (X, x) \rightarrow (Y, y)$ that is Lipschitz with respect to the induced metric defines a homogeneous Lipschitz differential $D_x^{sub}f : T_x^{sub}X \rightarrow T_y^{sub}Y$.

3. GROMOV-HAUSDORFF LIMIT IN A SUBANALYTIC FAMILY

This section is devoted to the proof of Theorem 1.1.

Proof. Let $X \subset \mathbb{R} \times \mathbb{R}^n$ be a compact subanalytic set such that $X_t := X \cap (\{t\} \times \mathbb{R})$ is connected.

We use the known fact ([13]) that for each $C > 1$ the set X can be decomposed as finite union $X = \cup_{i=1}^m X^i$ such that each fiber X_t^i is subanalytic, compact, connected with the property that length and induced metric on X_t^i differ by at most a factor C .

It follows immediately that the diameters of the metric spaces (X_t, d_{X_t}) , $t \in \mathbb{R}$ are uniformly bounded from above by some $0 < D < \infty$.

It also follows that the family is equicompact, i.e. for each $\epsilon > 0$, there exists $N_1(\epsilon)$, independent of t , such that each X_t can be covered by at most $N_1(\epsilon)$ balls of radius ϵ . Equivalently, there exists $N_2(\epsilon)$ such that each ϵ -separated net in X_t contains at most $N_2(\epsilon)$ points.

Consider two subanalytic curves $\gamma_1, \gamma_2 : (0, \epsilon) \rightarrow X$ with $\gamma_i(t) \in X_t$. We claim that the limit $\lim_{t \rightarrow 0+} d_{X_t}(\gamma_1(t), \gamma_2(t)) \in [0, \infty)$ exists. This follows from the theorem of Kurdyka-Orro ([13]) stating that for each $\eta > 0$ there exists a subanalytic distance $\tilde{d} : X \times X \rightarrow \mathbb{R}$ such that $\tilde{d}(x, y) \leq d_{X_t}(x, y) \leq (1 + \eta)\tilde{d}(x, y)$ for all $x, y \in X_t$. Since the limit $\lim_{t \rightarrow 0+} \tilde{d}(\gamma_1(t), \gamma_2(t))$ exists in $[0, 2D]$ (by properties of subanalytic functions), we obtain that $\lim_{t \rightarrow 0+} d_{X_t}(\gamma_1(t), \gamma_2(t)) \in [0, 2D]$ exists.

On the space Λ^{sub} of subanalytic curve germs $\gamma : (0, \epsilon) \rightarrow X$ with $\gamma(t) \in X_t$ this limit defines a pseudo-metric d_{\lim} .

Let $\gamma_1, \dots, \gamma_k$ be an ϵ -separated net in Λ^{sub} . Then, for small enough $t > 0$, $\gamma_1(t), \dots, \gamma_k(t)$ is a 2ϵ -separated net in X_t . By equicompactness,

we get $k \leq N_2(2\epsilon) < \infty$. The pseudo metric space (Λ^{sub}, d_{lim}) is therefore totally bounded and its completion (X_{lim}, d_{lim}) is a compact metric space (by the theorem of Hausdorff).

We claim that (X_{lim}, d_{lim}) is the Gromov-Hausdorff limit $\lim_{t \rightarrow 0+} X_t$. By total boundedness of Λ^{sub} , there exists a finite ϵ -dense net $\gamma_1, \dots, \gamma_k \in \Lambda^{sub}$. From the theorem of Kurdyka-Orro we infer the existence of a subanalytic distance \tilde{d} with $\tilde{d} \leq d_{X_t} \leq 2\tilde{d}$ for each t . If the subanalytic set $\{(t, x) : x \in X_t, \tilde{d}(x, \gamma_i(t)) > 2\epsilon, i = 1, \dots, k\}$ contains points with arbitrarily small $t > 0$, the curve selection lemma implies that there is a subanalytic curve $\gamma \in \Lambda^{sub}$ contained in it. But then $d_{lim}(\gamma, \gamma_i) \geq 2\epsilon$ for $i = 1, \dots, k$ which is a contradiction. It follows that $\gamma_1(t), \dots, \gamma_k(t)$ form, for t sufficiently small, a 4ϵ -dense net in X_t .

Since the Gromov-Hausdorff distance between a compact metric space and an ϵ -dense net is at most ϵ , and since the Gromov-Hausdorff distance between $(\{\gamma_1, \dots, \gamma_k\}, d_{lim})$ and $(\{\gamma_1(t), \dots, \gamma_k(t)\}, d_{X_t})$ tends to 0, the triangle inequality implies that $d_{G-H}(X_{lim}, X_t) \rightarrow 0$ for $t \rightarrow 0$. \square

Now we consider more closely the case of the tangent space. Let $X \subset \mathbb{R}^n$ be a closed subanalytic subset, $x \in X$. Without loss of generality we assume $x = 0$. Define $Y \subset \mathbb{R} \times \mathbb{R}^n$ as the set of all points $y = (t, x)$ with $t > 0$ and $\frac{1}{t}x \in X$. For $r > 0$ denote by Y^r the subset of all $y = (t, x) \in Y$ with $\|\frac{1}{t}x\| \leq r$. The fiber Y_t^r of the family Y^r is just the ball $B_{rt} \subset X$ with the metric rescaled by $\frac{1}{t}$. By the local conical structure of X , it is connected for each $r > 0$ and all sufficiently small t . Therefore the spaces Y_t^r considered with the inner metric are equicompact.

This and the proof of Theorem 1.1 above show, that the pointed spaces $(Y_t, d_{Y_t}, (t, 0)) = (X, \frac{1}{t}d_X, x)$ converge in the pointed Gromov-Hausdorff topology to the space Λ of all continuous subanalytic curves γ with $\gamma(t) \in Y_t$, where the metric is defined as in the proof above.

Let $\Lambda_x^{sub}X$ be the set of all continuous subanalytic curves in X starting in x such that $\lim_{t \rightarrow 0} \frac{\|\eta(t) - x\|}{t} < \infty$. Considering the map $\Lambda \rightarrow \Lambda_x^{sub}X$ defined by $\gamma \rightarrow \eta : \eta(t) = t\gamma(t) \in X$, we conclude from the arguments above, that $d(\eta_1, \eta_2) := \lim_{t \rightarrow 0} \frac{d_X(\eta_1(t), \eta_2(t))}{t}$ defines a metric on $\Lambda_x^{sub}X$ and that the spaces $(X, \frac{1}{t}d_X, x)$ converge to $\Lambda_x^{sub}X$ in the pointed Gromov-Hausdorff topology.

As was already mentioned in the introduction we get more than just an abstract Gromov-Hausdorff convergence.

Corollary 3.1. *Let $x \in X$, where X is closed subanalytic. Then the tangent cone of (X, d_X) at the point x exists and is given by completion of the pseudo-metric space $(\Lambda_x^{sub}X, d)$.*

Proof. The dilatations on $\Lambda_x^{sub}X$ are given by linear reparameterizations and induce natural dilatations on T_xX .

The “exponential maps” $\exp^{(t_i)} : \Lambda_x^{sub} X \rightarrow X_x^{(t_i)}, \eta \mapsto (\eta(t_i))$ extend to isometries on the completion $T_x X$. The commutation relations required in Subsection 2.4 are obviously satisfied. \square

Observe that each $\gamma \in \Lambda_x^{sub} X$ starting at x has a well defined initial direction $v \in T_x X$ (i.e. γ is differentiable at 0 as a map from the interval $[0, \epsilon]$ to (X, d_X) if γ is Lipschitz).

Let now $f : (X, x) \rightarrow (Y, y)$ be a subanalytic Lipschitz map (with respect to the inner metrics) between subanalytic sets. One gets a well defined homogeneous Lipschitz differential $D_x f : \Lambda_x^{sub} X \rightarrow \Lambda_y^{sub} Y, \gamma \mapsto f \circ \gamma$ which extends to a Lipschitz differential $D_x f : T_x X \rightarrow T_y Y$.

4. REGULARITY OF GEODESICS

Let $X \subset \mathbb{R}^n$ be a closed subanalytic set, $x \in X$ a point.

Lemma 4.1. *There are $C, \alpha, r > 0$ (depending on X and x) such that for each $z \in X$ with $\|z - x\| \leq r$ there is a Lipschitz curve γ in X of length at most $\|z - x\| + C\|z - x\|^{1+2\alpha}$ connecting x and z .*

Proof. Stratify X such that Whitney’s condition A is satisfied for each pair of strata. Consider on $X \setminus \{x\}$ the stratified vector field V such that $V(y)$ is the projection of $\frac{x-y}{\|x-y\|}$ onto $T_y S$, where S is the stratum containing y . Of course $\|V(y)\| \leq 1$. Define the subanalytic function $g(t) := \sup \left\{ \|V(y) - \frac{x-y}{\|x-y\|}\| : y \in X, y \neq x, \|y - x\| \leq t \right\}$. If $g(t)$ does not tend to 0 for $t \rightarrow 0$, there is a sequence (y_i) tending to x contained in one single stratum S such that the angle between the tangent space $T_{y_i} S$ and the line between x and y_i does not tend to 0, in contradiction to Whitney’s condition A. By Łojasiewicz’ inequality we get $g(t) \leq C_1 t^{2\alpha}$ for some constants $\alpha, C_1 > 0$.

Consider now a maximal integral curve γ of V starting at z in the stratum containing z . It converges to a unique point z_1 in a stratum of smaller dimension. Then continue on the maximal integral curve of V starting in z_1 and so on. After finitely many steps we get a Lipschitz curve γ connecting z and x . Let s be the smallest real such that $\gamma(s) = 0$ and set $\bar{\gamma}(t) := \gamma(s - t), t \in [0, s]$. From $\|V(y)\| \leq 1$ we get $L(\bar{\gamma}) = L(\gamma) \leq s$ and $f(t) := \|\bar{\gamma}(t) - x\| \leq t$ for $t \in [0, s]$.

Then

$$\begin{aligned} \frac{d}{dt} f^2 &= 2\langle \bar{\gamma}'(t), \bar{\gamma}(t) - x \rangle \\ &= 2\left\langle -V(\bar{\gamma}(t)) + \frac{x - \bar{\gamma}(t)}{\|x - \bar{\gamma}(t)\|}, \bar{\gamma}(t) - x \right\rangle - 2\left\langle \frac{x - \bar{\gamma}(t)}{\|x - \bar{\gamma}(t)\|}, \bar{\gamma}(t) - x \right\rangle \\ &\geq 2f(t) - 2C_1 t^{2\alpha} f(t) \end{aligned}$$

We conclude that (for $t > 0$) $f'(t) \geq 1 - C_1 t^{2\alpha}$ and therefore $\|z - x\| = f(s) = \int_0^s f'(t) dt \geq s - C_2 s^{1+2\alpha}$ with $C_2 := \frac{C_1}{1+2\alpha}$.

For s sufficiently small, $C_2 s^{2\alpha} \leq \frac{1}{2}$. Replacing this yields $s \leq 2\|z - x\|$ and finally $L(\gamma) \leq s \leq \|z - x\| + C\|z - x\|^{1+2\alpha}$ with $C := 2^{1+2\alpha}C_2$. \square

Remark 4.1. This lemma can be reformulated in terms of the identity map $\iota : (X, d_X) \rightarrow (X, d_e)$ as follows. For each $z \in X$ holds $d_e(\iota(z), \iota(x)) \geq d_X(x, z) - Cd_X(x, z)^{1+2\alpha}$, for some C, α depending on x . Moreover we see that by the pointed Gromov-Hausdorff convergence $(X, \frac{1}{t}d_X, x) \rightarrow T_x X$ the intersections $(S_t(x), \frac{1}{t}d_X)$ of the Euclidean spheres with X converge to the unit sphere in S in $T_x X$.

Now we derive:

PROPOSITION 4.2. *Let $\gamma : [0, t] \rightarrow X$ be a geodesic starting at x . Then γ , considered as a curve in \mathbb{R}^n , has a unique direction γ^+ at 0. Moreover, there are $r, C, \alpha > 0$ depending only on X and x such that for all $0 < t \leq r$*

$$\left\| \gamma^+ - \frac{\gamma(t) - x}{\|\gamma(t) - x\|} \right\| \leq Ct^\alpha.$$

Proof. We choose r, C, α as in Lemma 4.1 and $0 < t \leq r$. Let first s be a number with $\frac{t}{2} \leq s \leq t$. Put $z = \gamma(t)$ and $y = \gamma(s)$. In the triangle xyz we know $\|x - y\| \leq s$, $\|y - z\| \leq t - s$ and $\|x - z\| \geq t - Ct^{1+2\alpha}$, with $C = C(X, x)$.

Using the cosine law for the (Euclidean) triangle xyz we get that the angle at x between xz and xy is at most $\tilde{C}t^\alpha$, for some $\tilde{C} = \tilde{C}(C)$.

Thus the directions $v_t := \frac{\gamma(t) - x}{\|\gamma(t) - x\|}$ satisfy $\|v_t - v_s\| \leq Ct^\alpha$ for each $t > s \geq \frac{t}{2}$. From this we immediately conclude that v_t converge for $t \rightarrow 0$ to some v . Moreover we get $\|v_t - v\| \leq \sum_{i=0}^{\infty} \|v_{2^{-i}t} - v_{2^{-i-1}t}\| \leq Ct^\alpha \sum_{i=0}^{\infty} (2^\alpha)^{-i} = \tilde{C}t^\alpha$, with some \tilde{C} depending on C and α . \square

5. COMPARISON BETWEEN THE METRIC AND THE SUBANALYTIC TANGENT CONE

The natural embedding $\iota_X : (X, d_X) \rightarrow \mathbb{R}^n$ is subanalytic and 1-Lipschitz. Hence ι_X is differentiable at x with differential $D_x \iota_X : T_x X \rightarrow T_x \mathbb{R}^n = \mathbb{R}^n$ (see Section 3). The image $D_x \iota_X(T_x X)$ coincides with $T_x^{sub} X$. Due to Remark 4.1, $D_x \iota_X$ preserves distances to the origin. If $f : (X, x) \rightarrow (Y, y)$ is a subanalytic Lipschitz map with respect to the induced metrics, then f is also Lipschitz with respect to the length metrics and the differentials commute, i.e. $D_x^{sub} f \circ D_x \iota_X = D_y \iota_Y \circ D_x f$.

Fix $\epsilon > 0$ and let again $X = \cup_{j=1}^m X_j$ be a decomposition in subanalytic sets such that inner and induced metric agree up to a factor $1 + \epsilon$ on each of it. The injection $\tau_j : X_j \rightarrow X$ is subanalytic, hence it induces a 1-Lipschitz differential $D_x \tau_j : T_x X_j \rightarrow T_x X$. Denote by $\tilde{T}_x X_j$ the image $D_x \tau_j(T_x X_j) \subset T_x X$. Remark that $T_x X = \cup_{j=1}^m \tilde{T}_x X_j$.

Since the restriction of d_X to X_j is $(1 + \epsilon)$ -bilipschitz equivalent to the induced metric on X_j , the map $D_x \iota_X : \tilde{T}_x X_j \rightarrow D_x \iota_X(\tilde{T}_x X_j)$ is $(1 + \epsilon)$ -bilipschitz. In particular this restriction is injective. This shows that all

fibers of the map $D_x \iota_X : T_x X \rightarrow T_x^{sub} X$ have at most m elements. Moreover $\tilde{T}_x X$ has a finite decomposition such that each set of this decomposition is mapped $(1 + \epsilon)$ -bilipschitz under $D_x \iota_X$ onto its image in $T_x^{sub} X$. Since this holds for each $\epsilon > 0$ we see that $D_x \iota_X$ preserves lengths of curves. These observations complete the proof of Theorem 1.4.

Now we are going to prove that the tangent cone $T_x X$ is a Euclidean cone. This result is a direct consequence of Theorem 1.4, the fact that $T_x^{sub} X$ is a Euclidean cone as a subcone of \mathbb{R}^n and the following:

Lemma 5.1. *Let T be a metric cone that is in addition a geodesic metric space. Let $\iota : T \rightarrow CV$ be a homogeneous 1-Lipschitz arclengths preserving map onto an Euclidean cone CV , that preserves the distance to the origin. Then T is an Euclidean cone.*

Proof. Let S be the unit sphere in T . It is mapped by ι to V . For $x \in S$ the image $\iota(\delta_t(x)) = \delta_t(\iota(x))$ is a radial ray in CV . Since ι preserves the lengths of curves we see that $\delta_t(x)$ and $\delta(s)(x)$ have distance at most $|s - t|$. Therefore T is a radial cone.

The restriction $\iota : S \rightarrow V$ is again 1-Lipschitz and preserves lengths of curves. Let \tilde{S} be the set S considered with the inner metric. Consider the Euclidean cone $C\tilde{S}$ and the natural bijection $F : C\tilde{S} \rightarrow T$. Observe that the composition $\iota \circ F : C\tilde{S} \rightarrow CV$ preserves the lengths of curves. Since so does the map $\iota : T \rightarrow CV$ and since $C\tilde{S}$ and T are geodesic metric spaces, it is enough to see that the bijection F preserves the class of Lipschitz curves. But so does the natural 1-Lipschitz map $Id : C\tilde{S} \rightarrow CS$ and by Subsection 2.2 the natural map $F : CS \rightarrow T$ is bilipschitz. This finishes the proof. \square

6. CONNECTIVITY

Proof of Theorem 1.3, a). Let X be a closed subanalytic space, $x \in X$. By local conical structure of X , the (Euclidean) ball $B_r(x)$ around x is homeomorph to the cone over the (Euclidean) sphere $S_r(x)$ for $r > 0$ small enough. Suppose that $S_r(x)$ is not connected and let S_1, S_2, \dots, S_k be its connected components. They correspond to connected components B_1, \dots, B_k of $B_r(x) \setminus \{x\}$. Since each d_X -geodesic between points from different components B_i and B_j must run through x , we see that for the closed subcones $T_x B_i$ of $T_x X$ holds $T_x B_i \cap T_x B_j = \{0\}$. Therefore $T_x B_i \setminus \{0\}$ is open and closed in $T_x X \setminus \{0\}$ and the unit sphere in $T_x X$ is not connected.

Assume on the other hand that S_r is connected for small r . Observe that by the convergence $(X, \frac{1}{t}d_X, x) \rightarrow T_x X$ the spheres $(S_r, \frac{1}{r}d_X)$ converge to the unit sphere in $T_x X$. But $\frac{1}{r}S_r$ is a subanalytic family of bounded connected subanalytic subsets for $r > 0$. By the equicompactness of the family (see Section 3) the fibers S_r of the family have uniformly bounded diameters with respect to the inner metrics d_{S_r} . Hence the unit sphere in $T_x X$ is connected. \square

For a vector $v \in \mathbb{R}^n$ and $\rho > 0$ denote by $K(v, \rho)$ the set of all vectors $w \in \mathbb{R}^n$ with $\langle v, w \rangle \geq (1 - \rho)\|v\|\|w\|$. In the next section we will use:

Lemma 6.1. *Let X be a subanalytic space, $0 \in X$. Let $v \in T_0^{sub}X$ be a direction. For the canonical map $D_0\iota_X : T_0X \rightarrow T_0^{sub}X$ let I be the finite set $D_0\iota_X^{-1}(v)$. Then for some $\rho > 0$ the intersection Y of X with the cone $K(v, \rho)$ has the property, that different points of I lie in different components of $T_0Y \setminus \{0\}$*

Proof. If $v = 0$ then $D_0\iota_X^{-1}(v) = \{0\}$ and the claim is trivial. If $v \neq 0$ we may assume that $\|v\| = 1$. Let $5s$ be the minimal distance of two points in I . Let U_1, U_2, \dots be a sequence of neighborhoods of v in S^{sub} with diameters tending to 0. If for each $i = 1, 2, \dots$ there exists a point $w_i \in D_0\iota_X^{-1}(U_i)$ with $d(w_i, I) \geq s$, we can (by compactness of the unit sphere $S \subset T_xX$) extract a converging subsequence of (w_i) . If w denotes its limit, then $d(w, I) \geq s$ and $D_0\iota_X(w) = v$, contradiction. Therefore, for some i , the intersections of $D_0\iota_X^{-1}(U_i)$ with balls of radius $2s$ around points of I are open and closed in $D_0\iota_X^{-1}(U_i)$. Taking ρ so small that $K(v, \rho) \cap S^{sub}$ is contained in this neighborhood U_i , we obtain the result. \square

7. FINER PROPERTIES OF THE TANGENT CONE

Now we are going to compare the tangent cone with the Alexandrov cone at the given point. We start with:

Lemma 7.1. *Let X be a closed connected subanalytic space and suppose $0 \in X$. Let γ be a Lipschitz curve in X starting at 0, that has a Euclidean differential $v \in T_0^{sub}X$ at 0. Then γ is also differentiable at 0 as a curve into (X, d_X)*

Proof. Consider the cone $K(v, \rho)$ as in Lemma 6.1 and set $Y = X \cap K(v, \rho)$. Then a small starting part of γ is contained in Y and it is sufficient to prove that γ is differentiable at 0 as map into (Y, d_Y) . We may assume that $\|v\| \neq 0$. Then a beginning part of γ is contained in a connected component C of $(B_r(0) \cap Y) \setminus \{0\}$. By Theorem 1.3 a) and Lemma 6.1 the restriction $\iota : T_0C \rightarrow T_0^{sub}C$ has the property that $\iota^{-1}(v)$ has only one point w . But for each zero sequence t_i the point $(\gamma(t_i)) \in C_0^{(t_i)} = T_0C$ is mapped by ι onto v . Hence $(\gamma(t_i)) = w$ and we are done. \square

Together with Proposition 4.2 this shows that geodesics are differentiable as maps into (X, d_X) . In particular for geodesics γ_1 and γ_2 starting at x and for each $s \geq 0$ the generalized angle $\lim_{t \rightarrow 0} \frac{d(\gamma_1(st), \gamma_2(t))}{t}$ is well defined. Hence the exponential maps $\exp_x^{(t_i)} : C_xX \rightarrow T_xX$ are isometric embeddings. Since T_xX is a Euclidean cone, the same is true for C_xX and the usual angle between geodesics is well defined too.

The next lemma finishes the proof of Theorem 1.2:

Lemma 7.2. *In the above notations the exponential maps $\exp_x^{(t_i)} : C_x X \rightarrow X_x^{(t_i)} = T_x X$ are surjective.*

Proof. It is enough to prove that for a curve $\eta \in \Lambda_x^{sub}$, with starting direction v in the unit sphere S of $T_x X$, a geodesic γ_n between x and $\eta(\frac{1}{n})$ with starting direction $v_n \in S$ holds $v_n \rightarrow v$. Consider again the intersection Y of X with a small cone $K(\iota(v), \rho)$ as in the last lemma. Let C be the component of $(B_r(x) \cap Y) \setminus \{x\}$ containing η . Due to Lemma 4.1 the geodesics γ_n are contained in Y for big n . Moreover $\iota(v_n)$ converge to $\iota(v)$. Hence for each limit point w of v_n holds $\iota(w) = \iota(v)$. Since v is the only preimage point of $\iota(v)$ in $T_0 C$, we obtain $v = w$ and finish the proof. \square

Remark 7.1. The (natural) equality between the tangent cone $T_x X$ and the Alexandrov cone $C_x X$ implies directly (compare [15]), that each isometry between subanalytic spaces is differentiable at each point. This reflects the fact, that the tangent cones, although they are defined in subanalytic terms, are in fact purely metric invariants of X , not only as isometry classes (which is trivial) but as metric spaces.

8. MEASURE AND DIMENSION

Let X be a compact subanalytic set. Then due to the decomposition of X in pieces where d_X and d_e coincide up to some factor near 1, we see that the identity $\iota : (X, d_x) \rightarrow (X, d_e)$ preserves the Hausdorff measures \mathcal{H}^l , for each $l \in \mathbb{R}^+$.

On the other hand the Hausdorff dimension k of X coincides with its topological dimension and is given by the dimension of the maximal stratum in a stratification of X . Moreover the k -dimensional Hausdorff measure is finite on bounded subsets and is positive on each k -dimensional subanalytic subset.

The same statements hold true for the subanalytic tangent cone $T_x^{sub} X$. Since $T_x X$ has a finite decomposition such that $D_x \iota : T_x X \rightarrow T_x^{sub} X$ is almost 1-bilipschitz on pieces of the decomposition, we see that the Hausdorff dimension of $T_x X$ coincides with the Hausdorff dimension of $T_x^{sub} X$. Moreover if the restriction of $D_x \iota$ onto a subset U of $T_x X$ is injective, then $D_x \iota : U \rightarrow D_x \iota(U)$ preserves the Hausdorff measure. Finally restricting ι to the preimage of a maximal stratum of a stratification of $T_x^{sub} X$, we see that $T_x X$ contains an open subset homeomorph to a $\dim(T_x^{sub} X)$ -dimensional ball. Hence the topological dimension of $T_x X$ coincides with its Hausdorff dimension.

Proof of Theorem 1.3, b). Choose $\epsilon > 0$ and a decomposition $X = \cup_{i=1}^m X_i \cup Y$ such that X_1, \dots, X_m are ϵ -analytic pieces ([14]) and $\dim Y < k$. Then $\dim(T_x Y) = \dim(T_x^{sub} Y) \leq \dim Y < k$. Since $T_x X_i \cap T_x X_j \subset T_x X$ is contained in $T_x Y$ it is therefore enough to prove the proposition in the case where X is the closure of a single subset X_i . In this case $T_x X = T_x^{sub} X$ and the result was shown in ([14], Proposition 3.6. and Theorem 3.8). \square

Remark 8.1. Suppose that $v \in T_x^{sub}X$ is moreover contained in the pure tangent cone ([14]). Then the multiplicity $n(x)$ defined as in [14] equals the cardinality of the fiber of the map $D_x\iota : T_xX \rightarrow T_x^{sub}$ above v . This follows by a similar argument as above.

9. SEMIALGEBRAIC METRIC SPACES

In a naive sense, the Gromov-Hausdorff limit of a semialgebraic family will not be semialgebraic. Consider for instance a family of ellipsoids getting thinner and thinner. The limit space is a double disc, which is not isometric to the semialgebraic limit consisting of a single disc. But we will show that such a limit space can be obtained as inner metric space associated to a semialgebraic set over some real closed field R .

For the notions of semialgebraic sets, real spectrum and fiber of a semialgebraic family over a point in the real spectrum we refer to [5].

Let R be a real closed field. Let A be the convex hull of $\mathbf{Z} \subset R$. Then A is a valuation ring of R . We denote by m_A its maximal ideal and by $\pi : A \rightarrow A/m_A$ the canonical projection. The field A/m_A is archimedean and can therefore be uniquely identified with a subfield of \mathbb{R} . We define the real place $\lambda_R : R \rightarrow \mathbb{R} \cup \{\infty\}$ by setting $\lambda_R(x) = \infty$ if $x \notin A$ and $\lambda_R(x) := \pi(x)$ if $x \in A$.

An alternative way to define λ_R is given by $\lambda_R(x) = \inf\{r \in \mathbf{Q} : r > x\}$ (with the convention $\inf \emptyset = \inf \mathbf{Q} = \infty$).

Let $S \subset R^n$ be a closed connected semialgebraic set. A path in S is a continuous map $\gamma : [0, 1] \rightarrow S$, where $[0, 1] \subset R$ is the closed unit interval in R . The length of γ is defined by

$$l(\gamma) := \sup \left\{ \lambda_R \left(\sum_{i=0}^{k-1} \|\gamma(t_{i+1}) - \gamma(t_i)\| \right) : 0 = t_0 < t_1 < \dots < t_k = 1 \right\}$$

The distance between two points $x, y \in S$ is defined by

$$d_S(x, y) := \inf\{l(\gamma) : \gamma \text{ is a path between } x \text{ and } y\} \in \mathbb{R} \cup \{\infty\}$$

Note that this is a real number and not a number in R .

Definition 9.1. A metric space (X, d) is called *semialgebraic* if there exists a real closed field R , an integer n , a closed connected semialgebraic set $S \subset R^n$ such that (X, d) equals the completion of (S, d_S) .

Proof of Theorem 1.5. We first construct a semialgebraic set S and show afterwards that it is the Gromov-Hausdorff limit of the family. Let $R' := R(t)_{alg}^\wedge$ denote the (real closed) field of algebraic Puiseux-series in the parameter t . Equivalently, R' is the real closed field associated to the point 0_+ in the real spectrum of $R[t]$.

An element $\gamma \in R'$ can be identified with the germ at 0_+ of a continuous semialgebraic curve $\gamma : (0, \epsilon) \rightarrow R$ ($\epsilon \in R$).

Let $S := X_{0^+} \subset (R')^n$ be the fiber of X above 0^+ . S consists of those semialgebraic curve germs with $\gamma(t) \in X_t$ for all sufficiently small $t > 0$. We will show that (\bar{S}, \bar{d}_S) is the Gromov-Hausdorff limit of the family X .

Lemma 9.1. *Let $X \subset R \times R^n$ be a semialgebraic family which is closed, rationally bounded (i.e. there exists a natural number a with $X \subset B_a(0)$) and fiberwise semialgebraically connected. Then, for any rational number $C > 1$, there exists a decomposition $X = \cup_{i=1}^m X^i$ such that each X_t^i is connected and such that for $x, y \in X_t^i$*

$$\lambda_R(\|x - y\|) \leq d_{X_t^i}(x, y) \leq C\lambda_R(\|x - y\|)$$

and such that for $\gamma_1, \gamma_2 \in X_{0^+}^i$

$$\lambda_{R'}(\|\gamma_1 - \gamma_2\|) \leq d_{S^i}(\gamma_1, \gamma_2) \leq C\lambda_{R'}(\|\gamma_1 - \gamma_2\|).$$

The proof of the lemma is by extending the proof contained in [13] (which is based on [12]) to arbitrary real closed field. In most parts of the proof, one can just replace R by \mathbb{R} . This is not the case for the compactness of the Grassmannians used in [12], but this can be easily replaced by model completeness.

We continue the proof of the theorem. Let $\epsilon > 0$ be a rational number. Apply the above Lemma (with $C := 2$) to X . Let $x_1, \dots, x_k \in X_t^i$ be an ϵ -separated net (with respect to $d_{X_t^i}$). Then, if $j_1 \neq j_2$, $\lambda_R(\|x_{j_1} - x_{j_2}\|) \geq \frac{\epsilon}{2}$ which implies that $\|x_{j_1} - x_{j_2}\| \geq \frac{\epsilon}{4}$.

The size of an $\frac{\epsilon}{4}$ -separated net in $B(0, a)$ is bounded by a function of ϵ and a . This is trivial if $R = \mathbb{R}$ (by considering the volume) and follows by model completeness for all real closed R .

Therefore, the size of an ϵ -separated net in X_t is bounded by a number which only depends on ϵ, a, m , but not on t . It follows that the family of pseudo-metric spaces (X_t, d_{X_t}) is equicontact. In particular, each (X_t, d_{X_t}) is totally bounded, which implies (by the theorem of Hausdorff) that the completion $(\bar{X}_t, \bar{d}_{X_t})$ is compact.

By the same reasoning, the space (S, d_S) is totally bounded and its completion (\bar{S}, \bar{d}_S) compact.

Choose a rational $C > 1$ and a decomposition $X = \cup_{i=1}^m X^i$ as in the lemma. Similarly as in [13], we define a semialgebraic function $\tilde{d} : X \times X \rightarrow R$ by $\tilde{d}(x, y) := 0$ if x and y lie in different fibers and

$$\tilde{d}(x, y) := \inf \left\{ \sum_{i=0}^{m'-1} \|x_i - x_{i+1}\| : x = x_0, x_1, \dots, x_{m'} = y \text{ is a chain, } m' \leq m \right\}$$

Here the word “chain” means that two consecutive of the x_i lie in the closure of one of the X_t^i (where t is fixed). It is clear from the definition that \tilde{d} is bounded by the natural number $2ma$.

By the lemma, we get for $x, y \in X_t$

$$\lambda_R(\tilde{d}(x, y)) \leq d_{X_t}(x, y) \leq C\lambda_R(\tilde{d}(x, y))$$

and for $\gamma_1, \gamma_2 \in S = X_{0^+}$

$$\lambda_{R'}(\tilde{d}(\gamma_1, \gamma_2)) \leq d_S(\gamma_1, \gamma_2) \leq C\lambda_{R'}(\tilde{d}(\gamma_1, \gamma_2)).$$

Let γ_1, γ_2 be two points in X_{0^+} . Using the fact that the limit $\lim_{t \rightarrow 0^+} \tilde{d}(\gamma_1(t), \gamma_2(t)) \in R$ exists (since \tilde{d} is bounded by $2ma$ and semialgebraic and γ_1, γ_2 are semialgebraic) and using the alternative description of $\lambda_R, \lambda_{R'}$, we obtain

$$\lim_{t \rightarrow 0^+} \lambda_R(\tilde{d}(\gamma_1(t), \gamma_2(t))) = \lambda_{R'}(\tilde{d}(\gamma_1, \gamma_2)).$$

From this we conclude

$$\frac{1}{C} \limsup_{t \rightarrow 0^+} d_{X_t}(\gamma_1(t), \gamma_2(t)) \leq d_S(\gamma_1, \gamma_2) \leq C \liminf_{t \rightarrow 0^+} d_{X_t}(\gamma_1(t), \gamma_2(t)).$$

Since C was an arbitrary rational number with $C > 1$, it follows that

$$d_S(\gamma_1, \gamma_2) = \lim_{t \rightarrow 0^+} d_{X_t}(\gamma_1(t), \gamma_2(t)).$$

Now we continue the proof as in Section 3 (replacing “subanalytic” by “semialgebraic”) to see that (\bar{S}, \bar{d}_S) equals the Gromov-Hausdorff limit $\lim_{t \rightarrow 0^+} (\bar{X}_t, \bar{d}_{X_t})$. \square

Remark 9.1. The above proof, applied to a constant family, shows that the metric space associated to a rationally bounded, connected, closed semialgebraic set is compact. Since the Gromov-Hausdorff distance between two compact metric spaces vanishes if and only if they are isometric, the metric space associated to an extension of a semialgebraic S to another real closed field ([5]) gives rise to the same metric space.

Remark 9.2. The Hausdorff limit of the family X at 0^+ is given by $\lambda_R(S)$, endowed with the Euclidean metric ([7]). This shows that tangent cone and semialgebraic tangent cone at a point x of a closed semialgebraic set are given by the same semialgebraic set, but the former gets the length metric and the latter gets the Euclidean metric.

REFERENCES

- [1] *A. Bernig*, Scalar Curvature of definable Alexandrov spaces. *Advances in Geometry* **2** (2002), 29-55.
- [2] *A. Bernig*, Scalar curvature of definable CAT-spaces. *Advances in Geometry* **3** (2003), 23-43.
- [3] *E. Bierstone, P. D. Milman*, Semianalytic and subanalytic sets. *Inst. Hautes Études Sci. Publ. Math.* **67** (1988), 5-42.
- [4] *L. Birbrair, T. Mostowski*, Normal embeddings of semialgebraic sets. *Michigan Math. J.* **47** (2000), 125-132.
- [5] *J. Bochnak, M. Coste, M.-F. Roy*, Géométrie algébrique réelle. Springer Verlag Berlin, 1987.
- [6] *M. Bridson, A. Haefliger*, Metric Spaces of Non-Positive Curvature. Springer-Verlag Berlin-Heidelberg 1999.

- [7] *L. Bröcker*, Families of semialgebraic sets and limits. In: Real Algebraic Geometry, LNM 1524, Springer-Verlag Berlin 1992, 145-162.
- [8] *L. Bröcker, M. Kuppe, W. Scheufler*, Inner metric properties of 2-dimensional semi-algebraic sets. Rev. Mat. Univ. Complut. Madrid **10** (1997), 51-78.
- [9] *D. Burago, Yu. Burago, S. Ivanov*, A course in metric geometry. American Mathematical Society, Providence, RI 2001.
- [10] *M. Coste, M. Reguiat*, Trivialités en famille. In: Real Algebraic Geometry, LNM 1524, Springer-Verlag Berlin 1992, 193-204.
- [11] *R. Hardt*, Some analytic bounds for subanalytic sets. In: Differential geometric control theory, Progress in Math., vol 27, 1983, 259-267.
- [12] *K. Kurdyka*, On a subanalytic stratification satisfying a Whitney property with exponent 1. In: Real Algebraic Geometry, LNM 1524, Springer-Verlag Berlin 1992, 316-322.
- [13] *K. Kurdyka, P. Orro*, Distance géodésique sur un sous-analytique. Rev. Mat. Univ. Complut. Madrid **10** (1997), 173-182.
- [14] *K. Kurdyka, G. Raby*, Densité des ensembles sous-analytiques. Ann. Inst. Fourier **39** (1989), 753-771.
- [15] *A. Lytchak*, Differentiation in metric space. Preprint 2003.
- [16] *P. Orro*, Quelques propriétés de la distance géodésique. In: Real analytic and algebraic singularities, Pitman Res. Notes Math. Ser. 381, Longman, Harlow, 1998, 107-113.
- [17] *A. Parusiński*, Lipschitz stratification of subanalytic sets. Ann. scient. Ec. Norm. Sup. **27** (1994), 661-696.
- [18] *M. Pflaum*, Analytic and Geometric Study of Stratified Spaces. Springer-Verlag Berlin-Heidelberg 2001.
- [19] *L. Van den Dries, C. Miller*, Geometric Categories and o-minimal structures. Duke Math. Journal **84** (1996), 497-540.

MATHEMATISCHES INSTITUT, UNIVERSITÄT FREIBURG, ECKERSTR. 1, 79104 FREIBURG, GERMANY

E-mail address: `andreas.bernig@math.uni-freiburg.de`

MATHEMATISCHES INSTITUT, UNIVERSITÄT BONN, BERINGSTR. 1, 53115 BONN, GERMANY,

E-mail address: `lytchak@math.uni-bonn.de`