

# Emergence of the 16S rRNA Methylase RmtG in an Extended-Spectrum- $\beta$ -Lactamase-Producing and Colistin-Resistant *Klebsiella pneumoniae* Isolate in Chile

Laurent Poirel,<sup>a,b</sup> Jaime Labarca,<sup>b,c</sup> Helia Bello,<sup>d</sup> Maria Luisa Riosco,<sup>e</sup> Sandrine Bernabeu,<sup>b</sup> Patrice Nordmann<sup>a,b</sup>

Medical and Molecular Microbiology Unit, Department of Medicine, Faculty of Science, University of Fribourg, Fribourg, Switzerland<sup>a</sup>; INSERM U914, Emerging Resistance to Antibiotics, Le Kremlin-Bicêtre, France<sup>b</sup>; Department of Infectious Diseases, School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile<sup>c</sup>; Department of Microbiology, Facultad de Ciencias Biológicas, Universidad de Concepción, Concepción, Chile<sup>d</sup>; Clinical Microbiology Laboratory, Hospital de Puerto Montt, Puerto Montt, Chile<sup>e</sup>

While carbapenemases are disseminating worldwide in *Enterobacteriaceae*, and particularly in *Escherichia* and *Klebsiella pneumoniae* (1), therapeutic options increasingly rely on very few choices (2). Although colistin and tigecycline are now considered last-resort antibiotics for treating infections caused by carbapenemase producers, aminoglycosides may still be considered valuable treatment options, with amikacin remaining quite often active (3). While broad-spectrum  $\beta$ -lactamases such as extended-spectrum  $\beta$ -lactamases (ESBLs) or carbapenemases are spreading rapidly worldwide, the emergence of acquired 16S rRNA methylases has been also noticed (4). Those enzymes do confer high-level resistance to almost all aminoglycosides. The most frequent 16S rRNA methylases in *Enterobacteriaceae* are ArmA, RmtB to RmtF, and NpmA (4, 5). Those enzymes exhibit significant amino acid heterogeneity, sharing similar amino acid motifs but being overall distantly related. Nevertheless, they all confer the same type of resistance, whereas NpmA confers additional resistance to neomycin and apramycin (5).

Here, we report on a 22-year-old male patient who was admitted at the intensive care unit of the Puerto Montt Hospital located in Santiago, Chile, on 30 June 2012 for a traumatic subdural hematoma of the occipitotemporal region. He underwent surgery and was mechanically ventilated. Ceftriaxone and metronidazole were prescribed for 7 days (during which time cultures remained negative) as empirical treatment for pneumonia. Ten days after his admission, bronchial aspirate samples grew ceftriaxone-resistant *K. pneumoniae*, while an isolate with the same resistant pattern was recovered from his central venous catheter 3 days later. Therefore, ceftriaxone treatment was discontinued and switched to meropenem, and the patient was finally discharged after a good clinical response. Susceptibility testing performed and interpreted according to the updated CLSI guidelines (6) showed that it was resistant at a high level to all aminoglycosides (MICs of  $>256$   $\mu\text{g/ml}$  for amikacin, gentamicin, and tobramycin) and to most broad-spectrum  $\beta$ -lactams (MIC of cefotaxime and cefepime at 256  $\mu\text{g/ml}$ , MIC of ceftazidime at 8  $\mu\text{g/ml}$ ). It remained susceptible to carbapenems (MICs of imipenem and meropenem at 0.5 and 1  $\mu\text{g/ml}$ , respectively), and the Carba NP test remained negative (7). In addition, isolate 921 was resistant to tetracycline, chloramphenicol, sulfonamides, and to all tested fluoroquinolones. It remained susceptible to tigecycline (MIC of 1  $\mu\text{g/ml}$ ) according to EUCAST breakpoints (8), but the MIC of colistin was high as determined by Etest (4  $\mu\text{g/ml}$ ). According to the phenotypic test results (synergy between aztreonam and clavulanate), *K. pneumoniae* isolate 921 produced an ESBL.

PCR and sequencing performed as described previously (9)

revealed that *K. pneumoniae* isolate 921 harbored the *bla*<sub>CTX-M-2</sub> ESBL gene, in addition to the *bla*<sub>TEM-1</sub> and *bla*<sub>SHV-11</sub> narrow-spectrum  $\beta$ -lactamase genes. Since isolate 921 was resistant at a high level to all aminoglycosides, screening for 16S rRNA methylase genes was performed as reported (10). It failed to identify any known 16S rRNA methylase-encoding gene. Therefore, a shotgun cloning experiment was attempted as described previously (11), and selection of recombinant *Escherichia coli* strains was based on amikacin (30  $\mu\text{g/ml}$ ) and gentamicin (30  $\mu\text{g/ml}$ ). Recombinant strains expressing resistance to all aminoglycosides were obtained, suggesting the expression of a 16S rRNA methylase. Sequencing of the recombinant plasmid identified a putative open reading frame encoding a protein sharing 58% amino acid identity with the 16S rRNA methylase RmtD. However, it has a perfect identity with the newly described RmtG which was published while this study was in progress (12). The *rmtG* gene encoding high-level resistance to all aminoglycosides has been identified in a series of five *K. pneumoniae* isolates from Brazil, four coproducing the carbapenemase KPC-2, and all coproducing an ESBL of the CTX-M type (CTX-M-2, CTX-M-15, or CTX-M-59) (12).

Mating-out assays were performed as described previously (11) but remained unsuccessful. Therefore, an electrotransformation experiment was performed using a selection based on amikacin and gentamicin or on cefotaxime (30  $\mu\text{g/ml}$ ). It allowed us to obtain an *E. coli* transformant expressing RmtG, exhibiting resistance to all aminoglycosides, to sulfonamides, and to tetracycline but remaining susceptible to all  $\beta$ -lactams, and another type of *E. coli* transformant expressing CTX-M-2 and TEM-1. The *rmtG*-positive transformant harbored a single 80-kb plasmid that was of IncA/C type as identified by PCR-based replicon typing (13). The *bla*<sub>CTX-M-2</sub>-positive transformant harbored a single 170-kb plasmid that was not typeable.

Multilocus sequence typing was performed as described previously (14), and results were analyzed by eBURST (<http://pubmlst.org>). It showed that isolate 921 belonged to the ST11 sequence type, whereas the sequence types of the RmtG-positive *K. pneumoniae* isolates from Brazil were 340, 442, and 1046 (12). However, in the same study, Bueno et al. identified several ST11 isolates expressing the RmtD2 methylase (12).

Address correspondence to Patrice Nordmann, [patrice.nordmann@unifr.ch](mailto:patrice.nordmann@unifr.ch).

Of note, the patient did not receive any aminoglycoside-based treatment before or after his admission. Therefore, selection of this aminoglycoside-resistant isolate remains intriguing. It may be speculated that clones or plasmids carrying the *rmtG* gene are disseminating in South American countries such as Chile and Brazil (12). Occurrence of multidrug-resistant strains is extremely worrisome, and there is an urgent need to implement adequate screening strategies in order to control their spread. This study underlines that a focus on emerging antibiotic resistance mechanisms in Gram negatives shall be made not only for broad-spectrum  $\beta$ -lactams but also for broad-spectrum aminoglycosides, since both types of resistance will contribute to the emergence of pan-drug resistance.

## ACKNOWLEDGMENTS

This work was funded mostly by the INSERM, France, by the University of Fribourg, Switzerland, and by grants from the European Community (RGNOSIS, FP7/HEALTH-F3-2011-282512, and MAGIC-BULLET, FP7/HEALTH-F3-2001-278232).

## REFERENCES

1. Nordmann P, Naas T, Poirel L. 2011. Global spread of carbapenemase-producing *Enterobacteriaceae*. *Emerg. Infect. Dis.* 17:1791–1798. <http://dx.doi.org/10.3201/eid1710.110655>.
2. Nordmann P, Poirel L, Toleman MA, Walsh TR. 2011. Does broad spectrum  $\beta$ -lactam resistance due to NDM-1 herald the end of the antibiotic era for treatment of infections caused by Gram-negative bacteria? *J. Antimicrob. Chemother.* 66:689–692. <http://dx.doi.org/10.1093/jac/dkq520>.
3. Falagas ME, Karageorgopoulos DE, Nordmann P. 2011. Therapeutic options for infections with *Enterobacteriaceae* producing carbapenem-hydrolyzing enzymes. *Future Microbiol.* 6:653–666. <http://dx.doi.org/10.2217/fmb.11.49>.
4. Doi Y, Arakawa Y. 2007. 16S ribosomal RNA methylation: emerging resistance mechanism against aminoglycosides. *Clin. Infect. Dis.* 45:88–94. <http://dx.doi.org/10.1086/518605>.
5. Wachino J, Shibayama K, Kurokawa H, Kimura K, Yamane K, Suzuki S, Shibata N, Ike Y, Arakawa Y. 2007. Novel plasmid-mediated 16S rRNA m<sup>1</sup>A1408 methyltransferase, NpmA, found in a clinically isolated *Escherichia coli* strain resistant to structurally diverse aminoglycosides. *Antimicrob. Agents Chemother.* 51:4401–4409. <http://dx.doi.org/10.1128/AAC.00926-07>.
6. Clinical and Laboratory Standards Institute. 2013. Performance standards for antimicrobial susceptibility testing; 23rd informational supplement. M100-S23. CLSI, Wayne, PA.
7. Nordmann P, Poirel L, Dortet L. 2012. Rapid detection of carbapenemase-producing *Enterobacteriaceae*. *Emerg. Infect. Dis.* 18:1503–1507. <http://dx.doi.org/10.3201/eid1809.120355>.
8. EUCAST technical note on tigecycline. 2006. *Clin. Microb. Infect.* 12: 1147–1149. <http://dx.doi.org/10.1111/j.1469-0691.2006.01578.x>.
9. Poirel L, Dortet L, Bernabeu S, Nordmann P. 2011. Genetic features of bla<sub>NDM-1</sub>-positive *Enterobacteriaceae*. *Antimicrob. Agents Chemother.* 55: 5403–5407. <http://dx.doi.org/10.1128/AAC.00585-11>.
10. Berçot B, Poirel L, Nordmann P. 2011. Updated multiplex PCR for detection of 16S rRNA methylases; high prevalence among NDM-1 producers. *Diagn. Microbiol. Infect. Dis.* 71:442–445. <http://dx.doi.org/10.1016/j.diagmicrobio.2011.08.016>.
11. Doi Y, Yokoyama K, Yamane K, Wachino J, Shibata N, Yagi T, Shibayama K, Kato H, Arakawa Y. 2004. Plasmid-mediated 16S rRNA methylase in *Serratia marcescens* conferring high-level resistance to aminoglycosides. *Antimicrob. Agents Chemother.* 48:491–496. <http://dx.doi.org/10.1128/AAC.48.2.491-496.2004>.
12. Bueno MF, Francisco GR, O'Hara JA, de Oliveira Garcia D, Doi Y. 2013. Coproduction of 16S rRNA methyltransferase RmtD or RmtG with KPC-2 and CTX-M group extended-spectrum  $\beta$ -lactamases in *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.* 57:2397–2400. <http://dx.doi.org/10.1128/AAC.02108-12>.
13. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. 2005. Identification of plasmids by PCR-based replicon typing. *J. Microbiol. Methods* 63:219–228. <http://dx.doi.org/10.1016/j.mimet.2005.03.018>.
14. Diancourt L, Passet V, Verhoef J, Grimont PA, Brisse S. 2005. Multilocus sequence typing of *Klebsiella pneumoniae* nosocomial isolates. *J. Clin. Microbiol.* 43:4178–4182. <http://dx.doi.org/10.1128/JCM.43.8.4178-4182.2005>.