

Plateau's problem for integral currents in locally non-compact metric spaces

Stefan Wenger

Abstract. The purpose of this article is to prove existence of mass minimizing integral currents with prescribed possibly non-compact boundary in all dual Banach spaces and furthermore in certain spaces without linear structure, such as injective metric spaces and Hadamard spaces. We furthermore prove a weak*-compactness theorem for integral currents in dual spaces of separable Banach spaces. Our theorems generalize results of Ambrosio–Kirchheim, Lang, the author, and recent results of Ambrosio–Schmidt.

Keywords. Plateau's problem, area minimizers, Banach spaces, compactness.

2010 Mathematics Subject Classification. Primary 49Q15, 49Q05, 46B50; secondary 53C23.

1 Introduction

In this paper we study the generalized Plateau problem in the context of locally non-compact metric spaces. Roughly speaking, this problem concerns the question of existence of an $(m + 1)$ -dimensional generalized surface of least volume with prescribed m -dimensional boundary in a given metric space. A suitable notion of surface in the context of area minimization problems is provided by the theory of integral currents. In the setting of Euclidean space, this theory was developed by Federer–Fleming in [4], who solved Plateau's problem in the class of integral currents in Euclidean space. In [1], Ambrosio–Kirchheim extended Federer–Fleming's theory to the setting of complete metric spaces. They then solved the generalized Plateau problem in the class of integral currents in compact metric spaces (provided that given a boundary the family of fillings is not empty) and furthermore in dual spaces of separable Banach spaces, provided that the prescribed boundary lies in a compact set. This result was generalized in [8] to all dual Banach spaces as well as to Hadamard spaces (for Hadamard spaces, the result is due to U. Lang), still requiring that the prescribed boundary remain in a compact set. By definition, Hadamard spaces are complete simply-connected metric spaces of

Partially supported by NSF grants DMS 0956374 and DMS 1056263.

non-positive curvature in the sense of Alexandrov, see e.g. [3, Section II.1]. It was only shown very recently by Ambrosio–Schmidt [2] that the condition on compact boundary can be dropped in the case of separable dual Banach spaces. The aim of the present paper is to remove the separability condition made in [2]. In fact, we will show that the generalized Plateau problem can be solved for non-compact boundaries in a class of metric spaces which includes e.g. all dual Banach spaces (also non-separable ones), all injective metric spaces (i.e. absolute 1-Lipschitz retracts), and all Hadamard spaces. We therefore generalize corresponding results in [1, 2, 8]. We furthermore partially generalize weak*-compactness theorems proved in [1, 2] to the setting of dual spaces of separable Banach spaces.

We now give precise formulations of our main results. Given a complete metric space X and $m \geq 0$, we will denote by $\mathbf{M}_m(X)$ and $\mathbf{I}_m(X)$ the spaces of metric m -currents of finite mass and of integral m -currents, respectively, in the sense of Ambrosio–Kirchheim [1]. Given $T \in \mathbf{M}_m(X)$, the mass of T will be denoted by $\mathbf{M}(T)$ and, in case $m \geq 1$, the boundary of T by ∂T . We refer to Section 2 for the basic definitions from the theory of metric currents. Our first theorem gives solutions to Plateau’s problem and to the corresponding free boundary problem in the context of Banach spaces.

Theorem 1.1. *Let X be a Banach space which is 1-complemented in a dual Banach space. Let $m \geq 0$. Then*

- (i) *for every $S \in \mathbf{I}_m(X)$ with $\partial S = 0$ there exists $T \in \mathbf{I}_{m+1}(X)$ with $\partial T = S$ and such that*

$$\mathbf{M}(T) \leq \mathbf{M}(T')$$

for all $T' \in \mathbf{I}_{m+1}(X)$ with $\partial T' = S$,

- (ii) *for every $S \in \mathbf{M}_m(X)$ there exists $T \in \mathbf{I}_{m+1}(X)$ such that*

$$\mathbf{M}(T) + \mathbf{M}(\partial T - S) \leq \mathbf{M}(T') + \mathbf{M}(\partial T' - S)$$

for all $T' \in \mathbf{I}_{m+1}(X)$.

If $m = 0$, then the condition $\partial S = 0$ in (i) should be replaced by $S(1) = 0$, see Section 2. In the above, a Banach space X is said to be 1-complemented in a dual Banach space if X is (isometric to) a subspace W of a dual Banach space Y such that there is a norm 1 projection of Y onto W . Particular examples of such spaces are dual Banach spaces, L^1 -spaces, and L -embedded Banach spaces ([5]). Note that no compactness assumption on $\text{spt } S$ or separability assumption on X is made, and that furthermore S in (ii) is only required to be a current of finite mass. Theorem 1.1 generalizes [1, Theorem 10.6], [8, Theorem 1.5], and [2, Theorems 1.1 and 1.3].

Our next result provides a weak*-compactness theorem which partially generalizes corresponding results in [1] and [2].

Theorem 1.2. *Let X be either a reflexive Banach space or the dual space of a separable Banach space. Let $m \geq 0$ and let $(T_n) \subset \mathbf{I}_m(X)$ be a sequence satisfying*

$$\sup_n [\mathbf{M}(T_n) + \mathbf{M}(\partial T_n)] < \infty \quad (1.1)$$

and

$$\lim_{r \rightarrow \infty} \left[\sup_n \|T_n\|(X \setminus B(0, r)) \right] = 0.$$

Then there exists a subsequence T_{n_j} which w^ -converges to some $T \in \mathbf{I}_m(X)$.*

Here, $B(0, r)$ denotes the (open) ball in X of radius r and center 0. If $m = 0$, then (1.1) should be replaced by $\sup_n \mathbf{M}(T_n) < \infty$. For the definition of w^* -convergence see Definition 2.2. We remark that Theorem 1.2 fails in general if X is the dual space of a non-separable Banach space, see Example 3.4. Our theorem generalizes [1, Theorem 6.6] and [2, Theorem 1.4] in the case of integral currents. While our theorem applies to a larger class of Banach spaces, [1, Theorem 6.6] and [2, Theorem 1.4] apply to normal currents as well. It is interesting to note however that Theorem 1.2 cannot be generalized to normal currents without the additional assumption that X be separable, as was shown in [2, Example B.1].

We now briefly turn to metric spaces without a vector space structure. The methods which we use are not restricted to the setting of Banach spaces. Indeed, in Theorem 3.3 we will generalize Theorem 1.1 to a class of metric spaces. We will, in particular, obtain the following result as a consequence.

Theorem 1.3. *Let X be a Hadamard space or an injective metric space and let $m \geq 0$. Then assertions (i) and (ii) of Theorem 1.1 hold for X .*

Theorem 1.3 generalizes a corresponding result for compact boundaries in Hadamard spaces which goes back to U. Lang and which was published in [8, Theorem 1.6].

Finally, we mention that our approach differs from the one taken in [1] and [2]. Our methods combine arguments in the spirit of [8] with a variant of a compactness theorem recently proved in [10] and [7].

The paper is structured as follows. In Section 2 we recall the definitions from the theory of currents needed for the paper. We furthermore define the notion of local weak convergence and establish a relationship with weak convergence. In Section 3 we prove a variant of the compactness theorems established in [7, 10] and apply it in the proofs of the theorems stated above. We will furthermore state and prove some generalizations of the above results, see Theorem 3.3 and Theorem 3.5.

2 Preliminaries

In this section we recall the basic definitions from the theory of metric currents developed in [1] which we will need in the sequel. With the exception of the definition of local weak convergence, its relationship with weak convergence, and Lemma 2.3, all definitions and results appear in [1].

Let (X, d) be a complete metric space. We denote by $\text{Lip}(X)$ and $\text{Lip}_b(X)$ the spaces of real-valued Lipschitz functions and bounded Lipschitz functions on X , respectively. The Lipschitz constant of a Lipschitz function f will be denoted by $\text{Lip}(f)$.

Definition 2.1. Let $m \geq 0$. An m -dimensional metric current T on X is a multilinear functional $T : \text{Lip}_b(X) \times \text{Lip}^m(X) \rightarrow \mathbb{R}$ satisfying the following properties:

- (i) If $\pi_i^j \rightarrow \pi_i$ pointwise as $j \rightarrow \infty$ and if $\sup_{i,j} \text{Lip}(\pi_i^j) < \infty$, then

$$T(f, \pi_1^j, \dots, \pi_m^j) \rightarrow T(f, \pi_1, \dots, \pi_m).$$

- (ii) If $\{x \in X : f(x) \neq 0\}$ is contained in the union $\bigcup_{i=1}^m B_i$ of Borel sets B_i and if π_i is constant on B_i , then

$$T(f, \pi_1, \dots, \pi_m) = 0.$$

- (iii) There exists a finite Borel measure μ on X such that

$$|T(f, \pi_1, \dots, \pi_m)| \leq \prod_{i=1}^m \text{Lip}(\pi_i) \int_X |f| d\mu \quad (2.1)$$

for all $(f, \pi_1, \dots, \pi_k) \in \text{Lip}_b(X) \times \text{Lip}^m(X)$.

The space of m -dimensional metric currents on X is denoted by $\mathbf{M}_m(X)$ and the minimal Borel measure μ satisfying (2.1) is called mass of T and denoted by $\|T\|$. We also call mass of T the number $\|T\|(X)$ which we denote by $\mathbf{M}(T)$. The support of T is the closed set

$$\text{spt } T = \{x \in X : \|T\|(B(x, r)) > 0 \text{ for all } r > 0\}.$$

As in [1] we will assume throughout this paper that the cardinality of any set is an Ulam number. This is consistent with the standard ZFC set theory. We then have that $\text{spt } T$ is separable and furthermore that $\|T\|$ is concentrated on a σ -compact set, i.e. $\|T\|(X \setminus C) = 0$ for a σ -compact set $C \subset X$ (see [1]). This will be relevant in the proof of statement (ii) of Theorem 1.1.

In the following we will often abbreviate $\pi = (\pi_1, \dots, \pi_m)$ and write $T(f, \pi)$ instead of $T(f, \pi_1, \dots, \pi_m)$. The restriction of $T \in \mathbf{M}_m(X)$ to a Borel set $A \subset X$ is given by

$$(T \llcorner A)(f, \pi) := T(f\chi_A, \pi).$$

This expression is well-defined since T can be extended to a functional on tuples for which the first argument lies in $L^\infty(X, \|T\|)$. Furthermore, $T \llcorner A \in \mathbf{M}_m(X)$ by [1, Theorem 3.5]. If $m \geq 1$ and $T \in \mathbf{M}_m(X)$, then the boundary of T is the functional

$$\partial T(f, \pi_1, \dots, \pi_{m-1}) := T(1, f, \pi_1, \dots, \pi_{m-1});$$

it satisfies conditions (i) and (ii) in Definition 2.1. If it moreover satisfies condition (iii) in Definition 2.1, then T is called a normal current. By convention, elements of $\mathbf{M}_0(X)$ are also called normal currents. If $m = 0$ and $T \in \mathbf{M}_0(X)$, then we say $\partial T = 0$ if $T(1) = 0$. The push-forward of $T \in \mathbf{M}_m(X)$ under a Lipschitz map φ from X to another complete metric space Y is given by

$$\varphi_\# T(g, \tau) := T(g \circ \varphi, \tau \circ \varphi)$$

for $(g, \tau) \in \text{Lip}_b(Y) \times \text{Lip}^m(Y)$. This defines an m -dimensional current on Y . It follows directly from the definitions that $\partial(\varphi_\# T) = \varphi_\#(\partial T)$.

Definition 2.2. A sequence of currents $T_n \in \mathbf{M}_m(X)$ is said to converge weakly to $T \in \mathbf{M}_m(X)$ if

$$T_n(f, \pi) \rightarrow T(f, \pi) \tag{2.2}$$

for every $(f, \pi) \in \text{Lip}_b(X) \times \text{Lip}^m(X)$. If (2.2) only holds for those (f, π) for which f also has bounded support, we say that T_n converges locally weakly to T . If X is a dual Banach space and (2.2) holds for those (f, π) for which the f and π_i are also weak*-continuous, we say that T_n w*-converges to T .

It is clear that if T_n converges locally weakly to T and

$$\lim_{r \rightarrow \infty} \left[\sup_n \|T_n\|(X \setminus B(x_0, r)) \right] = 0$$

for some $x_0 \in X$, then T_n converges weakly to T . We furthermore have the following easy lemma.

Lemma 2.3. Let X, Y be complete metric spaces, $m \geq 0$, and $S \in \mathbf{M}_m(X)$. Let (φ_j) be a sequence of Lipschitz maps $\varphi_j : \text{spt } S \rightarrow Y$ with $\sup_j \text{Lip}(\varphi_j) < \infty$. If φ_j converges pointwise to a Lipschitz map $\varphi : \text{spt } S \rightarrow Y$, then $\varphi_{j\#} S$ converges weakly to $\varphi_\# S$.

Proof. Let $(f, \pi) \in \text{Lip}_b(Y) \times \text{Lip}^m(Y)$. We may assume without loss of generality that $|f| \leq 1$ and that each π_i is 1-Lipschitz. Set $A := \sup_j \text{Lip}(\varphi_j)$ and let $\varepsilon > 0$. By σ -compactness of $\text{spt } S$ there exists $C \subset \text{spt } S$ compact such that $\|S\|(X \setminus C) \leq \varepsilon/(4A^m)$. We then obtain for each j that

$$\begin{aligned} |\varphi_{\#}S(f, \pi) - \varphi_{j\#}S(f, \pi)| &\leq |(S \llcorner C)(f \circ \varphi, \pi \circ \varphi) - (S \llcorner C)(f \circ \varphi, \pi \circ \varphi_j)| \\ &\quad + |(S \llcorner C)(f \circ \varphi - f \circ \varphi_j, \pi \circ \varphi_j)| \\ &\quad + 2A^m \|S\|(X \setminus C) \\ &\leq |(S \llcorner C)(f \circ \varphi, \pi \circ \varphi) - (S \llcorner C)(f \circ \varphi, \pi \circ \varphi_j)| \\ &\quad + A^m \int_C |f \circ \varphi - f \circ \varphi_j| d\|S\| + \frac{\varepsilon}{2}. \end{aligned}$$

The first term after the last inequality sign converges to 0 as $j \rightarrow \infty$ by the continuity property of currents, property (i) in Definition 2.1. The second term converges to 0 since $f \circ \varphi_j$ converges uniformly to $f \circ \varphi$ on the compact set C . Therefore, the sum of the three terms is $\leq \varepsilon$ for all j large enough. This shows that $\varphi_{j\#}S$ converges weakly to $\varphi_{\#}S$. \square

In this paper we will mainly be concerned with integral currents. An element $T \in \mathbf{M}_0(X)$ is called integer rectifiable if there exist finitely many points

$$x_1, \dots, x_n \in X \quad \text{and} \quad \theta_1, \dots, \theta_n \in \mathbb{Z} \setminus \{0\}$$

such that

$$T(f) = \sum_{i=1}^n \theta_i f(x_i)$$

for all bounded Lipschitz functions f . A current $T \in \mathbf{M}_m(X)$ with $m \geq 1$ is said to be integer rectifiable if the following properties hold:

- (i) $\|T\|$ is concentrated on a countably \mathcal{H}^m -rectifiable set and vanishes on all \mathcal{H}^m -negligible Borel sets, where \mathcal{H}^m denotes the Hausdorff m -measure.
- (ii) For any Lipschitz map $\varphi : X \rightarrow \mathbb{R}^m$ and any open set $U \subset X$ there exists $\theta \in L^1(\mathbb{R}^m, \mathbb{Z})$ such that $\varphi_{\#}(T \llcorner U) = \llbracket \theta \rrbracket$, where

$$\llbracket \theta \rrbracket(f, \pi) := \int_{\mathbb{R}^m} \theta f \det(\nabla \pi) d\mathcal{H}^m.$$

Integer rectifiable normal currents are called integral currents. The corresponding space is denoted by $\mathbf{I}_m(X)$.

Recently, variants of Ambrosio-Kirchheim's theory that do not rely on the finite mass axiom have been developed by Lang in [6] and by Lang and the author in [7]. We will not need any definitions or results from these theories, however.

3 Proofs of the main results

The principal tool in the proofs of our main theorems is the following compactness result, which is a variant of the compactness theorem proven by the author in [10] and of a generalization proven by Lang and the author in [7].

Lemma 3.1. *Let X be a complete metric space, $\Omega \subset X$ separable, and $m \geq 0$. Let $(T_n) \subset \mathbf{I}_m(X)$ be a sequence satisfying*

$$\sup_n [\mathbf{M}(T_n) + \mathbf{M}(\partial T_n)] < \infty.$$

Then there exist a subsequence (n_j) , a complete metric space Z , and isometric embeddings $\varphi_j : X \hookrightarrow Z$ such that $\varphi_{j\#}T_{n_j}$ converges locally weakly to some element $T \in \mathbf{I}_m(Z)$ and $\varphi_j|_{\Omega}$ converges pointwise to an isometric embedding $\varphi : \Omega \hookrightarrow Z$.

The statement of Lemma 3.1 can be strengthened to apply also to the locally integral currents of [7] and local weak convergence can be replaced by convergence in the local flat topology. However, the above version suffices for all the applications in this paper. Note that, on the one hand, the results in [7, 10] apply more generally to sequences (T_n) of integral currents in a sequence (X_n) of metric spaces. On the other hand, these results do not yield convergence of the isometric embeddings φ_j as Lemma 3.1 does.

The proof of Lemma 3.1 relies on the constructions used in [7, 10]. Roughly speaking, the idea is to decompose each current T_n into a sum $T_n = T_n^1 + T_n^2 + \dots$ of currents T_n^i with mass growth $\geq \gamma_i r^m$ for some $\gamma_i > 0$ independent of n . This mass growth implies that for fixed i the sequence $(\text{spt } T_n^i)$ of supports is a uniformly compact sequence of metric spaces in the sense that for every $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that every set $\text{spt } T_n^i$, $n = 1, 2, \dots$, can be covered by at most N balls of radius ε . One can then use a variant of Gromov's compactness theorem together with the closure and compactness theorems for integral currents in a compact metric space [1] to produce a desired metric space Z and a limit T as in Lemma 3.1. The decomposition procedure alluded above was proved in [10, Lemma 5.1, Theorem 1.2]. It was summarized in [7, Proposition 3.1] in a form which is suitable for our purposes.

In the proof below we will follow the proof of [7, Theorem 1.1]. We would like to emphasize, however, that even though the paper [7] deals with locally integral currents we will only need integral currents in the sense of [1] and that we could instead follow the arguments given in the proof of [10, Theorem 1.2].

Proof. Fix $x_0 \in \Omega$ and choose numbers $0 < R_1 < R_2 < \dots \rightarrow \infty$ such that, after passing to a subsequence, we have

$$T_n \llcorner B(x_0, R_r) \in \mathbf{I}_m(X)$$

with

$$\sup_n \mathbf{M}(\partial(T_n \llcorner B(x_0, R_r))) < \infty$$

for each $r \in \mathbb{N}$. Existence of such an R_r follows from [1, Theorem 5.6] together with Fatou's lemma. Set $R_0 := 0$, and define $A_r := B(x_0, R_r) \setminus B(x_0, R_{r-1})$ and

$$T_{r,n} := T_n \llcorner A_r$$

for $r \in \mathbb{N}$; clearly $T_{r,n} \in \mathbf{I}_m(X)$ and

$$\sup_n [\mathbf{M}(T_{r,n}) + \mathbf{M}(\partial T_{r,n})] < \infty.$$

Fix integers $1 \leq j_1 < j_2 < \dots$ and positive numbers $\frac{1}{2} > \delta_1 > \delta_2 > \dots$ with

$$\sum_i \delta_i < \infty.$$

By possibly replacing X by $\ell^\infty(X)$ we may assume that X admits isoperimetric inequalities of Euclidean type for integral currents, see [8, Corollary 1.3]. Now, let

$$T_{r,n} = T_{r,n}^1 + \dots + T_{r,n}^{j_n+1} + U_{r,n}^1 + \dots + U_{r,n}^{j_n+1}$$

be a decomposition with $T_{r,n}^1, \dots, T_{r,n}^{j_n+1}, U_{r,n}^1, \dots, U_{r,n}^{j_n+1} \in \mathbf{I}_m(X)$ as in [7, Proposition 3.1] for $T_{r,n}$, R_r , x_0 , and X . Let $\{y_0, y_1, y_2, \dots\} \subset \Omega$ be a countable dense subset of Ω with $y_0 = x_0$ and define $\Omega_s := \{y_0, \dots, y_s\}$ for $s \in \mathbb{N}$. For $n, s \in \mathbb{N}$, define closed sets

$$B_n^s := \Omega_s \cup \bigcup_{r=1}^s \bigcup_{i=1}^{\min\{s, j_n\}} (\text{spt } T_{r,n}^i \cup \text{spt } U_{r,n}^i)$$

and note that $B_n^1 \subset B_n^2 \subset \dots \subset X$. According to [7, Proposition 3.1 (i)], for each s , the sequence (B_n^s) is uniformly compact. By [10, Proposition 5.2], after passage to a subsequence, there exist isometric embeddings $\varphi_n : X \hookrightarrow Z$ and compact subsets $Y^1 \subset Y^2 \subset \dots \subset Z$, for some complete metric space Z , such that

$$\varphi_n(B_n^s) \subset Y^s$$

for all n and s . It can be shown exactly as in the proof of [10, Theorem 1.2] or the proof of [7, Theorem 1.1] that, after passing to a further subsequence, $\varphi_{n\#} T_n$ converges locally weakly to some $T \in \mathbf{I}_m(Z)$. Note that the proof in [7] shows that after possibly replacing Z by $\ell^\infty(Z)$ the $\varphi_{n\#} T_n$ converge to T even in the local flat topology (which implies local weak convergence).

We are left to prove that, after passing to a further subsequence, φ_n converges to an isometric embedding $\varphi : \Omega \hookrightarrow Z$. For this, note first that since $\Omega_s \subset B_n^s$ we obtain $\varphi_n(\Omega_s) \subset Y^s$ for all n and s . Since each Y^s is compact, we may assume,

after passing to a subsequence, that $\varphi_n(y_s)$ converges to some $z_s \in Y^s$ as $n \rightarrow \infty$. By density of $\{y_0, y_1, \dots\}$ in Ω and by the fact that $d(z_s, z_r) = d(y_s, y_r)$ for all r, s it follows that there exists an isometric embedding $\varphi : \Omega \hookrightarrow Z$ such that $\varphi(y_s) = z_s$ for all s . It finally follows that $\varphi_n|_\Omega$ converges to φ , which concludes the proof. \square

In the proofs of our main results we will use non-principal ultrafilters and ultra-limits of sequences. Recall for this that a non-principal ultrafilter on \mathbb{N} is a finitely additive probability measure ω on \mathbb{N} (together with the σ -algebra of all subsets) such that ω takes values in $\{0, 1\}$ only and $\omega(A) = 0$ whenever $A \subset \mathbb{N}$ is finite. Existence of non-principal ultra-filters on \mathbb{N} follows from Zorn's lemma. It is not difficult to prove that if (Y, τ) is a compact Hausdorff topological space, then for every sequence $(y_n)_{n \in \mathbb{N}} \subset Y$ there exists a unique point $y \in Y$ such that

$$\omega(\{n \in \mathbb{N} : y_n \in U\}) = 1$$

for every $U \in \tau$ containing y . We will call this point y the ultralimit of the sequence (y_n) and denote it by $\lim_\omega y_n$.

We are now ready to prove Theorems 1.1–1.2 stated in the introduction.

Proof of Theorem 1.1. We first prove (i). Let $S \in \mathbf{I}_m(X)$ with $\partial S = 0$ and set

$$s := \inf\{\mathbf{M}(T') : T' \in \mathbf{I}_{m+1}(X), \partial T' = S\}.$$

Note that the set appearing on the right hand side is non-empty. Indeed, if $m = 0$ then this follows from the fact that X is geodesic. If $m \geq 1$, then this follows from the isoperimetric inequality [8, Corollary 1.3]. Let now $(T_n) \subset \mathbf{I}_{m+1}(X)$ be a sequence satisfying $\partial T_n = S$ for all $n \in \mathbb{N}$ and such that $\mathbf{M}(T_n) \rightarrow s$. Clearly,

$$\sup_n [\mathbf{M}(T_n) + \mathbf{M}(\partial T_n)] < \infty.$$

Set $\Omega := \text{spt } S$ and note that Ω is separable. Let n_j , Z , φ_j , φ , and T be as in Lemma 3.1, where m is replaced by $m + 1$. By Lemma 2.3, $\varphi_{j\#} S$ converges weakly to $\varphi_{\#} S$ and hence $\partial T = \varphi_{\#} S$. Now, view X as a subspace of a dual Banach space Y and let ω be a non-principal ultrafilter on \mathbb{N} . Define a map $\psi : \text{spt } T \rightarrow Y$ as follows. Let $z \in \text{spt } T$. Since $\varphi_{j\#} T_{n_j}$ converges locally weakly to T , there exists a sequence $(x_j) \subset X$ with $x_j \in \text{spt } T_{n_j}$ for all $j \in \mathbb{N}$ and such that $\varphi_j(x_j) \rightarrow z$. Clearly, (x_j) is a bounded sequence in Y . Since closed balls of finite radius in Y , endowed with the weak*-topology, are compact and Hausdorff, it follows that (x_j) has an ultralimit $\lim_\omega x_j$ in Y . Define $\psi(z) := \lim_\omega x_j$. It follows from the lower semi-continuity of the norm in Y with respect to weak*-convergence that $\psi(z)$ is independent of the choice of sequence (x_j) and that ψ is 1-Lipschitz. Since $\varphi_j(x) \rightarrow \varphi(x)$ for all $x \in \text{spt } S$, it follows furthermore that $\psi \circ \varphi = \text{id}_{\text{spt } S}$.

Finally, if $P : Y \rightarrow X$ is a projection of norm 1, then $\hat{T} := (P \circ \psi)_\# T$ satisfies $\hat{T} \in \mathbf{I}_{m+1}(X)$ and

$$\partial \hat{T} = (P \circ \psi)_\#(\varphi_\# S) = P_\# S = S$$

and $\mathbf{M}(\hat{T}) \leq \mathbf{M}(T) \leq \liminf \mathbf{M}(T_{n_j}) = s$. This completes the proof of (i).

The proof of statement (ii) is analogous up to some minor modifications. Indeed, let $S \in \mathbf{M}_m(X)$ and set

$$s := \inf\{\mathbf{M}(T') + \mathbf{M}(\partial T' - S) : T' \in \mathbf{I}_{m+1}(X)\}.$$

Clearly, the set appearing on the right hand side is non-empty and thus s is finite. Let $(T_n) \subset \mathbf{I}_{m+1}(X)$ be a sequence satisfying $\mathbf{M}(T_n) + \mathbf{M}(\partial T_n - S) \rightarrow s$. It follows that

$$\sup_n [\mathbf{M}(T_n) + \mathbf{M}(\partial T_n)] < \infty.$$

Set $\Omega := \text{spt } S$ and note that $\text{spt } S$ is separable. Let n_j , Z , φ_j , φ , and T be as in Lemma 3.1, where m is replaced by $m + 1$. By Lemma 2.3, $\varphi_{j\#} S$ converges weakly to $\varphi_\# S$. Let Y and $P : Y \rightarrow X$ be as in (i). Define $\psi : \text{spt } T \cup \text{spt } S \rightarrow E$ in a similar way as ψ was defined in (i). Then ψ is 1-Lipschitz and $\psi \circ \varphi = \text{id}_{\text{spt } S}$. It follows that $\hat{T} := (P \circ \psi)_\# T$ satisfies $\hat{T} \in \mathbf{I}_{m+1}(X)$ and

$$\partial \hat{T} - S = (P \circ \psi)_\#(\partial T - \varphi_\# S).$$

We thus obtain from the lower semi-continuity of mass with respect to (local) weak convergence that

$$\begin{aligned} \mathbf{M}(\hat{T}) + \mathbf{M}(\partial \hat{T} - S) &\leq \mathbf{M}(T) + \mathbf{M}(\partial T - \varphi_\# S) \\ &\leq \liminf_{j \rightarrow \infty} [\mathbf{M}(T_{n_j}) + \mathbf{M}(\partial T_{n_j} - S)] = s. \end{aligned}$$

This completes the proof of (ii). \square

Remark 3.2. We remark that the use of non-principal ultrafilters and ultralimits can be avoided in the above proof in the case that X is the dual of a separable Banach space since in this case one can pass to a subsequence and use sequential weak*-compactness of closed bounded balls, see also the proof of Theorem 1.2.

The proof of Theorem 1.1 easily generalizes to the following context. Let (X, d) be a metric space and let ω denote a non-principal ultrafilter on \mathbb{N} . A sequence $(x_n)_{n \in \mathbb{N}} \subset X$ is called bounded if $\sup_n d(x_n, x_1) < \infty$. Define an equivalence relation \sim on bounded sequences in X by considering (x_n) and (x'_n) equivalent if $\lim_\omega d(x_n, x'_n) = 0$, and denote by $[(x_n)]$ the equivalence class of (x_n) with respect to \sim . The ultra-completion X_ω of X with respect to ω is the metric space

given by the set

$$X_\omega := \{[(x_n)] : (x_n) \subset X \text{ with } \sup d(x_n, x_1) < \infty\}$$

with the metric $d_\omega([(x_n)], [(x'_n)]) := \lim_\omega d(x_n, x'_n)$. We note that X isometrically embeds into X_ω by the map which assigns to $x \in X$ the equivalence class of the constant sequence (x) . We may therefore view X as a subset of X_ω . In the following we will say that a metric space X is 1-complemented in some ultra-completion of X if there exists a non-principal ultrafilter ω on \mathbb{N} such that X , viewed as a subset of X_ω , admits a 1-Lipschitz retraction of X_ω onto X . Note that if X is a dual Banach space, then X is 1-complemented in every ultra-completion of X . Consequently, if X is a Banach space which is 1-complemented in some dual Banach space (in the terminology established before the statement of Theorem 1.1), then X is 1-complemented in every ultra-completion of X . The following result generalizes Theorem 1.1.

Theorem 3.3. *Let X be a complete metric space and suppose that X is 1-complemented in some ultra-completion of X . Let $m \geq 0$. Then*

- (i) *for every $V \in \mathbf{I}_{m+1}(X)$ there exists $T \in \mathbf{I}_{m+1}(X)$ with $\partial T = \partial V$ and*

$$\mathbf{M}(T) \leq \mathbf{M}(T')$$

for all $T' \in \mathbf{I}_{m+1}(X)$ with $\partial T' = \partial V$,

- (ii) *for every $S \in \mathbf{M}_m(X)$ there exists $T \in \mathbf{I}_{m+1}(X)$ such that*

$$\mathbf{M}(T) + \mathbf{M}(\partial T - S) \leq \mathbf{M}(T') + \mathbf{M}(\partial T' - S)$$

for all $T' \in \mathbf{I}_{m+1}(X)$.

Note that the statement in (i) of Theorem 3.3 is slightly weaker than the statement (i) in Theorem 1.1 inasmuch as we assume that there exists a filling of ∂V . This is needed since in the generality considered in Theorem 3.3 integral currents without boundary need not have a filling.

Proof. The proof is analogous to the proof of Theorem 1.1. □

Theorem 1.3 now comes as a consequence of Theorem 3.3. Indeed, if X is a Hadamard space, then so is every ultra-completion X_ω of X and, as a closed convex subspace of X_ω , the nearest point projection from X_ω to X is 1-Lipschitz, see e.g. [3, Proposition II.2.4]. Therefore, in the terminology established above, X is 1-complemented in every ultra-completion of X . Statement (ii) of Theorem 1.3 therefore follows from statement (ii) of Theorem 3.3. As for (i), if $S \in \mathbf{I}_m(X)$

is such that $\partial S = 0$, then by [8, Corollary 1.4] there exists $V \in \mathbf{I}_{m+1}(X)$ with $\partial V = S$. Statement (i) of Theorem 1.3 now follows from this together with statement (i) of Theorem 3.3. If X is an injective space, then X is an absolute 1-Lipschitz retract and thus there exists a 1-Lipschitz projection of X_ω onto X and thus X is 1-complemented in every ultra-completion of X . Statements (i) and (ii) now follow as above.

We finally turn to the proof of our w^* -compactness theorem.

Proof of Theorem 1.2. Let (T_n) be as in the statement of the theorem. Set $\Omega := \{0\}$ and let n_j , Z , φ_j , φ , and T be as in Lemma 3.1. Then $\varphi_j(0)$ converges to some $z^0 \in Z$. Let $(z^k) \subset \text{spt } T$ be a dense sequence. For each $k \in \mathbb{N}$ choose a sequence $(x_j^k) \subset X$ with $x_j^k \in \text{spt } T_{n_j}$ for all j and such that $\varphi_j(x_j^k) \rightarrow z^k$. Since

$$\|x_j^k\| = d(\varphi_j(x_j^k), \varphi_j(0)) \rightarrow d(z^k, z^0) \quad \text{as } j \rightarrow \infty,$$

it follows that for k fixed the sequence (x_j^k) is bounded. After passing to a subsequence we may therefore assume that the sequence $\{x_j^k\}$ converges to some x^k in the weak*-topology of X . (Note that for this separability of a predual or reflexivity of X is needed.) Define $\psi(z^k) := x^k$ and note that ψ is 1-Lipschitz by the lower semi-continuity of the norm on X with respect to w^* -convergent sequences. Since the sequence (z^k) is dense in $\text{spt } T$, we can extend ψ to a 1-Lipschitz map $\psi : \text{spt } T \rightarrow X$. Set $\hat{T} := \psi_\# T$ and note that $\hat{T} \in \mathbf{I}_m(X)$. We will show that T_{n_j} is w^* -convergent to \hat{T} . For this let $f, \pi_1, \dots, \pi_m \in \text{Lip}(X)$ be weak*-continuous and such that $|f| \leq C$ for some C . For each j let $f^j : Z \rightarrow \mathbb{R}$ be a Lipschitz extension of $f \circ \varphi_j^{-1}$ with $\text{Lip}(f^j) = \text{Lip}(f)$ and such that $|f^j| \leq C$. Similarly, let $\pi_i^j : Z \rightarrow \mathbb{R}$ be a Lipschitz extension of $\pi_i \circ \varphi_j^{-1}$ with $\text{Lip}(\pi_i^j) = \text{Lip}(\pi_i)$. For each k we have

$$\begin{aligned} |f \circ \psi(z^k) - f^j(z^k)| &\leq |f(x^k) - f(x_j^k)| + |f^j(\varphi_j(x_j^k)) - f^j(z^k)| \\ &\leq |f(x^k) - f(x_j^k)| + \text{Lip}(f) \cdot d(\varphi_j(x_j^k), z^k), \end{aligned}$$

from which it follows together with the weak*-continuity of f that f^j converges pointwise to $f \circ \psi$ on $\text{spt } T$. Analogously, π_i^j converges to $\pi_i \circ \psi$ on $\text{spt } T$. Now, since $\varphi_{j\#} T_{n_j}$ converges locally weakly to T and

$$\lim_{r \rightarrow \infty} \left[\sup_n \|T_n\|(X \setminus B(0, r)) \right] = 0,$$

it follows that $\varphi_{j\#} T_{n_j}$ converges weakly to T . By possibly replacing Z by $\ell^\infty(Z)$ we may assume by [9, Theorem 1.4] that $\varphi_{j\#} T_{n_j}$ converges even in the flat norm to T . In particular, there exist $U_j \in \mathbf{I}_m(Z)$ and $V_j \in \mathbf{I}_{m+1}(Z)$ such that

$$T - \varphi_{j\#} T_{n_j} = U_j + \partial V_j \quad \text{and} \quad \mathbf{M}(U_j) + \mathbf{M}(V_j) \rightarrow 0.$$

We now obtain

$$\begin{aligned}
& |\hat{T}(f, \pi) - T_{n_j}(f, \pi)| \\
& \leq |T(f \circ \psi, \pi \circ \psi) - T(f^j, \pi^j)| + |T(f^j, \pi^j) - \varphi_{j\#} T_{n_j}(f^j, \pi^j)| \\
& \leq |T(f \circ \psi, \pi \circ \psi) - T(f \circ \psi, \pi^j)| + |T(f \circ \psi - f^j, \pi^j)| \\
& \quad + |U_j(f^j, \pi^j)| + |V_j(1, f^j, \pi^j)| \\
& \leq |T(f \circ \psi, \pi \circ \psi) - T(f \circ \psi, \pi^j)| \\
& \quad + \prod_{i=1}^m \text{Lip}(\pi_i) \int_Z |f \circ \psi - f^j| d\|T\| \\
& \quad + \prod_{i=1}^m \text{Lip}(\pi_i) [C + \text{Lip}(f)] \cdot (\mathbf{M}(U_j) + \mathbf{M}(V_j)).
\end{aligned}$$

Since each π_i^j converges pointwise to $\pi_i \circ \psi$ on $\text{spt } T$ with bounded Lipschitz constants, the first term after the last inequality sign converges to 0 by the continuity property of currents. Since f^j converges uniformly to $f \circ \psi$ on compact subsets of $\text{spt } T$ and $\text{spt } T$ is σ -compact, it follows that the second term converges to 0 as well. Since also the third term converges to 0, it follows that T_{n_j} indeed w^* -converges to \hat{T} as claimed. This concludes the proof. \square

The following example shows that in general the assumption in the statement of Theorem 1.2 that a predual of X be separable cannot be dropped.

Example 3.4. Let X denote the dual space of ℓ^∞ . For $n \geq 1$ define $x_n \in X$ by $x_n(a) := a_n$ for $a = (a_1, a_2, \dots) \in \ell^\infty$, and let the current $T_n \in \mathbf{I}_0(X)$ be given by $T_n(f) := f(x_n)$ for every $f \in \text{Lip}_b(X)$. It follows that T_n is supported in the closed unit ball in X and that $\mathbf{M}(T_n) = 1$ for every n . It is easy to show that there cannot exist a subsequence (n_j) such that T_{n_j} is w^* -convergent to some T . Indeed, given a subsequence (n_j) , let $a = (a_1, a_2, \dots) \in \ell^\infty$ be defined by $a_k = 1$ if $k = n_j$ for some j even and $a_k = 0$ otherwise. Define a weak*-continuous function $f \in \text{Lip}_b(X)$ by $f(x) := \varphi(x(a))$, where φ is the truncation function given by $\varphi(t) = \max\{-1, \min\{1, t\}\}$ for $t \in \mathbb{R}$. Clearly, $T_{n_j}(f) = f(x_{n_j})$ does not converge as $j \rightarrow \infty$.

We note that the only facts about dual spaces X of separable Banach spaces which are used in the proof of Theorem 1.2 are the sequential weak*-compactness of closed bounded balls in X and the lower semi-continuity of the norm on X with respect to weak*-convergent sequences. The proof therefore easily gives the following generalization of Theorem 1.2.

Theorem 3.5. *Let X be a Banach space and \mathcal{W} be a vector space topology on X such that closed bounded balls in X are sequentially \mathcal{W} -compact and such that the norm on X is lower semi-continuous with respect to \mathcal{W} -convergent sequences. Then the conclusion of Theorem 1.2 holds when w^* -convergence is replaced by \mathcal{W} -convergence.*

We note that \mathcal{W} -convergence of a sequence T_{n_j} to T means by definition that $T_{n_j}(f, \pi) \rightarrow T(f, \pi)$ for all $f, \pi_1, \dots, \pi_m \in \text{Lip}(X)$ which are also \mathcal{W} -continuous and such that f is bounded. Theorem 3.5 generalizes [2, Theorem A.1] for integral currents.

Bibliography

- [1] L. Ambrosio and B. Kirchheim, Currents in metric spaces, *Acta Math.* **185** (2000), no. 1, 1–80.
- [2] L. Ambrosio and T. Schmidt, Compactness results for normal currents and the Plateau problem in dual Banach spaces, *Proc. Lond. Math. Soc.* (3) (2012), DOI 10.1112/plms/pds066.
- [3] M. Bridson and A. Haefliger, *Metric Spaces of Non-Positive Curvature*, Grundlehren der Mathematischen Wissenschaften 319, Springer-Verlag, Berlin, 1999.
- [4] H. Federer and W.H. Fleming, Normal and integral currents, *Ann. of Math.* (2) **72** (1960), 458–520.
- [5] P. Harmand, D. Werner and W. Werner, *M-Ideals in Banach Spaces and Banach Algebras*, Lecture Notes in Mathematics 1547, Springer-Verlag, Berlin, 1993.
- [6] U. Lang, [Local currents in metric spaces](#), *J. Geom. Anal.* **21** (2011), 683–742.
- [7] U. Lang and S. Wenger, The pointed flat compactness theorem for locally integral currents, *Comm. Anal. Geom.* **19** (2011), no. 1, 159–190.
- [8] S. Wenger, Isoperimetric inequalities of Euclidean type in metric spaces, *Geom. Funct. Anal.* **15** (2005), no. 2, 534–554.
- [9] S. Wenger, Flat convergence for metric integral currents, *Calc. Var. Partial Differential Equations* **28** (2007), no. 2, 139–160.
- [10] S. Wenger, Compactness for manifolds and integral currents with bounded volume and diameter, *Calc. Var. Partial Differential Equations* **40** (2011), no. 3–4, 423–448.

Author information

Stefan Wenger, Department of Mathematics, University of Fribourg,
Chemin du Musée 23, 1700 Fribourg, Switzerland.
E-mail: stefan.wenger@unifr.ch