


**Appendix S1.** Bayesian phylogeny based on rbcL DNA sequence data illustrating the phylogenetic relationships among non-native plants included in this study. Species names within brackets are those lacking DNA sequence data and were replaced by taxa within the same genus indicated outside the brackets. Plant families are indicated above branches. High nodal support (posterior probabilities >0.9) is indicated as black rectangular boxes at nodes.



**Appendix S2.** List of studies for meta-analysis on non-native plant species impact on local plant and animal species richness.

- Adams SN, Engelhardt KAM (2009) Diversity declines in *Microstegium vimineum* (Japanese stiltgrass) patches. – *Biological Conservation* 142: 1003–1010.
- Aguilera AG, Alpert P, Dukes JS (2010) Impacts of the invasive plant *Fallopia japonica* (Houtt.) on plant communities and ecosystem processes. – *Biological Invasions* 12: 1243-1252.
- Andreu J, Manzano E, Bartomeus I, Dana ED, Vilà M (2010) Vegetation response after removal of the invader *Carpobrotus* spp. in coastal dunes. – *Ecological Restoration* 28: 440–448.
- Angeloni NL, Jankowski KJ, Tuchman NC, Kelly JJ (2006) Effects of an invasive cattail species (*Typha x glauca*) on sediment nitrogen and microbial community composition in a freshwater wetland. – *Fems Microbiology Letters* 263: 86–92.
- Auerbach M, Simberloff D (1988) Rapid leaf-miner colonization of introduced trees and shifts in sources of herbivore mortality. – *Oikos* 52: 41–50.
- Badano EI, Jones CG, Cavieres LA, Wright JP (2006) Assessing impacts of ecosystem engineers on community organization: a general approach illustrated by effects of a high-Andean cushion plant. – *Oikos* 115: 369–385.
- Bakker J, Wilson S (2001) Competitive abilities of introduced and native grasses. – *Plant Ecology* 157: 117–125.
- Bartomeus I, Vilà M, Santamaria L (2008) Contrasting effects of invasive plants in plant-pollinator networks. – *Oecologia* 155: 761–770.
- Baskett CA, Emery SM, Rudgers JA (2011) Pollinator Visits to Threatened Species Are Restored Following Invasive Plant Removal. – *International Journal of Plant Sciences* 172: 411-422.
- Belnap J, Phillips SL, Sherrod SK, Moldenke A (2005) Soil biota can change after exotic plant invasion: does this affect ecosystem processes? – *Ecology* 86: 3007–3017.
- Belnap J, Phillips SL, Troxler T (2006) Soil lichen and moss cover and species richness can be highly dynamic: The effects of invasion by the annual exotic grass *Bromus tectorum*, precipitation, and temperature on biological soil crusts in SE Utah. – *Applied Soil Ecology* 32: 63–76.

- Brandon AL, Gibson DJ, Middleton BA (2004) Mechanisms for dominance in an early successional old field by the invasive non-native *Lespedeza cuneata* (Dum. Cours.) G. Don. – *Biological Invasions* 6: 483–493.
- Chabrierie O, Loinard J, Perrin S, Saguez R, Decocq G (2010) Impact of *Prunus serotina* invasion on understory functional diversity in a European temperate forest. – *Biological Invasions* 12: 1891–1907.
- Christian JM, Wilson SD (1999) Long-term ecosystem impacts of an introduced grass in the northern great plains. – *Ecology* 80: 2397–2407.
- Cuevas YA, Zalba SM (2010) Recovery of Native Grasslands after Removing Invasive Pines. – *Restoration Ecology* 18: 711–719.
- Cushman JH, Gaffney KA (2010) Community-level consequences of invasion: impacts of exotic clonal plants on riparian vegetation. – *Biological Invasions* 12: 2765–2776.
- Dassonville N, Vanderhoeven S, Gruber W, Meerts P (2007) Invasion by *Fallopia japonica* increases topsoil mineral nutrient concentrations. – *Ecoscience* 14: 230–240.
- Davies KW, Svejcar TJ (2008) Comparison of medusahead-invaded and noninvaded Wyoming big sagebrush steppe in southeastern Oregon. – *Rangeland Ecology Management* 61: 623–629.
- de Groot M, Kleijn D, Jogan N (2007) Species groups occupying different trophic levels respond differently to the invasion of semi-natural vegetation by *Solidago canadensis*. – *Biological Conservation* 136: 612–617.
- De Meester JE, Richter DdeB (2010) Restoring restoration: removal of the invasive plant *Microstegium vimineum* from a North Carolina wetland. – *Biological Invasions* 12: 781–793.
- Domenech R, Vilà M (2008) Response of the invader *Cortaderia selloana* and two coexisting natives to competition and water stress. – *Biological Invasions* 10: 903–912.
- Douglas MM, O'Connor RA (2003) Effects of the exotic macrophyte, para grass (*Urochloa mutica*), on benthic and epiphytic macroinvertebrates of a tropical floodplain. – *Freshwater Biology* 48: 962–971.
- Dunbar KR, Facelli JM (1999) The impact of a novel invasive species, *Orbea variegata* (African carrion flower), on the chenopod shrublands of South Australia. – *Journal of Arid Environments* 41: 37–48.

- Durst SL, Theimer TC, Paxton EH, Sogge MK (2008) Temporal variation in the arthropod community of desert riparian habitats with varying amounts of saltcedar (*Tamarix ramosissima*). – *Journal of Arid Environment* 72: 1644–1653.
- Fischer LK, von der Lippe M, Kowarik I (2009) Tree invasion in managed tropical forest facilitates endemic species. – *Journal of Biogeography* 36: 2251–2263.
- Flory SL, Clay K (2009) Invasive plant removal method affects native community diversity responses. – *Journal of Applied Ecology* 4: 434–442.
- Flory SL, Clay K (2010) Non-native grass invasion alters native plant composition in experimental communities. – *Biological Invasions* 12: 1285–1294.
- Franklin K, Molina-Freaner F (2010) Consequences of Buffelgrass Pasture Development for Primary Productivity, Perennial Plant Richness, and Vegetation Structure in the Drylands of Sonora, Mexico. – *Conservation Biology* 24: 1664–1673.
- French K, Major RE (2001) Effect of an exotic *Acacia* (Fabaceae) on ant assemblages in South African fynbos. – *Austral Ecology* 26: 303–310.
- French K, Mason TJ, Sullivan N (2011) Recruitment limitation of native species in invaded coastal dune communities. – *Plant Ecology* 212: 601–609.
- Gaertner M, Richardson DM, Privett SDJ (2011) Effects of Alien Plants on Ecosystem Structure and Functioning and Implications for Restoration: Insights from Three Degraded Sites in South African Fynbos. – *Environmental Management* 48: 57–69.
- Gan X, Cai Y, Choi C, Ma Z, Chen J, Li B (2009) Potential impacts of invasive *Spartina alterniflora* on spring bird communities at Chongming Dongtan, a Chinese wetland of international importance. *Estuarine, Coastal and Shelf Science* 83: 211–218.
- García-de-Lomas J, Cózar A, Dana ED, Hernández I (2009) Invasiveness of *Galenia pubescens* (Aizoaceae): A new threat to Mediterranean-climate coastal ecosystems. – *Acta Oecologica* 36: 39–45.
- Giantomasi A, Tecco PA, Funes G, Gurvich DE, Cabido M (2008) Canopy effects of the invasive shrub *Pyracantha angustifolia* on seed bank composition, richness and density in a montane shrubland (Cordoba, Argentina). – *Austral Ecology* 33: 68–77.

- Gooden, B, French, K, Turner P (2009) Invasion and management of a woody plant, *Lantana camara* L., alters vegetation diversity within wet sclerophyll forest in south-eastern Australia. – *Forest Ecology and Management* 257: 960–967.
- Gratton C, Denno RF (2005) Restoration of arthropod assemblages in a *Spartina* salt marsh following removal of the invasive plant *Phragmites australis*. – *Restoration Ecology* 13: 358–372.
- Greenwood H, O'Dowd DJ, Lake PS (2004). Willow (*Salix x rubens*) invasion of the riparian zone in south-eastern Australia: reduced abundance and altered composition of terrestrial arthropods. – *Diversity Distrib.* 10: 485–492.
- Hagen EN, Barker JD, Gara RI (2010) Aerial Arthropod Communities of Native and Invaded Forests, Robinson Crusoe Island, Chile. – *Environmental Entomology* 39: 1159-1164.
- Hanula JL, Horn S (2011) Removing an exotic shrub from riparian forests increases butterfly abundance and diversity. – *Forest Ecology and Management* 262: 674–680.
- Hejda M (2012) What Is the Impact of *Impatiens parviflora* on Diversity and Composition of Herbal Layer Communities of Temperate Forests? *PLoS ONE* 7: e39571.
- Hejda M, Pyšek P (2006) What is the impact of *Impatiens glandulifera* on species diversity of invaded riparian vegetation? – *Biological Conservation* 132: 143–152.
- Hejda M, Pyšek P, Jarošík V (2009) Impact of invasive plants on the species richness, diversity and composition of invaded communities. – *Journal of Ecology* 97: 393–403.
- Henderson DC, Naeth MA (2005) Multi-scale impacts of crested wheatgrass invasion in mixed-grass prairie. – *Biological Invasions* 7: 639–650.
- Herrera AM, Dudley TL (2003) Reduction of riparian arthropod abundance and diversity as a consequence of giant reed (*Arundo donax*) invasion. – *Biological Invasions* 5: 167–177.
- Holland-Clift S, O'Dowd DJ, Mac Nally R (2011) Impacts of an invasive willow (*Salix x rubens*) on riparian bird assemblages in south-eastern Australia. – *Austral Ecology* 36: 511–520.
- Holmes PM, Cowling RM (1997) The effects of invasion by *Acacia saligna* on the guild structure and regeneration capabilities of South African fynbos shrublands. – *Journal of Applied Ecology* 34: 317–332.

- Holmquist JG, Schmidt-Gengenbach J, Slaton MR (2011) Influence of invasive palms on terrestrial arthropod assemblages in desert spring habitat. – *Biological Conservation* 144: 518–525.
- Houston WA, Duivenvoorden LJ (2002) Replacement of littoral native vegetation with the ponded pasture grass *Hymenachne amplexicaulis*: effects on plant, macroinvertebrate and fish biodiversity of backwaters in the Fitzroy River, Central Queensland, Australia. – *Marine Freshwater Research* 53: 1235–1244.
- Hua C , Jian L, Yongli Z, Qiang W, Xiuli G, Yinghua W, Renqing W (2011) Influence of invasive plant *Coreopsis grandiflora* on functional diversity of soil microbial communities. – *Journal of Environmental Biology* 32: 567-572.
- Hulme PE, Bremner ET (2006) Assessing the impact of *Impatiens glandulifera* on riparian habitats: partitioning diversity components following species removal. – *Journal of Applied Ecology* 43: 43–50.
- Jackson J (2005) Is there a relationship between herbaceous species richness and buffel grass (*Cenchrus ciliaris*)? – *Austral Ecology* 30: 505–517.
- Jäger H, Kowarik I (2010) Resilience of Native Plant Community Following Manual Control of Invasive *Cinchona pubescens* in Galápagos. – *Restoration Ecology* 18: 103–112.
- Jäger H, Tye A, Kowarik I (2007) Tree invasion in naturally treeless environments: Impacts of quinine (*Cinchona pubescens*) trees on native vegetation in Galápagos. – *Biological Conservation* 140: 297–307.
- Jeddi K, Cortina J, Chaie M (2009) *Acacia salicina*, *Pinus halepensis* and *Eucalyptus occidentalis* improve soil surface conditions in arid southern Tunisia. – *Journal of Arid Environments* 73: 1005–1013.
- Kappes H, Lay R, Topp W (2007) Changes in different trophic levels of litter-dwelling macrofauna associated with giant knotweed invasion. – *Ecosystems* 10: 734–744.
- Kaur R, Malhorta S, Inderjit (2012) Effects of invasion of *Mikania micrantha* on germination of rice seedlings, plant richness, chemical properties and respiration of soil. – *Biological Fertilization Soils* 48: 481–488.
- Kwiatkowska AJ, Spalik K, Michalak E, Palińska A, Panufnik D (1997) Influence of the size and density of *Carpinus betulus* on the spatial distribution and rate of deletion of forest-floor species in thermophilous oak forest. – *Plant Ecology* 129: 1–10.

- Lindsay EA, French K (2006) The impact of the weed *Chrysanthemoides monilifera* ssp. *rotundata* on coastal leaf litter invertebrates. – *Biological Invasions* 8: 177–192.
- Ma Z, Gan X, Cai Y, Chen J, Li B (2011) Effects of exotic *Spartina alterniflora* on the habitat patch associations of breeding saltmarsh birds at Chongming Dongtan in the Yangtze River estuary, China. – *Biological Invasions* 13: 1673–1686.
- Marchante H, Freitas H, Hoffmann JH (2011) Post-clearing recovery of coastal dunes invaded by *Acacia longifolia*: is duration of invasion relevant for management success? – *Journal of Applied Ecology* 48: 1295–1304.
- Marchante H, Marchante E, Freitas H (2003) Invasion of the Portuguese dune ecosystems by the exotic species *Acacia longifolia* (Andrews) Willd.: Effects at the community level. – In: *Plant Invasions: Ecological Threats and Management Solutions*, L.E. Child, J.H. Brock, G. Brundu, K. Prach, P. Pyšek, P.M. Wade, M. Williamson (Eds). Backhuys Publishers, Leiden, The Netherlands, pp 75–85.
- Martin PH (1999) Norway Maple (*Acer platanoides*) invasion of a natural forest stand: understory consequence and regeneration pattern. – *Biological Invasions* 1: 215–222.
- Mason TJ, French K (2008) Impacts of a woody invader vary in different vegetation communities. – *Diversity and Distributions* 14: 829–838.
- Mason TJ, French K, Russell KG (2007) Moderate impacts of plant invasion and management regimes in coastal hind dune seed banks. – *Biological Conservation* 134: 428–439.
- Mealor BA, Hild AL, Shaw NL (2004) Native plant community composition and genetic diversity associated with long-term weed invasions. – *Western North American Naturalist* 64: 503–513.
- Meffin R, Miller AL, Hulme PE, Duncan RP (2010) Experimental introduction of the alien plant *Hieracium lepidulum* reveals no significant impact on montane plant communities in New Zealand. – *Diversity and Distributions* 16: 804–815.
- Meggaro Y, Vilà M (2002) Distribución de dos especies arbóreas exóticas *Ailanthus altissima* y *Robinia pseudoacacia* en el Parque de Collserola (Barcelona). – *Montes* 68: 25–33.
- Meyerson L, Saltonstall K, Windham L, Kiviat E, Findlay S (2000) A comparison of *Phragmites australis* in freshwater and brackish marsh environments in North America. – *Wetlands Ecology Management* 8: 89–103.

- Minden V, Hennenberg KJ, Porembski S, Boehmer HJ (2010) Invasion and management of alien *Hedychium gardnerianum* (kahili ginger, Zingiberaceae) alter plant species composition of a montane rainforest on the island of Hawai'i. – *Plant Ecology* 206: 321–333.
- Mitchell ME, Lishawa SC, Geddes P, Larkin DJ, Treering D, Tuchman NC (2011) Time-Dependent Impacts of Cattail Invasion in a Great Lakes Coastal Wetland Complex. *Wetlands* 31: 1143–1149.
- Moroń D, Lenda M, Skórka P, Szentgyörgyi H, Settele J, Woyciechowski M (2009) Wild pollinator communities are negatively affected by invasion of alien goldenrods in grassland landscapes. – *Biological Conservation* 142: 1322–1332.
- Mummey DL, Rillig MC (2006) The invasive plant species *Centaurea maculosa* alters arbuscular mycorrhizal fungal communities in the field. – *Plant and Soil* 288: 81–90.
- Muralli KS, Setty RS (2001) Effect of weeds *Lantana camara* and *Chromelina odorata* growth on the species diversity, regeneration and stem density of tree and shrub layer in BRT sanctuary. – *Current Science* 80: 675–678.
- Odat N, Al Khateeb W, Muhaidat R, Al U'datt M, Irshiad L (2011) The effect of exotic *Acacia saligna* tree on plant biodiversity of Northern Jordan. – *International Journal of Agricultural Biology* 13: 823–826.
- Ostoja SM, Schupp EW (2009) Conversion of sagebrush shrublands to exotic annual grasslands negatively impacts small mammal communities. *Diversity and Distributions* 15: 863–870.
- Ostoja SM, Schupp EW (2009) Conversion of sagebrush shrublands to exotic annual grasslands negatively impacts small mammal communities. - *Diversity and Distributions* 15: 863–870.
- Oswalt CM, Oswalt SN, Clatterbuck WK (2007) Effects of *Microstegium vimineum* (Trin.) A. Camus on native woody species density and diversity in a productive mixed-hardwood forest in Tennessee. – *Forest Ecology and Management* 242: 727–732.
- Parr CL, Ryan BJ, Setterfield SA (2010) Habitat Complexity and Invasive Species: The Impacts of Gamba Grass (*Andropogon gayanus*) on Invertebrates in an Australian Tropical Savanna. – *Biotropica* 4: 688–696.
- Paterson ID, Coetzee JA, Hill MP, Downie DD (2011) A pre-release assessment of the relationship between the invasive alien plant, *Pereskia aculeata* Miller (Cactaceae), and native plant biodiversity in South Africa. – *Biological Control*

57: 59–65.

- Pawson SM, McCarthy JK, Ledgard NJ, Didham RK (2010) Density-dependent impacts of exotic conifer invasion on grassland invertebrate assemblages. – *Journal of Applied Ecology* 47: 1053–1062.
- Petillon J, Ysnel F, Canard A, Lefeuvre JC (2005) Impact of an invasive plant (*Elymus athericus*) on the conservation value of tidal salt marshes in western France and implications for management: Responses of spider populations. – *Biological Conservation* 126: 103–117.
- Pyšek P, Pyšek A (1995) Invasion by *Heracleum mantegazzianum* in different habitats in the Czech Republic. – *Journal of Vegetation Science*: 711–718.
- Rascher KG, Große-Stoltenberg A, Máguas C, Meira-Neto JAA, Werner C (2011) *Acacia longifolia* invasion impacts vegetation structure and regeneration dynamics in open dunes and pine forests. – *Biological Invasions* 13: 1099–1113.
- Rayamajhi MB, Pratt PD, Center TD, Tipping PW, Van TK (2009) Decline in exotic tree density facilitates increased plant diversity: the experience from *Melaleuca quinquenervia* invaded wetlands. – *Wetlands Ecology and Management* 17: 455–467.
- Reinhart KO, Green E, Callaway RM (2005) Effects of *Acer platanoides* invasion on understory plant communities and tree regeneration in the northern Rocky Mountains. – *Ecography* 28: 573–582.
- Richburg JA, Patterson WA, Lowenstein F (2001) Effects of road salt and *Phragmites australis* invasion on the vegetation of a western Massachusetts calcareous lake-basin fen. – *Wetlands* 21: 247–255.
- Robertson MR, Harris KR, Coetzee JA, Foxcroft LC, Dippenaar-Schoeman AS, von Rensburg BJ (2011) Assessing local scale impacts of *Opuntia stricta* (Cactaceae) invasion on beetle and spider diversity in Kruger National Park, South Africa. – *African Zoology* 46: 205–223.
- Robertson SG, Hickman KR (2012) Aboveground plant community and seed bank composition along an invasion gradient. – *Plant Ecology* 213: 1461–1475.
- Scharfy D, Eggenschwiler H, Olde Venterink H, Edwards PJ, Güsewell S (2009) The invasive alien plant species *Solidago gigantea* alters ecosystem properties across habitats with differing fertility. – *Journal of Vegetation Science* 20: 1072–1085.

- Schutzenhofer MR, Valone TJ (2006) Positive and negative effects of exotic *Erodium cicutarium* in an arid ecosystem. – *Biological Conservation* 132: 376–381.
- Shaben J, Myers JH (2010) Relationships between Scotch broom (*Cytisus scoparius*), soil nutrients, and plant diversity in the Garry oak savannah ecosystem. – *Plant Ecology* 207: 81–91.
- Simao MCM, Flory SL, Rudgers JA (2010) Experimental plant invasion reduces arthropod abundance and richness across multiple trophic levels. – *Oikos* 119: 1553–1562.
- Skórka P, Lenda M, Tryjanowski P (2010) Invasive alien goldenrods negatively affect grassland bird communities in Eastern Europe. – *Biological Conservation* 143: 856–861.
- Spyreas G, Wilm BW, Plocher AE, Ketzner DM, Matthews JW, Ellis JL, Heske EJ (2010) Biological consequences of invasion by reed canary grass (*Phalaris arundinacea*). – *Biological Invasions* 12: 1253–1267.
- Standish RJ (2004) Impact of an invasive clonal herb on epigeic invertebrates in forest remnants in New Zealand. – *Biological Conservation* 116: 49–58.
- Stinson K, Kaufman S, Durbin L, Lowenstein F (2007) Impacts of garlic mustard invasion on a forest understory community. – *Northeastern Naturalist* 14: 73–88.
- Timsina B, Shrestha BB, Rokaya MB, Münzbergová Z (2011) Impact of *Parthenium hysterophorus* L. invasion on plant species composition and soil properties of grassland communities in Nepal. – *Flora* 206: 233–240.
- Treberg MA, Husband BC (1999) Relationship between the abundance of *Lythrum salicaria* (purple loosestrife) and plant species richness along the Bar River, Canada. – *Wetlands* 19: 118–125.
- Truscott AM, Palmer SC, Soulsby C, Westaway S, Hulme PE (2008) Consequences of invasion by the alien plant *Mimulus guttatus* on the species composition and soil properties of riparian plant communities in Scotland. – *Perspectives in Plant Ecology, Evolution and Systematics* 10: 231–240.
- Turner PJ, Scott JK, Spafford H (2008) The ecological barriers to the recovery of bridal creeper (*Asparagus asparagoides* (L.) Druce) infested sites: Impacts on vegetation and the potential increase in other exotic species. – *Austral Ecology* 33: 713–722.

- Valtonen A, Jantunen J, Saarinen K (2006) Flora and lepidoptera fauna adversely affected by invasive *Lupinus polyphyllus* along road verges. – *Biological Conservation* 133: 389–396.
- Vilà M, Tessier M, Suehs CM, Brundu G, Carta L, Galanidis A, Lambdon P, Manca M, Medail F, Moragues E, Traveset A, Troumbis AY, Hulme PE (2006) Local and regional assessments of the impacts of plant invaders on vegetation structure and soil properties of Mediterranean islands. – *Journal of Biogeography* 33: 853–861.
- Vincent Florens FB, Mauremootoo JR, Fowler SV, Winder L, Baider C (2010) Recovery of indigenous butterfly community following control of invasive alien plants in a tropical island's wet forests. – *Biodiversity Conservation* 19: 3835–3848.
- Watling JI, Hickman CR, Orrock JL (2011) Invasive shrub alters native forest amphibian communities. – *Biological Conservation* 144: 2597–2601.
- Wearne LJ, Morgan JW (2004) Community-level changes in Australian subalpine vegetation following invasion by the non-native shrub *Cytisus scoparius*. – *Journal of Vegetation Science* 15: 595–604.
- Wilcox J, Beck CW (2007) Effects of *Ligustrum sinense* Lour. (Chinese privet) on abundance and diversity of songbirds and native plants in a southeastern nature preserve. – *Southeastern Naturalist* 6: 535–550.
- Woods KD (1993) Effects of invasion by *Lonicera tatarica* L on herbs and tree seedlings in 4 New-England forests. – *The American Midland Naturalist* 130: 62–74.
- Wootton LS, Halsey SD, Bevaart K, McGough A, Ondreicka J, Patel P (2005) When invasive species have benefits as well as costs: Managing *Carex kobomugi* (Asiatic sand sedge) in New Jersey's coastal dunes. – *Biological Invasions* 7: 1017–1027.
- Wu YT, Wang CH, Zhang XD, Zhao B, Jiang LF, Chen JK, Li B (2009) Effects of saltmarsh invasion by *Spartina alterniflora* on arthropod community structure and diets. – *Biological Invasions* 11: 635–649.
- Wyckoff PH, Webb SL (1996) Understory influence of the invasive Norway Maple (*Acer platanoides*). – *Bulletin of the Torrey Botanical Club* 123: 197–205.
- Yoshida K, Oka S (2004) Invasion of *Leucaena leucocephala* and its effects on the native plant community in the Ogasawara (Bonin) Islands. – *Weed Technology* 18: 1371–1375.

Young HS, Raab TK, McCauley DJ, Briggs AA, Dirzo R (2010) The cocconut palm, *Cocos nucifera*, impacts forest composition and soil characteristics at Palmyra Atoll, Central Pacific. – Journal of Vegetation Science 21: 1058–1068.

**Appendix S3.** R code to test the effect sizes of the impact of non-native plant species on resident plant species richness.

```
rm(list=ls())

#load needed libraries
library(ape)
library(nlme)
library(MASS)

#functions for the Meta-Analysis

model_fit <- function(formula,tree,d){
  #fit of a Meta-Anyalsis based on phylogenetic regression

  mint <- gls(formula,
correlation=corGrafen(0.5,tree,fixed=FALSE),method='ML',weights=varFixed(~w),data
=d)

  r <- d$y - fitted(mint)
  Q <- sum(d$w*r^2)
  df <- length(r) - 1
  C <- sum(d$w) - sum(d$w^2) / sum(d$w)
  T2 <- (Q - df)/C
  new_w <- 1/(1/d$w + T2)
  d$new_w <- new_w

  m <- gls(formula,
correlation=corGrafen(0.5,tree,fixed=FALSE),method='ML',weights=varFixed(~new_w)
,data=d)
}

#load phylogenetic tree
tr_plant <- read.tree(file='tree_plant.txt')

#load data set
d_plant <- read.table(file='dataset_plant.txt',header=TRUE)

#compute effect size and the weight
y <- log(d_plant$mean_control / d_plant$mean_invaded)
```

```
names(y) <- rownames(d_plant)
sdy <- (d_plant$SD_control / d_plant$mean_control + d_plant$SD_invaded /
d_plant$mean_invaded)
w <- 1/sdy^2
```

```
#covariables
```

```
x1 <- factor(d_plant$Life_form)
x2 <- factor(d_plant$N_fixing)
x3 <- factor(d_plant$Clonal)
x4 <- factor(d_plant$Biogeographical_region)
x5 <- factor(d_plant$Insularity.)
x6 <- factor(d_plant$Habitat_type)
```

```
#creat the data frame made of response variable, cavariables, and weights
```

```
d2 <- as.data.frame(x = y)
d2$x1 <- x1
d2$x2 <- x2
d2$x3 <- x3
d2$x4 <- x4
d2$x5 <- x5
d2$x6 <- x6
d2$w <- w
```

```
#full model
```

```
formula = y ~ -1 + x1 + x2 + x3 + x4 + x5 + x6
m <- model_fit(formula,tr_plant,d2)
AIC(m)
```

```
#model selection procedure
```

```
#first step
```

```
#model without x1
```

```
formula = y ~ -1 + x2 + x3 + x4 + x5 + x6
m_x1 <- model_fit(formula,tr_plant,d2)
AIC(m_x1)
```

```
#model without x2
```

```
formula = y ~ -1 + x1 + x3 + x4 + x5 + x6
m_x2 <- model_fit(formula,tr_plant,d2)
AIC(m_x2)
```

```
#model without x3
```

```
formula = y ~ -1 + x1 + x2 + x4 + x5 + x6
m_x3 <- model_fit(formula,tr_plant,d2)
AIC(m_x3)
```

```

#model without x4
formula = y ~ -1 + x1 + x2 + x3 + x5 + x6
m_x4 <- model_fit(formula,tr_plant,d2)
AIC(m_x4)
#model without x5
formula = y ~ -1 + x1 + x2 + x3 + x4 + x6
m_x5 <- model_fit(formula,tr_plant,d2)
AIC(m_x5)
#model without x6
formula = y ~ -1 + x1 + x2 + x3 + x4 + x5
m_x6 <- model_fit(formula,tr_plant,d2)
AIC(m_x6)

AIC(m,m_x1,m_x2,m_x3,m_x4,m_x5,m_x6)

```

```

#based on AIC, we remove the covariable x4
formula = y ~ -1 + x1 + x2 + x3 + x5 + x6
m <- model_fit(formula,tr_plant,d2)
AIC(m)

```

```

#second step
#model without x1
formula = y ~ -1 + x2 + x3 + x5 + x6
m_x1 <- model_fit(formula,tr_plant,d2)
AIC(m_x1)
#model without x2
formula = y ~ -1 + x1 + x3 + x5 + x6
m_x2 <- model_fit(formula,tr_plant,d2)
AIC(m_x2)
#model without x3
formula = y ~ -1 + x1 + x2 + x5 + x6
m_x3 <- model_fit(formula,tr_plant,d2)
AIC(m_x3)
#model without x5
formula = y ~ -1 + x1 + x2 + x3 + x6
m_x5 <- model_fit(formula,tr_plant,d2)
AIC(m_x5)
#model without x6

formula = y ~ -1 + x1 + x2 + x3 + x5

m_x6 <- model_fit(formula,tr_plant,d2)
AIC(m_x6)

```

```
AIC(m,m_x1,m_x2,m_x3,m_x5,m_x6)
```

```
#based on AIC, we remove the covariable x1  
formula = y ~ -1 + x2 + x3 + x5 + x6  
m <- model_fit(formula,tr_plant,d2)  
AIC(m)
```

```
#third step  
#model without x2  
formula = y ~ -1 + x3 + x5 + x6  
m_x2 <- model_fit(formula,tr_plant,d2)  
AIC(m_x2)  
#model without x3  
formula = y ~ -1 + x2 + x5 + x6  
m_x3 <- model_fit(formula,tr_plant,d2)  
AIC(m_x3)  
#model without x5  
formula = y ~ -1 + x2 + x3 + x6  
m_x5 <- model_fit(formula,tr_plant,d2)  
AIC(m_x5)  
#model without x6  
formula = y ~ -1 + x2 + x3 + x5  
m_x6 <- model_fit(formula,tr_plant,d2)  
AIC(m_x6)
```

```
AIC(m,m_x2,m_x3,m_x5,m_x6)
```

```
#based on AIC, we remove the covariable x6  
formula = y ~ -1 + x2 + x3 + x5  
m <- model_fit(formula,tr_plant,d2)  
AIC(m)
```

```
#fourth step  
#model without x2  
formula = y ~ -1 + x3 + x5  
m_x2 <- model_fit(formula,tr_plant,d2)  
AIC(m_x2)  
#model without x3  
formula = y ~ -1 + x2 + x5  
m_x3 <- model_fit(formula,tr_plant,d2)
```

```
AIC(m_x3)
#model without x5
formula = y ~ -1 + x2 + x3
m_x5 <- model_fit(formula,tr_plant,d2)
AIC(m_x5)
```

```
AIC(m,m_x2,m_x3,m_x5)
```

```
#based on AIC, we remove the covariable x5
formula = y ~ -1 + x2 + x3
m <- model_fit(formula,tr_plant,d2)
AIC(m)
```

```
#fifth step
#model without x2
formula = y ~ -1 + x3
m_x2 <- model_fit(formula,tr_plant,d2)
AIC(m_x2)
#model without x3
formula = y ~ -1 + x2
m_x3 <- model_fit(formula,tr_plant,d2)
AIC(m_x3)
```

```
AIC(m,m_x2,m_x3)
#based on AIC, the procedure has converged
#####
## compute the delta AIC
```

```
formula = y ~ -1 + x2 + x3
mfull <- model_fit(formula,tr_plant,d2)
AIC(mfull)
summary(mfull)
```

```
#-x2
formula = y ~ -1 + x3
mfull_x2 <- model_fit(formula,tr_plant,d2)
AIC(mfull_x2)
```

```
#-x3
formula = y ~ -1 + x2
mfull_x3 <- model_fit(formula,tr_plant,d2)
AIC(mfull_x3)
```

```
#-phylo
```

```
mfull_phylo <- gls(y ~ -1 + x2 + x3,  
correlation=corPagel(0,tr_plant,fixed=TRUE),method='ML',weights=varFixed(~w))  
r <- y - fitted(mfull_phylo)  
Q <- sum(w*r^2)  
df <- length(r) - 1  
C <- sum(w) - sum(w^2) / sum(w)  
T2 <- (Q - df)/C  
new_w <- 1/(1/w + T2)  
mfull_phylo <- gls(y ~ -1 + x1 + x3 + x2 + x6,  
correlation=corPagel(0,tr_plant,fixed=TRUE),method='ML',weights=varFixed(~new_w)  
)  
AIC(mfull_phylo)  
  
AIC(mfull_x2,mfull_x3,mfull_phylo)-AIC(mfull)
```

**Appendix S4.** R code to test the effect sizes of the impact of non-native plant species on resident animal species richness.

```
rm(list=ls())

#load needed libraries
library(ape)
library(nlme)
library(MASS)

#functions for the Meta-Analysis

model_fit <- function(formula,tree,d){
  #fit of a Meta-Anyalsis based on phylogenetic regression

  mint <- gls(formula,
correlation=corGrafen(0.5,tree,fixed=FALSE),method='ML',weights=varFixed(~w),data
=d)

  r <- d$y - fitted(mint)
  Q <- sum(d$w*r^2)
  df <- length(r) - 1
  C <- sum(d$w) - sum(d$w^2) / sum(d$w)
  T2 <- (Q - df)/C
  new_w <- 1/(1/d$w + T2)
  d$new_w <- new_w

  m <- gls(formula,
correlation=corGrafen(0.5,tree,fixed=FALSE),method='ML',weights=varFixed(~new_w)
,data=d)
}

model_fit_no_phylo <- function(formula,d){
  #fit of a Meta-Anyalsis based with a standard linear regression

  mint <- gls(formula,method='ML',weights=varFixed(~w),data=d)

  r <- d$y - fitted(mint)
  Q <- sum(d$w*r^2)
  df <- length(r) - 1
  C <- sum(d$w) - sum(d$w^2) / sum(d$w)
```

```

T2 <- (Q - df)/C
new_w <- 1/(1/d$w + T2)
d$new_w <- new_w

m <- gls(formula,method='ML',weights=varFixed(~new_w),data=d)

}

#load phylogenetic tree
tr_animal <- read.tree(file='tree_animal.txt')

#load data set
d_animal <- read.table(file='dataset_animal.txt',header=TRUE)

#compute effect size and the weight
y <- log(d_animal$mean_control / d_animal$mean_invaded)
names(y) <- rownames(d_animal)
sdy <- (d_animal$SD_control / d_animal$mean_control + d_animal$SD_invaded /
d_animal$mean_invaded)
w <- 1/sdy^2

#covariables
x1 <- factor(d_animal$Life_form)
x2 <- factor(d_animal$N_fixing)
x3 <- factor(d_animal$Clonal)
x4 <- factor(d_animal$Biogeographical_region)
x5 <- factor(d_animal$Insularity.)
x6 <- factor(d_animal$Habitat_type)

#creat the data frame made of response variable, cavariables, and weights
d2 <- as.data.frame(x = y)
d2$x1 <- x1
d2$x2 <- x2
d2$x3 <- x3
d2$x4 <- x4
d2$x5 <- x5
d2$x6 <- x6
d2$w <- w

```

```

#null model
#-phylo
mnull <- gls(y ~ 1, method='ML',weights=varFixed(~w))
r <- y - fitted(mnull)
Q <- sum(w*r^2)
df <- length(r) - 1
C <- sum(w) - sum(w^2) / sum(w)
T2 <- (Q - df)/C
new_w <- 1/(1/w + T2)
mnull <- gls(y ~ 1, method='ML',weights=varFixed(~new_w))
AIC(mnull)

```

```

#model selection procedure

```

```

#first step
#model with phylogeny only
formula = y ~ 1
m <- model_fit(formula,tr_animal,d2)
AIC(m)
#model with x1
formula = y ~ -1 + x1
m_x1 <- model_fit_no_phylo(formula,d2)
AIC(m_x1)
#model with x2
formula = y ~ -1 + x2
m_x2 <- model_fit_no_phylo(formula,d2)
AIC(m_x2)
#model with x3
formula = y ~ -1 + x3
m_x3 <- model_fit_no_phylo(formula,d2)
AIC(m_x3)
#model with x4
formula = y ~ -1 + x4
m_x4 <- model_fit_no_phylo(formula,d2)
AIC(m_x4)
#model with x5
formula = y ~ -1 + x5
m_x5 <- model_fit_no_phylo(formula,d2)
AIC(m_x5)
#model with x6
formula = y ~ -1 + x6
m_x6 <- model_fit_no_phylo(formula,d2)
AIC(m_x6)

```

```
AIC(mnull,m,m_x1,m_x2,m_x3,m_x4,m_x5,m_x6)
```

```
#based on AIC, we add x5
```

```
formula = y ~ -1 + x5
```

```
m2 <- model_fit_no_phylo(formula,d2)
```

```
#second step
```

```
#with x1
```

```
formula = y ~ -1 + x5 + x1
```

```
m2_x1 <- model_fit_no_phylo(formula,d2)
```

```
#with x2
```

```
formula = y ~ -1 + x5 + x2
```

```
m2_x2 <- model_fit_no_phylo(formula,d2)
```

```
#with x3
```

```
formula = y ~ -1 + x5 + x3
```

```
m2_x3 <- model_fit_no_phylo(formula,d2)
```

```
#with x4
```

```
formula = y ~ -1 + x5 + x4
```

```
m2_x4 <- model_fit_no_phylo(formula,d2)
```

```
#with x6
```

```
formula = y ~ -1 + x5 + x6
```

```
m2_x6 <- model_fit_no_phylo(formula,d2)
```

```
#with phylo
```

```
formula = y ~ -1 + x5
```

```
m2_phylo <- model_fit(formula,tr_animal,d2)
```

```
AIC(m2,m2_x1,m2_x2,m2_x3,m2_x4,m2_x6,m2_phylo)
```

```
#based on AIC, we add the phylogeny
```

```
formula = y ~ -1 + x5
```

```
m3 <- model_fit(formula,tr_animal,d2)
```

```
#third step
```

```
#with x1
```

```
formula = y ~ -1 + x5 + x1
```

```
m3_x1 <- model_fit(formula,tr_animal,d2)
```

```
#with x2
```

```
formula = y ~ -1 + x5 + x2
```

```
m3_x2 <- model_fit(formula,tr_animal,d2)
```

```
#with x3
```

```
formula = y ~ -1 + x5 + x3
```

```
m3_x3 <- model_fit(formula,tr_animal,d2)
```

```
#with x4
```

```
formula = y ~ -1 + x5 + x4
```

```
m3_x4 <- model_fit(formula,tr_animal,d2)
#with x6
formula = y ~ -1 + x5 + x6
m3_x6 <- model_fit(formula,tr_animal,d2)

AIC(m3,m3_x1,m3_x2,m3_x3,m3_x4,m3_x6)
#based on AIC, the procedure has converged

#####
compte the delta AIC

#full model
formula = y ~ 1 + x5
mfull <- model_fit(formula,tr_animal,d2)

#-x5
formula = y ~ 1
mfull_x5 <- model_fit(formula,tr_animal,d2)

#-phylo
formula = y ~ -1 + x5
mfull_phylo <- model_fit_no_phylo(formula,d2)

AIC(mfull_phylo,mfull_x5)-AIC(mfull)
```