

1 **In Vitro Prediction of the Evolution of the GES-1 β -Lactamase**
2 **Hydrolytic Activity**

3 **S  verine Bontron,¹ Laurent Poirel^{1,2*}, and Patrice Nordmann^{1,2,3}**

4
5 ¹*Medical and Molecular Microbiology Unit, Department of Medicine, Faculty of Science,*
6 *University of Fribourg, Fribourg, Switzerland, ²INSERM U914 «Emerging Resistance to*
7 *Antibiotics», Facult   de M  decine et Universit   Paris Sud, K.-Bic  tre, France, and ³HFR-*
8 *H  pital Cantonal de Fribourg, Fribourg, Switzerland*

9
10 Running title: Directed evolution of GES-type β -lactamases

11 Keywords: GES, β -lactamase, carbapenems, cephalosporins, evolution

12
13
14
15
16 *Corresponding author. Mailing address: Medical and Molecular Microbiology
17 Unit, Department of Medicine, Faculty of Science, University of Fribourg, rue Albert
18 Gockel 3, CH-1700 Fribourg, Switzerland. Phone: 41-26-300-9582. E-mail:
19 laurent.poirel@unifr.ch

20
21

22 **Abstract**

23 **Resistance to β -lactams is constantly increasing, due to the emergence of totally new**
24 **enzymes, but also to the evolution of pre-existing β -lactamases. GES-1 is a clinically-**
25 **relevant extended-spectrum β -lactamase (ESBL) hydrolyzing penicillins and broad-**
26 **spectrum cephalosporins, but sparing monobactams and carbapenems. However,**
27 **several GES-1 variants (i.e. GES-2 and GES-5) previously identified among clinical**
28 **isolates display an extended spectrum of activity toward carbapenems. To study the**
29 **evolution potential of the GES-1 β -lactamase, this enzyme was submitted to in-vitro**
30 **directed evolution, with selection on increasing concentrations of the cephalosporin**
31 **cefotaxime, the monobactam aztreonam, or the carbapenem imipenem. The highest**
32 **resistance levels were conferred by the combination of up to four substitutions. The**
33 **A6T, E104K, G243A variant selected on cefotaxime, and the A6T, E104K, T237A,**
34 **G243A variant selected on aztreonam, conferred high resistance to cefotaxime,**
35 **ceftazidime, and aztreonam. Conversely, the A6T, G170S variant selected on imipenem**
36 **conferred high resistance to imipenem and cefoxitin. Noteworthy, the A6T substitution**
37 **involved in higher MICs for all β -lactams is located in the leader peptide of the GES**
38 **enzyme, therefore not present in the mature protein. Acquired cross resistance was not**

39 **observed since selection with CTX or ATM did not select for resistance to IPM and vice**
40 **versa. Here we demonstrated that β -lactamase GES-1 exhibits peculiar properties with a**
41 **significant potential to gain activity toward broad-spectrum cephalosporins,**
42 **monobactams, and carbapenems.**

43

44

INTRODUCTION

45

46 The main mechanism of resistance to β -lactams in Gram negatives is the production of β -
47 lactamases, which are classified into four molecular classes, namely A, B, C, and D based on
48 protein sequence analysis (1). Enzymes belonging to class A, C, and D, are serine enzymes,
49 while those belonging to class B are metallo-enzymes requiring zinc ions for activity (2, 3).

50 Over the past 70 years, since the introduction of penicillins and cephalosporins, the massive
51 use of broad-spectrum β -lactams has been at the origin of the selection of β -lactamases with
52 broadened hydrolytic activities. Emergence of those broad-spectrum β -lactamases may
53 correspond to three main phenomena; *i*) the evolution of pre-existing narrow-spectrum β -
54 lactamases with an extension of their hydrolytic profile due to key amino-acid substitutions,
55 as observed for TEM and SHV β -lactamases, *ii*) the acquisition of enzymes possessing an
56 intrinsic broad-spectrum hydrolytic activity, as observed for all CTX-M-type extended-
57 spectrum β -lactamases (ESBLs) (4, 5), and *iii*) the evolution of broad-spectrum enzymes to
58 expand or increase their hydrolytic activity to carbapenems.

59 GES-1 is an ESBL firstly identified in a *Klebsiella pneumoniae* (6), and then extensively
60 reported from clinical isolates in *Pseudomonas aeruginosa* and *Acinetobacter baumannii* (7-
61 10), but also frequently from the environmental (11-13). Similarly to other ESBLs, GES-1

62 hydrolyses penicillins and broad-spectrum cephalosporins, spares carbapenems, and is
63 susceptible to the activity of clavulanic acid as inhibitor. However, by contrast to most
64 ESBLs, GES-1 has a low activity toward the broad-spectrum cephalosporin cefotaxime
65 (CTX), and does not hydrolyze monobactams (6). The GES family comprises 24 variants
66 (GES-1 to GES-24) identified from clinical isolates, some of them having amino acid
67 substitutions conferring peculiar hydrolytic properties (Table 1). The G170N and G170S
68 substitutions (Ambler numbering [1]), located in the omega-loop of the enzyme and first
69 described in GES-2 and GES-5, confer extended activity against carbapenems and cefoxitin
70 (FOX) (this latter only for G170S), decreased hydrolysis of broad-spectrum cephalosporins,
71 and decreased susceptibility to β -lactam inhibitors (Table 1) (9, 14-16). On the other hand, the
72 E104K, G243A, and G243S substitutions, identified in several GES variants, have been
73 shown to confer higher activity toward broad-spectrum cephalosporins and the monobactam
74 aztreonam (ATM), together with an increased susceptibility to β -lactam inhibitors (Table 1).
75 Given its ability to evolve, GES-1 was chosen as a model enzyme for testing the
76 diversification potential of ESBLs. Therefore, the GES-1 enzyme was subjected to directed
77 evolution; this method consists of iterative rounds of random mutagenesis and selection, and
78 is commonly used for altering or optimizing protein function (17, 18). GES-1 was submitted

79 to three different β -lactam-based selective pressures, namely the broad-spectrum
80 cephalosporin CTX, ATM, or the carbapenem imipenem (IPM). A series of variants with
81 increased and/or broadened specificity was selected, from which the causality between the
82 increased hydrolytic activity and the mutations could be inferred, and future evolutionary
83 trajectories predicted.

84

85

MATERIALS AND METHODS

86 **Construction and selection of mutagenized GES-1 libraries.** The pBSKSII-kanR-
87 GES-1 plasmid was used as the reference plasmid coding for GES-1. Plasmid pBSKSII-kanR
88 is derived from pBluescriptII (a high copy number plasmid, 500 copies per cell), and encodes
89 resistance to kanamycin. The entire *bla*_{GES-1} coding region (6) was amplified, the amplicon
90 including 31 bp upstream of the ATG start, with primers No50 (GES-1-*HindIII*-F, 5'-
91 gatgatAAGCTTACAAAGATAATTTCCATCTCAAGG-3') and No51 (GES-1-*NotI*-R, 5'-
92 gatgatGCGGCCGCCTATTTGTCCGTGCTCAGGATG-3'), and cloned into the *HindIII*/*NotI*
93 restriction sites of pBSKSII-kanR. The construct was verified by sequencing. Random
94 mutagenesis was performed with the GenMorph II Random Mutagenesis Kit (Agilent
95 Technologies, Santa Clara, CA), with primers No50 and No51, following the manufacturer

96 recommendations. The PCR amplified mutagenized product was purified, digested, and
97 ligated into the HindIII/NotI restriction sites of pBSKSII-kanR. After purification, the ligation
98 mixture was transformed into TOP10 electro-competent *E. coli* cells (Life Technologies, Zug,
99 Switzerland). The library was plated on Luria broth plates supplemented with 25 µg/ml of
100 kanamycin. At each round, the complexity of the library was at least 10⁵ independent clones.
101 The mean substitution rate was of 2 nucleotides per molecule, based on sequencing of 10
102 clones. This corresponds to at least 6x10⁴ distinct sequences, as calculated with the library
103 statistics program PEDEL (19). For selection, the libraries were plated on increasing doses of
104 the indicated antibiotic, with a 2-fold increment. Plasmids recovered from the clones obtained
105 with the highest antibiotic concentrations (usually 100 to 300 clones) were isolated, re-
106 transformed, and plated again on the same antibiotic concentration. At this step, at least 4
107 clones were analyzed by sequencing, and the rest of the clones was isolated as a pool, and
108 used as a basis for the next round of random mutagenesis. Constructs with different
109 combinations of mutations were made by sub-cloning, or with the Q5 Site-Directed
110 Mutagenesis Kit (New England BioLabs, Ipswich, MA).

111 **MICs, specific hydrolytic activities, and IC₅₀s measurements.** MICs were measured
112 by Etest (bioMérieux, Marcy l'Etoile, France). Specific hydrolytic activities were measured

113 from whole cell extracts of recombinant *E. coli* strains producing the different GES variants
114 as described (6). Assays were performed in 500 μ l total volume, using 150 μ M (for
115 benzylpenicillin [PEN], ATM, IPM, FOX) or 75 μ M (for CTX and ceftazidime [CAZ]) of
116 substrate. Hydrolysis measurements were performed at room temperature with a JENWAY
117 spectrophotometer (Staffordshire, UK). The protein concentrations were measured with
118 Bradford Reagent (Sigma-Aldrich, Buchs, Switzerland). The results were expressed in nmoles
119 $\times \text{min}^{-1} \times \mu\text{g}^{-1}$ extract. Experiments were made in triplicates from 3 independent cultures.
120 Inhibitory concentrations inhibiting 50% of the hydrolysis activity (IC_{50}) were measured for
121 clavulanic acid using PEN as substrate. A 3-min long preincubation step with clavulanic acid
122 was used before adding PEN. Those experiments were performed in triplicates with three
123 independent cultures.

124

125

RESULTS

126 **Selection of GES variants conferring increased resistance to CTX, ATM, and**
127 **IPM by directed evolution.** GES mutants conferring increased resistance to CTX, ATM, or
128 IPM were recovered after three (for ATM and IPM) or four (for CTX) rounds of mutagenesis
129 and selection. Over the rounds the variants were successively selected with 1, 2, 4, and

130 16 µg/ml of CTX, with 1, 16, 128, and 256 µg/ml of ATM, or with 0.1, 0.125, and 0.25 µg/ml
131 of IPM. Additional mutagenesis steps could not select variants with higher MICs. At the end
132 of each round, at least four clones were sequenced. Mutations common to several clones were
133 anticipated to be the phenotypically relevant ones, but additional substitutions, silent or not,
134 did accumulate throughout the mutagenesis rounds. Therefore, some constructs were
135 specifically generated to definitely correlate amino acid substitutions to resistance phenotypes
136 (data not shown). The corresponding so-called cured variants (GES-C1 to -C5 for selection
137 with CTX, GES-A1 to -A5 for selection with ATM, and GES-I1 to -I3 for selection with
138 IPM) harboring the corresponding amino acid changes are depicted in Table 2A-C. The
139 sequences of the originally isolated clones are listed in Table S1.

140 **MICs of β -lactams for the selected mutants.** Upon selection with CTX, the A6T
141 substitution (clone GES-C1), located in the signal peptide, was the only mutation selected
142 after round 1 (Table 2A). At round 2, substitutions E104K or G243S were added to A6T
143 (clones GES-C2 and GES-C3), and GES-C4 (A6T, G243A) was selected at round 3.
144 Combination of substitutions A6T, E104K, G243A was selected at round 4 (clone GES-C5).
145 MICs of CTX for these constructs gradually increased during the directed evolution, from
146 0.75 µg/ml for GES-1 to 48 µg/ml for GES-C5 (Table 2A). MICs of ATM and ceftazidime

147 (CAZ) increased concomitantly to those of CTX, while MICs of FOX, IPM, and the
148 carbapenem ertapenem (ETP) remained unchanged (Table 2A). Four additional constructs
149 were made to dissect the role of each amino acid change. Single E104K, G243S, or G243A
150 substitutions (clones GES-C6, GES-C7, and GES-C8, respectively) conferred a lower
151 resistance than A6T (clone GES-C1). Combination of E104K and G243A (clone GES-C9)
152 was slightly more efficient than single mutations (Table 2A). Clone GES-C5 exhibiting
153 substitutions A6T, E104K, and G243A combined three changes that individually conferred
154 modest MIC increases, but when combined together resulted into a variant for which the MIC
155 of CTX reached 48 $\mu\text{g/ml}$.

156 When selecting with ATM, single E104K (clone GES-A1) or G243A (clone GES-A2)
157 mutations were selected in round 1, while a combination of those two was selected at round 2
158 (clone GES-A3). Ultimately, clones selected on ATM-128 additionally harbored a T237A
159 substitution (E104K, T237A, G243A, clone GES-A4), and those selected on ATM-256
160 harbored the A6T substitution in addition to the three other changes (clone GES-A5) (Table
161 2B). MICs of ATM for these constructs increased during the directed evolution, from 0.25
162 $\mu\text{g/ml}$ for GES-1 to $>256 \mu\text{g/ml}$ for clones GES-A4 and GES-A5. In parallel MICs of CAZ
163 increased sharply, while those of CTX more modestly. No change in MICs of FOX, IPM, or

164 ETP was observed (Table 2B). Each of the three E104K, T237A, or G243A substitutions
165 (clones GES-C6, GES-A6, and GES-A2, respectively) had slightly increased MICs of ATM
166 (from 1.5 to 3 µg/ml). Any dual combination resulted into higher MICs of ATM (from 12 or
167 32 µg/ml, clones GES-A3, -A7 and -A8), while the triple mutant had an MIC of ATM of
168 >256 µg/ml (clone GES-A4) (Table 2B). Of note, while the T237A substitution correlated
169 with increased MICs of ATM and CAZ, it was systematically deleterious for the MIC of
170 CTX.

171 When selecting with IPM, substitution G170S was selected at round 1 (clone GES-I1). Then
172 substitution c-1t lying 1 bp before the ATG start codon (clone GES-I2) and finally
173 substitution A6T (clone GES-I3) were selected at rounds 2 and 3 (Table 2C). Overall, MICs
174 of IPM increased from 0.25 µg/ml for wild-type GES-1 to 2 µg/ml for clone GES-I3. MICs of
175 ETP and FOX increased concomitantly but conversely, MICs of ATM, CTX, and CAZ were
176 lowered once the G170S mutation was selected (Table 2C).

177 **In-vitro specific hydrolytic activities of the GES variants correlating with MICs**
178 **values.** In order to confirm that the higher MICs observed for the selected clones were indeed
179 related to higher catalytic activities or higher amount of the different GES enzymes selected,
180 and not to a non-enzymatic resistance mechanism, in-vitro specific hydrolytic activities of a

181 set of GES variants were determined. Overall, for all enzymes the increased hydrolytic
182 activities toward CTX, CAZ, ATM, IPM, and FOX correlated with the higher MICs of the
183 corresponding substrates (Tables 2A-C).

184 **Differential inhibition of the GES selected variants by clavulanic acid.**

185 Determination of IC_{50} s of clavulanic acid was performed for a representative set of the GES
186 variants. Clones GES-C1 and GES-C5 showed similar IC_{50} values of clavulanic acid
187 compared to the wild-type GES-1 (Table 2A). However, clones GES-A4 and GES-A5 showed
188 10-fold lower IC_{50} values (0.6 μ M) (Table 2B) likely due to the T237A substitution.
189 Conversely clones GES-I3 and GES-I4 selected on IPM and harboring the G170S substitution
190 showed significantly higher IC_{50} values (Table 2C). Overall, mutants selected on CTX
191 showed similar susceptibility to clavulanic acid as GES-1, while clones selected on ATM
192 showed increased, and clones selected on IPM lower susceptibility to clavulanic acid.

193

194 **DISCUSSION**

195 The directed evolution procedure used here combined the generation of mutagenized libraries
196 with antibiotic selection, and allowed the selection of GES variants that may be categorized
197 into two classes, being those selected on CTX or ATM conferring high resistance to CTX,

198 ATM, and CAZ on one hand, and those selected on IPM conferring high resistance to IPM
199 and FOX on the other hand.

200 Interestingly, the A6T substitution located into the signal sequence (18 amino-acid long for
201 GES-1) was selected through all three evolutionary routes. Noteworthy, signal sequences are
202 required for translocation to the periplasm (20). The A6T substitution in GES-1 systematically
203 conferred a 3- to 8-fold increase in the MICs of CTX, ATM, and IPM (Table 2A-C). In
204 accordance with our observation, a study including experimental mutagenesis of a consensus
205 signal sequence fused to a β -lactamase gene increased ampicillin tolerance level of the host
206 cell up to 8-fold (21). To the best of our knowledge, it has never been clearly assessed that
207 mutations in the signal sequence of any β -lactamase may confer increased resistance to β -
208 lactams. The impact of such substitution should therefore be further investigated as a
209 mechanism leading to reduced susceptibility or even resistance to β -lactams.

210 By selecting with IPM, the G170S mutation located in the Ω -loop of the catalytic site (amino
211 acids 159 to 182), a highly conserved motif among the β -lactamases, was recovered. This
212 substitution was previously shown to confer a 100-fold increased catalytic activity against
213 IPM when compared to GES-1 (14, 22, 23). The E104 residue is exposed near the entrance to
214 the binding site, and the E104K substitution is commonly found in the TEM family, where

215 this change participates to the expansion of the β -lactamase spectrum, more strikingly when
216 associated with other substitutions such as R164S or G238S (24-26). Despite several studies,
217 the mechanism of this synergism remains poorly understood (26). The G243A is not
218 conserved among β -lactamases and this change might create subtle rearrangements in the
219 disulfide bond. The T237 amino acid, together with the S70 residue, forms an oxyanion hole,
220 which houses the β -lactam carbonyl of the acyl-enzymes intermediate (27). Position 237,
221 usually occupied by an Ala or Ser in most class A β -lactamases, corresponds to a Thr residue
222 in GES-1, but also in the PER-1 ESBL and in the class A KPC-2 carbapenemase. It was
223 experimentally shown with KPC-2 that a T237A change resulted into lower hydrolysis of
224 CTX (28). Conversely in TEM, the natural or experimentally generated A237T substitution
225 confers an increased hydrolysis of CTX and a decreased hydrolysis of CAZ and ATM (22,
226 30). These observations correlate with the detrimental effect of the T237A substitution in
227 GES for the hydrolysis of CTX, and the beneficial effect for the hydrolysis of CAZ and ATM.
228 Substitutions E104K, G170S, T237A, G243S, and G243A selected through our study were
229 previously described in GES alleles identified from clinical isolates (Table 1). Noticeably,
230 substitution G170S, increasing carbapenem hydrolysis, was previously identified in the
231 natural carbapenemase GES-5. Substitutions E104K, T237A, and G243A were also identified

232 in natural GES alleles, either alone (GES-9, G243A), associated to phenotypically
233 uncharacterized mutations (GES-3, GES-7, GES-19, GES-22), or combined with other
234 substitutions (GES-12 and GES-17) (Table 1). As inferred from our in-vitro directed
235 evolution experiments, GES-5, GES-12 and GES-17 may therefore be prone to evolve into
236 more active variants. On the other hand, some GES alleles recovered from clinical isolates
237 combined the G170S mutation conferring higher activity toward carbapenems, together with
238 another mutation conferring increased activity toward CTX, ATM, and CAZ (see GES-6,
239 E104K, G170S, and GES-14, G243A, G170S). Comparative studies showed that the
240 increased hydrolysis of CTX, CAZ, or ATM mediated by E104K or G243A was abolished by
241 the additional presence of G170S (29, 31, 32). This precludes that the natural alleles GES-6
242 and GES-14 have been selected under successive distinct selective pressures. Similarly,
243 detailed analysis of the CTX-M-type ESBL potential evolutionary trajectories showed that the
244 diversification process of the CTX-M variants could only be explained by a selection with at
245 least two antibiotics (33).

246 Our study showed that GES enzymes can evolve into two types of variants conferring higher
247 resistance to CTX, ATM, or to IPM. Spontaneous evolution of antibiotic resistance is a
248 multifactorial phenomenon, given the diversity of the genetic support of resistance genes, of

249 the bacterial strain (including potential changes of membrane permeability or of the
250 penicillin-binding proteins), and of the nature and the concentration of the antibiotic. As a
251 consequence, a higher number of possible mutations and evolutionary trajectories are
252 possible, although constrained by intramolecular interactions (34). Nevertheless, a good
253 correlation between in-vitro prediction inferred from the analysis of the selected variants, and
254 those found in clinical isolates has been established here. Predictions regarding the occurrence
255 of very efficient natural variants in term of catalytic efficiency using an in-vitro directed
256 evolution was previously demonstrated for the TEM β -lactamase (34-37). From our study we
257 may speculate that selection with cephalosporins or monobactam might not select for GES
258 variants possessing carbapenemase activity, and conversely selection with the carbapenem
259 IPM might not select for GES variants possessing increased hydrolytic activity toward CTX,
260 CAZ, or ATM. Interestingly, as previously noticed with CTX-M-type ESBLs possessing
261 increased activity toward broad-spectrum cephalosporins (38), some antagonistic pleiotropy
262 was observed, such as a decreased susceptibility to β -lactamase inhibitors of some GES
263 variants that exhibited increased catalytic activity toward carbapenems. Overall, such an
264 approach sheds light on how clinical alleles have been selected, and might predict the future
265 evolutionary trajectories of the β -lactamases according to the antibiotic selection pressure.

266

267

ACKNOWLEDGEMENTS

268 This work was supported by the University of Fribourg, Fribourg, Switzerland.

269

270

REFERENCES

- 271 1. **Ambler RP, Coulson AF, Frère JM, Ghuysen JM, Joris B, Forsman M, Lévesque**
272 **RC, Tiraby G, Waley SG.** 1991. A standard numbering scheme for the class A β -
273 lactamases. *Biochem. J.* **276**:269-270.
- 274 2. **Bush K.** 2013. The ABCD's of beta-lactamase nomenclature. *J. Infect. Chemother.*
275 **19**:549-559.
- 276 3. **Bush K.** 2010. Alarming β -lactamase-mediated resistance in multidrug-resistant
277 Enterobacteriaceae. *Curr. Opin. Microbiol.* **13**:558-564.
- 278 4. **Gutkind GO, Di Conza J, Power P, Radice M.** 2013. β -lactamase-mediated
279 resistance: a biochemical, epidemiological and genetic overview. *Curr. Pharm. Des.*
280 **19**:164-208.
- 281 5. **Naas T, Poirel L, Nordmann P.** 2008. Minor extended-spectrum β -lactamases. *Clin.*
282 *Microbiol. Infect.* **14 Suppl 1**:42-52.
- 283 6. **Poirel L, Le Thomas I, Naas T, Karim A, Nordmann P.** 2000. Biochemical
284 sequence analyses of GES-1, a novel class A extended-spectrum β -lactamase, and the
285 class 1 integron In52 from *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.*
286 **44**:622-632.

- 287 7. **Poirel L, Bonnin RA, Nordmann P.** 2012. Genetic support and diversity of acquired
288 extended-spectrum β -lactamases in Gram-negative rods. *Infect. Genet. Evol.* **12**:883-
289 893.
- 290 8. **Poirel L, Brinas L, Fortineau N, Nordmann P.** 2005. Integron-encoded GES-type
291 extended-spectrum β -lactamase with increased activity toward aztreonam in
292 *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **49**:3593-3597.
- 293 9. **Poirel L, Weldhagen GF, Naas T, De Champs C, Dove MG, Nordmann P.** 2001.
294 GES-2, a class A β -lactamase from *Pseudomonas aeruginosa* with increased
295 hydrolysis of imipenem. *Antimicrob. Agents Chemother.* **45**:2598-2603.
- 296 10. **Bonnin RA, Nordmann P, Potron A, Lecuyer H, Zahar JR, Poirel L.** 2011.
297 Carbapenem-hydrolyzing GES-type extended-spectrum β -lactamase in *Acinetobacter*
298 *baumannii*. *Antimicrob. Agents Chemother.* **55**:349-354.
- 299 11. **Girlich D, Poirel L, Szczepanowski R, Schluter A, Nordmann P.** 2012.
300 Carbapenem-hydrolyzing GES-5-encoding gene on different plasmid types recovered
301 from a bacterial community in a sewage treatment plant. *Appl. Environ. Microbiol.*
302 **78**:1292-1295.
- 303 12. **Manageiro V, Ferreira E, Canica M, Manaia CM.** 2014. GES-5 among the β -
304 lactamases detected in ubiquitous bacteria isolated from aquatic environment samples.
305 *FEMS Microbiol. Lett.* **351**:64-69.
- 306 13. **Ribeiro VB, Zavascki AP, Rozales FP, Pagano M, Magagnin CM, Nodari CS, da**
307 **Silva RC, Dalarosa MG, Falci DR, Barth AL.** 2014. Detection of *bla*_{GES-5} in
308 carbapenem-resistant *Kluyvera intermedia* isolates recovered from the hospital
309 environment. *Antimicrob. Agents Chemother.* **58**:622-623.

- 310 14. **Smith CA, Caccamo M, Kantardjieff KA, Vakulenko S.** 2007. Structure of GES-1
311 at atomic resolution: insights into the evolution of carbapenamase activity in the class
312 A extended-spectrum β -lactamases. *Acta Crystallogr. D. Biol. Crystallogr.* **63**:982-
313 992.
- 314 15. **Picao RC, Poirel L, Gales AC, Nordmann P.** 2009. Diversity of β -lactamases
315 produced by ceftazidime-resistant *Pseudomonas aeruginosa* isolates causing
316 bloodstream infections in Brazil. *Antimicrob. Agents Chemother.* **53**:3908-3913.
- 317 16. **Bae IK, Lee YN, Jeong SH, Hong SG, Lee JH, Lee SH, Kim HJ, Youn H.** 2007.
318 Genetic and biochemical characterization of GES-5, an extended-spectrum class A β -
319 lactamase from *Klebsiella pneumoniae*. *Diagn. Microbiol. Infect. Dis.* **58**:465-468.
- 320 17. **Goldsmith M, Tawfik DS.** 2012. Directed enzyme evolution: beyond the low-
321 hanging fruit. *Curr. Opin. Struct. Biol.* **22**:406-412.
- 322 18. **Romero PA, Arnold FH.** 2009. Exploring protein fitness landscapes by directed
323 evolution. *Nat. Rev. Mol. Cell. Biol.* **10**:866-876.
- 324 19. **Patrick WM, Firth AE, Blackburn JM.** 2003. User-friendly algorithms for
325 estimating completeness and diversity in randomized protein-encoding libraries.
326 *Protein Eng.* **16**:451-457.
- 327 20. **Izard JW, Kendall DA.** 1994. Signal peptides: exquisitely designed transport
328 promoters. *Mol. Microbiol.* **13**:765-773.
- 329 21. **Heggeset TM, Kucharova V, Naerdal I, Valla S, Sletta H, Ellingsen TE, Brautaset**
330 **T.** 2013. Combinatorial mutagenesis and selection of improved signal sequences and

- 331 their application for high-level production of translocated heterologous proteins in
332 *Escherichia coli*. Appl. Environ. Microbiol. **79**:559-568.
- 333 22. **Frase H, Shi Q, Testero SA, Mobashery S, Vakulenko SB.** 2009. Mechanistic basis
334 for the emergence of catalytic competence against carbapenem antibiotics by the GES
335 family of β -lactamases. J. Biol. Chem. **284**:29509-29513.
- 336 23. **Smith CA, Frase H, Toth M, Kumarasiri M, Wiafe K, Munoz J, Mobashery S,**
337 **Vakulenko SB.** 2012. Structural basis for progression toward the carbapenemase
338 activity in the GES family of β -lactamases. J. Am. Chem. Soc. **134**:19512-19515.
- 339 24. **Gniadkowski M.** 2008. Evolution of extended-spectrum beta-lactamases by mutation.
340 Clin. Microbiol. Infect. **1**:11-32.
- 341 25. **Page MG.** 2008. Extended-spectrum beta-lactamases: structure and kinetic
342 mechanism. Clin. Microbiol. Infect. **1**:64-74.
- 343 26. **Salverda ML, De Visser JA, Barlow M.** 2010. Natural evolution of TEM-1 β -
344 lactamase: experimental reconstruction and clinical relevance. FEMS Microbiol. Rev.
345 **34**:1015-1036.
- 346 27. **Murphy BP, Pratt RF.** 1988. Evidence for an oxyanion hole in serine β -lactamases
347 and DD-peptidases. Biochem. J. **256**:669-672.
- 348 28. **Papp-Wallace KM, Taracila M, Hornick JM, Hujer AM, Hujer KM, Distler AM,**
349 **Endimiani A, Bonomo RA.** 2010. Substrate selectivity and a novel role in inhibitor
350 discrimination by residue 237 in the KPC-2 β -lactamase. Antimicrob. Agents
351 Chemother. **54**:2867-2877.

- 352 29. **Kotsakis SD, Miriagou V, Tzelepi E, Tzouvelekis LS.** 2010. Comparative
353 biochemical and computational study of the role of naturally occurring mutations at
354 Ambler positions 104 and 170 in GES β -lactamases. *Antimicrob. Agents Chemother.*
355 **54**:4864-4871.
- 356 30. **Giakkoupi P, Hujer AM, Miriagou V, Tzelepi E, Bonomo RA, Tzouvelekis LS.**
357 2001. Substitution of Thr for Ala-237 in TEM-17, TEM-12 and TEM-26: alterations
358 in β -lactam resistance conferred on *Escherichia coli*. *FEMS Microbiol. Lett.* **201**:37-
359 40.
- 360 31. **Delbruck H, Bogaerts P, Kupper MB, Rezende de Castro R, Bennink S,**
361 **Glupczynski Y, Galleni M, Hoffmann KM, Bebrone C.** 2012. Kinetic and
362 crystallographic studies of extended-spectrum GES-11, GES-12, and GES-14 β -
363 lactamases. *Antimicrob. Agents Chemother.* **56**:5618-5625.
- 364 32. **Bogaerts P, Naas T, El Garch F, Cuzon G, Deplano A, Delaire T, Huang TD,**
365 **Lissoir B, Nordmann P, Glupczynski Y.** 2010. GES extended-spectrum β -
366 lactamases in *Acinetobacter baumannii* isolates in Belgium. *Antimicrob. Agents*
367 *Chemother.* **54**:4872-4878.
- 368 33. **Novais A, Comas I, Baquero F, Canton R, Coque TM, Moya A, Gonzalez-**
369 **Candelas F, Galan JC.** 2010. Evolutionary trajectories of β -lactamase CTX-M-1
370 cluster enzymes: predicting antibiotic resistance. *PLoS Pathog.* **6**:e1000735.
- 371 34. **Weinreich DM, Delaney NF, Depristo MA, Hartl DL.** 2006. Darwinian evolution
372 can follow only very few mutational paths to fitter proteins. *Science* **312**:111-114.

- 373 35. **Stemmer WP.** 1994. DNA shuffling by random fragmentation and reassembly: in
374 vitro recombination for molecular evolution. Proc. Natl. Acad. Sci. U.S.A. **91**:10747-
375 10751.
- 376 36. **Zaccolo M, Gherardi E.** 1999. The effect of high-frequency random mutagenesis on
377 in vitro protein evolution: a study on TEM-1 β -lactamase. J. Mol. Biol. **285**:775-783.
- 378 37. **Orencia MC, Yoon JS, Ness JE, Stemmer WP, Stevens RC.** 2001. Predicting the
379 emergence of antibiotic resistance by directed evolution and structural analysis. Nat.
380 Struct. Biol. **8**:238-242.
- 381 38. **Ripoll A, Baquero F, Novais A, Rodríguez-Domínguez MJ, Turrientes MC,**
382 **Cantón R, Galán JC.** 2011. In vitro selection of variants resistant to β -lactams plus β -
383 lactamase inhibitors in CTX-M β -lactamases: predicting the *in vivo* scenario?
384 Antimicrob. Agents Chemother. **55**:4530-4536.
- 385 39. **Wachino J, Doi Y, Yamane K, Shibata N, Yagi T, Kubota T, Ito H, Arakawa Y.**
386 2004. Nosocomial spread of ceftazidime-resistant *Klebsiella pneumoniae* strains
387 producing a novel class a β -lactamase, GES-3, in a neonatal intensive care unit in
388 Japan. Antimicrob. Agents Chemother. **48**:1960-1967.
- 389 40. **Bebrone C, Bogaerts P, Delbruck H, Bennink S, Kupper MB, Rezende de Castro**
390 **R, Glupczynski Y, Hoffmann KM.** 2013. GES-18, a new carbapenem-hydrolyzing
391 GES-Type β -lactamase from *Pseudomonas aeruginosa* that contains Ile80 and Ser170
392 residues. Antimicrob. Agents Chemother. **57**:396-401.
- 393 41. **Giakkoupi P, Tzouveleakis LS, Tsakris A, Loukova V, Sofianou D, Tzelepi E.**
394 2000. IBC-1, a novel integron-associated class A β -lactamase with extended-spectrum

- 395 properties produced by an *Enterobacter cloacae* clinical strain. Antimicrob. Agents
396 Chemother. **44**:2247-2253.
- 397 42. **Mavroidi A, Tzelepi E, Tsakris A, Miriagou V, Sofianou D, Tzouvelekis LS.** 2001.
398 An integron-associated β -lactamase (IBC-2) from *Pseudomonas aeruginosa* is a
399 variant of the extended-spectrum β -lactamase IBC-1. J. Antimicrob. Chemother.
400 **48**:627-630.
- 401 43. **Moubareck C, Bremont S, Conroy MC, Courvalin P, Lambert T.** 2009. GES-11, a
402 novel integron-associated GES variant in *Acinetobacter baumannii*. Antimicrob.
403 Agents Chemother. **53**:3579-3581.
- 404 44. **Kotsakis SD, Papagiannitsis CC, Tzelepi E, Legakis NJ, Miriagou V, Tzouvelekis**
405 **LS.** 2010. GES-13, a β -lactamase variant possessing Lys-104 and Asn-170 in
406 *Pseudomonas aeruginosa*. Antimicrob. Agents Chemother. **54**:1331-1333.
- 407

Table 1. GES variants isolated from clinical isolates, in relation to their hydrolysis profile

Variant ^a	Amino acid position ^b														Hydrolysis profile ^c					References	
	11	12	44	55	62	80	81	104	125	130	167	169	170	237	243	CTX	ATM	CAZ	IPM		FOX
GES-1	G	I	Q	I	M	V	F	E	A	S	P	M	G	T	G	+	-	++	-	-	(6)
GES-2													N		+/-	-	+	+	-	(9, 29)	
GES-3					T		K								+	+	++	-	-	(39)	
GES-4					T		K						S		+	+	++	+	+	(39)	
GES-5													S		+/-	-	+	+	+	(29, 32, 40)	
GES-6							K						S		+/-	-	++	+	+	(29)	
GES-7							K			L					+	++	+++	-	-	(29, 41, 42)	
GES-8										L					+	-	++	-	-	(42)	
GES-9													S		++	+	+++	-	-	(8, 32)	
GES-10		T			T						C										
GES-11													A		++	+	+++	-	-	(31, 32, 43)	
GES-12													A	A	++	+	+++	-	-	(31, 32)	
GES-13							K						N		+	+	++	-	-	(44)	
GES-14											S		S	A	+/-	-	++	+	+	(31, 32)	
GES-15													S								
GES-16			E										S								
GES-17								K						A							
GES-18						I							S		-	-	+	+	+	(40)	
GES-19	A													A							
GES-20	A												S								
GES-21							L						S								
GES-22											L			A							
GES-23				L																	
GES-24					T								S								

^a GES-1 to GES-24 clinical variants are listed according to <http://lahey.org/studies/other.asp>.

^b Amino acid positions were assigned according to Ambler. Amino acid changes as compared to wild type GES-1 are indicated.

^c When available, the hydrolysis profile of each variant was estimated from published MICs: + and -, hydrolysis and no hydrolysis, respectively, with dark grey related to an increased hydrolysis, and light grey to a decreased hydrolysis, as compared to wild type GES-1. For each variant, the amino acid substitution responsible for the change in the hydrolytic profile is framed. Of note, these comparisons are indicative, given the bacterial species and the genetic support differ according to the study.

Table 2. Selected GES variants, related MICs and in vitro specific hydrolytic activities of CTX, ATM, CAZ, IPM, ETP, FOX, and IC₅₀ of clavulanic acid

Table 2A. Clones selected on CTX

Variant ^a	Amino acid position ^b			MIC ^c (µg/ml)						In vitro specific hydrolytic activity ^d (nmoles x min ⁻¹ x µg ⁻¹ extract)						IC ₅₀ clav. ac. (µM)	
	6	104	243	CTX	ATM	CAZ	IPM	ETP	FOX	PEN	CTX	ATM	CAZ	IPM	FOX		
Wild-type	GES-1	A	E	G	0.75	0.25	12	0.25	0.006	4	3.4±0.6	1.0±0.2	<0.1	<0.1	<0.1	<0.1	7.7±0.5
Round 1	GES-C1	T			4	1.5	128	0.38	0.016	6	9.1±0.6	2.8±0.5	<0.1	<0.1	<0.1	<0.1	10.3±3.8
Round 2	GES-C2	T	K		8	16	>256	0.38	0.016	6	-	-	-	-	-	-	-
	GES-C3	T		S	8	12	>256	0.38	0.016	3	-	-	-	-	-	-	-
Round 3	GES-C4	T		A	16	24	>256	0.38	0.032	3	5.2±0.9	12.1±1.0	1.8±0.7	1.5±0.3	<0.1	<0.1	-
Round 4	GES-C5	T	K	A	48	>256	>256	0.38	0.032	4	4.6±0.6	17.4±1.6	24.2±2.8	6.6±0.6	<0.1	<0.1	5.3±0.1
Constructs	GES-C6		K		1.5	3	>256	0.25	0.008	4	-	-	-	-	-	-	-
	GES-C7			S	1.5	2	48	0.25	0.008	4	-	-	-	-	-	-	-
	GES-C8			A	3	3	128	0.38	0.012	3	-	-	-	-	-	-	-
	GES-C9		K	A	4	32	>256	0.25	0.012	4	-	-	-	-	-	-	-

Table 2B. Clones selected on ATM

Variant ^a	Amino acid position ^b				MIC ^c (µg/ml)						In vitro specific hydrolytic activity ^d (nmoles x min ⁻¹ x µg ⁻¹ extract)						IC ₅₀ clav. ac. (µM)	
	6	104	237	243	CTX	ATM	CAZ	IPM	ETP	FOX	PEN	CTX	ATM	CAZ	IPM	FOX		
Wild-type	A	E	T	G	0.75	0.25	12	0.25	0.006	4	3.4±0.6	1.0±0.2	<0.1	<0.1	<0.1	<0.1	7.7±0.5	
GES-1		K			1.5	3	>256	0.25	0.008	4	-	-	-	-	-	-	-	
Round 1				A	3	3	128	0.38	0.012	3	-	-	-	-	-	-	-	
GES-A2					4	32	>256	0.25	0.012	4	1.3±0.2	4.1±0.6	4.7±0.5	1.4±0.3	<0.1	<0.1	-	
GES-A3		K	A		4	32	>256	0.25	0.012	4	1.3±0.2	4.1±0.6	4.7±0.5	1.4±0.3	<0.1	<0.1	-	
Round 2		K	A	A	2	>256	>256	0.38	0.025	4	8.7±2.3	4.5±1.1	14.2±3.0	4.0±0.6	<0.1	<0.1	0.6±0.07	
GES-A4		T	K	A	A	12	>256	>256	0.25	0.064	6	23.6±2.5	12.2±1.2	38.0±2.5	10.9±0.7	<0.1	<0.1	0.5±0.04
Round 4					0.38	1.5	48	0.25	0.012	3	-	-	-	-	-	-	-	
GES-A6				A	0.75	32	>256	0.25	0.012	4	-	-	-	-	-	-	-	
Conducts		K	A		1	12	>256	0.38	0.025	4	-	-	-	-	-	-	-	
GES-A7				A	1	12	>256	0.38	0.025	4	-	-	-	-	-	-	-	
GES-A8					1	12	>256	0.38	0.025	4	-	-	-	-	-	-	-	

Table 2C. Clones selected on IPM

Variant ^a	Amino acid position ^b			MIC (µg/ml) ^c						In vitro specific hydrolytic activity (nmoles x min ⁻¹ x µg ⁻¹ extract) ^d						IC ₅₀ clav. ac. (µM) ^e	
	nt-1	6	170	CTX	ATM	CAZ	IPM	ETP	FOX	PEN	CTX	ATM	CAZ	IPM	FOX		
Wild-type	GES-1	c	A	G	0.75	0.25	12	0.25	0.006	4	3.4±0.6	1.0±0.2	<0.1	<0.1	<0.1	7.7±0.5	
Round 1	GES-11			S	0.125	0.094	1.5	0.5	0.064	12	7.7±1.2	<0.1	<0.1	<0.1	0.15±0.0006	0.16±0.04	86±24
Round 2	GES-12	t		S	0.19	0.125	2	0.75	0.094	24	12.4±3.2	<0.1	<0.1	<0.1	0.18±0.02	0.27±0.06	-
Round 3	GES-13	t	T	S	1	0.19	12	2	0.25	>256	37.8±2.1	<0.1	<0.1	<0.1	0.64±0.04	0.97±0.13	150±12
Constructs	GES-14		T		4	1.5	128	0.38	0.016	6	-	-	-	-	-	-	-
	GES-15		T	S	0.5	0.19	4	0.75	0.19	48	-	-	-	-	-	-	-

^a Variants selected with CTX and derivative constructs are designated with a "C", those on ATM with an "A", and those on IPM with an "I". The round of directed evolution on which the variants were selected is indicated on the left. Antibiotic concentrations used for selection were the following: on CTX, round 1: 1µg/ml, round 2: 2µg/ml, round 3: 4µg/ml, round 4: 16µg/ml; on ATM, round 1: 1µg/ml, round 2: 16µg/ml, round 3: 128µg/ml, round 4: 256µg/ml; on IPM, round 1: 0.1µg/ml, round 2: 0.125µg/ml, round 3: 0.25µg/ml.

^b Amino acid positions were assigned according to Ambler, except for mutations before the ATG start codon, where the nucleotide number relative to it is indicated. Amino acid changes as compared to wild type GES-1 are indicated.

^c All the GES alleles are expressed in the highly susceptible *E. coli* TOP10.

^d Specific activity values were measured by U.V. spectrophotometry from crude extracts of *E. coli* TOP10 producing the indicated variant, for each of the indicated antibiotics (PEN, CTX, CAZ, ATM, FOX, and IPM). The mean and the S.D. are indicated. (-) not determined.

^e IC₅₀ of clavulanic acid were measured with PEN as a substrate. The mean and the S.D. are indicated.