

Emerging broad-spectrum resistance in *Pseudomonas aeruginosa* and *Acinetobacter baumannii*: Mechanisms and epidemiology

Anaïs Potron^a, Laurent Poirel^{b,*}, Patrice Nordmann^{b,c}

^a Laboratoire de Bactériologie, Faculté de Médecine-Pharmacie, Centre Hospitalier Régional Universitaire, Université de Franche-Comté, Besançon, France

^b Emerging Antibiotic Resistance Medical and Molecular Microbiology Unit, Department of Medicine, Faculty of Science, University of Fribourg, Fribourg, Switzerland

^c HFR – Hôpital Cantonal de Fribourg, Fribourg, Switzerland

Multidrug resistance is quite common among non-fermenting Gram-negative rods, in particular among clinically relevant species including *Pseudomonas aeruginosa* and *Acinetobacter baumannii*. These bacterial species, which are mainly nosocomial pathogens, possess a diversity of resistance mechanisms that may lead to multidrug or even pandrug resistance. Extended-spectrum β -lactamases (ESBLs) conferring resistance to broad-spectrum cephalosporins, carbapenemases conferring resistance to carbapenems, and 16S rRNA methylases conferring resistance to all clinically relevant aminoglycosides are the most important causes of concern. Concomitant resistance to fluoroquinolones, polymyxins (colistin) and tigecycline may lead to pandrug resistance. The most important mechanisms of resistance in *P. aeruginosa* and *A. baumannii* and their most recent dissemination worldwide are detailed here.

1. Introduction

The emergence and spread of bacteria resistant to multiple antibiotics and at the origin of severe infections is currently of great concern. This is particularly true for nosocomial pathogens isolated in hospitals, where these superbugs may compromise advanced medicine, including surgery, transplantation, efficient treatment of immunocompromised and haematological patients, etc. Among the increasingly reported and commonly identified multidrug-resistant or even pandrug-resistant bacteria, the lactose-non-fermenting Gram-negative pathogens *Acinetobacter baumannii* and *Pseudomonas aeruginosa* occupy an important place. These bacterial species are quick to become multidrug-resistant owing to their additional intrinsic resistance mechanisms. They are responsible for hospital-acquired infections (bloodstream, urinary tract, pulmonary and device-related infections) and are frequently isolated from immunocompromised patients hospitalised in the intensive care unit. Resistance to multiple antibiotic classes, and notably to the β -lactam cephalosporins and carbapenems, is on the

rise worldwide. In this review, the emerging antibiotic resistance mechanisms in *A. baumannii* and *P. aeruginosa* are highlighted, with a special focus on the most prescribed antimicrobial agents, i.e. β -lactams, aminoglycosides and fluoroquinolones.

2. Resistance to β -lactams

2.1. Class A β -lactamases

2.1.1. Extended-spectrum β -lactamases (ESBLs)

The class A ESBLs confer resistance to expanded-spectrum cephalosporins and are inhibited in vitro by clavulanic acid and tazobactam [1]. They have been extensively identified in members of the Enterobacteriaceae family but are also reported from non-fermenters.

2.1.1.1. *Acinetobacter baumannii*. The most common ESBLs described in *A. baumannii* are the PER-, GES- and VEB-type β -lactamases. The first ESBL identified in *A. baumannii* was PER-1, being later widely detected in Turkey [2]. PER-1-producing *A. baumannii* are also considered to be widespread in South Korea [3], Hungary [4], Romania [5], Russia [6], Belgium [7] and the USA [8]. They have also been identified in Bulgaria, India, China, Iran and Kuwait [9–13] (Table 1). In *A. baumannii*, the *bla*_{PER-1} gene is part of a composite transposon named Tn1213, bracketed by two different insertion sequences (*ISPa12* and *ISPa13*) sharing similar

* Corresponding author. Present address: Medical and Molecular Microbiology Unit, Department of Medicine, Faculty of Science, University of Fribourg, rue Albert Gockel 3, CH-1700 Fribourg, Switzerland. Tel.: +41 26 300 9582; fax: +41 33670023872.

E-mail address: laurent.poirel@unifr.ch (L. Poirel).

Table 1Ambler class A extended-spectrum β -lactamases (ESBLs) known in *Acinetobacter baumannii* and *Pseudomonas aeruginosa*.

β -Lactamase	Host	Genetic support ^a	Country of isolation	Reference(s)
PER-1	<i>Acinetobacter baumannii</i>	C, P	Turkey	[2]
		C, P	South Korea	[3]
		?	Hungary	[4]
		C	Romania	[5]
		C	Russia	[6]
		C	Belgium	[7]
		?	USA	[8]
		?	Bulgaria	[9]
		C	India	[10]
		?	China	[11]
		?	Iran	[12]
		P	Kuwait	[13]
		C	France	[42]
		C, P	Turkey	[2]
	?	Belgium	[43]	
	C	Italy	[44]	
	C	Spain	[45]	
	C	Poland	[46]	
	P	Hungary	[47]	
	C	Serbia	[47]	
	?	Tunisia	[48]	
?	Japan	[49]		
?	China	[50]		
C	Greece	[51]		
?	Iran	[52]		
PER-2	<i>Acinetobacter baumannii</i>	?	Argentina	[15]
PER-3	<i>Pseudomonas aeruginosa</i>	?	Bolivia	[38]
PER-7	<i>Acinetobacter baumannii</i>	C	Egypt	[16]
PER-8	<i>Acinetobacter baumannii</i>	C	France	[17]
VEB-1	<i>Acinetobacter baumannii</i>	P	United Arab Emirates	[18]
VEB-1	<i>Acinetobacter baumannii</i>	?	Nepal	Accession no. AB985401
VEB-1	<i>Acinetobacter baumannii</i>	C	France	[19–21]
		C	Belgium	[7]
		?	Argentina	[15]
		?	Iran	[12]
		C	France	[53]
		C, P	Thailand	[54]
	<i>Pseudomonas aeruginosa</i>	C, P	Kuwait	[55]
		C	India	[56]
		?	Bulgaria	[57]
		?	UK	[58]
		?	Denmark	[59]
		?	Iran	[12]
		C	Thailand	[54]
VEB-2	<i>Pseudomonas aeruginosa</i>	C	Thailand	[54]
VEB-3	<i>Acinetobacter baumannii</i>	?	Taiwan	[23]
VEB-7	<i>Pseudomonas aeruginosa</i>	?	China	[60]
VEB-7	<i>Acinetobacter baumannii</i>	?	USA	Accession no. FJ825622
GES-1	<i>Pseudomonas aeruginosa</i>	C	France	[61]
		C	Brazil	[62]
		?	Argentina	[63]
GES-8 (IBC-2)	<i>Pseudomonas aeruginosa</i>	C	Greece	[64]
GES-9	<i>Pseudomonas aeruginosa</i>	C	France	[66]
GES-11	<i>Acinetobacter baumannii</i>	P	France	[25]
		P	Belgium	[26]
		?	Sweden	[27]
		P	Kuwait	[28]
		?	Turkey	[29]
		P	Tunisia	[30]
GES-12	<i>Acinetobacter baumannii</i>	P	Belgium	[26]
GES-13	<i>Pseudomonas aeruginosa</i>	C	Greece	[65]
GES-22	<i>Acinetobacter baumannii</i>	P	Turkey	[32]
SHV-2a	<i>Pseudomonas aeruginosa</i>	C	France	[67]
		C	Tunisia	[68]
SHV-5	<i>Acinetobacter baumannii</i>	C	France (from USA)	[33]
		C	Greece	[71]
SHV-12	<i>Acinetobacter baumannii</i>	P	The Netherlands	[35]
		C	Thailand	[69]
		C	Japan	[70]
TEM-4	<i>Pseudomonas aeruginosa</i>	C	France	[72]
TEM-21	<i>Pseudomonas aeruginosa</i>	C	France	[73]

Table 1 (Continued)

β-Lactamase	Host	Genetic support ^a	Country of isolation	Reference(s)
TEM-24	<i>Pseudomonas aeruginosa</i>	P	France	[74]
TEM-42	<i>Pseudomonas aeruginosa</i>	P	France	[75]
TEM-92	<i>Acinetobacter baumannii</i>	?	Italy	[34]
TEM-116	<i>Acinetobacter baumannii</i>	P	The Netherlands	[35]
CTX-M-1	<i>Pseudomonas aeruginosa</i>	?	The Netherlands	[76]
CTX-M-2	<i>Acinetobacter</i>	P	Japan	[36]
	<i>baumannii</i>	?	USA	[37]
	<i>Pseudomonas</i>	C	Brazil	[77,78]
	<i>aeruginosa</i>	?	Bolivia	[38]
CTX-M-3	<i>Pseudomonas aeruginosa</i>	?	China	[50]
CTX-M-14	<i>Pseudomonas aeruginosa</i>	?	China	[50]
CTX-M-15	<i>Acinetobacter baumannii</i>	P	India	[39]
		C	Haiti	[40]
	<i>Pseudomonas aeruginosa</i>	?	China	[50]
CTX-M-43	<i>Acinetobacter baumannii</i>	?	Bolivia	[38]
	<i>Pseudomonas aeruginosa</i>	?	Bolivia	[38]
RTG-4	<i>Acinetobacter baumannii</i>	P	France	[41]
BEL-1	<i>Pseudomonas aeruginosa</i>	C	Belgium	[79,80]
BEL-2	<i>Pseudomonas aeruginosa</i>	C	Belgium	[81]
BEL-3	<i>Pseudomonas aeruginosa</i>	?	Spain	[82]
PME-1	<i>Pseudomonas aeruginosa</i>	P	USA	[83]

^a C, chromosome; P, plasmid;?, unknown.

inverted repeat sequences [14]. Recently, the *bla*_{PER-1} gene was identified in a composite transposon made of two copies of *ISPa12* in an *A. baumannii* isolate from Kuwait [13]. PER-2, which is quite distantly related to PER-1 (86% amino acid identity), has so far been found exclusively in South America [15]. Recently, the *bla*_{PER-3} gene, initially identified from *Aeromonas punctata* and *Aeromonas caviae*, has been identified in a single *A. baumannii* isolate in Egypt [16]. In addition, PER-7 (four amino acid substitutions compared with PER-1) has been identified in a single *A. baumannii* clinical isolate in France [17] and in the United Arab Emirates (UAE) [18]. The *bla*_{PER-7} gene was associated with the insertion sequence (IS) element *ISCR1* that was also involved in its expression [17]. PER variants identified in *A. baumannii* are summarised Table 1.

Another important ESBL in *A. baumannii* is the Vietnamese extended-spectrum β-lactamase (VEB). VEB-1 is distantly related to other ESBLs, sharing only 38% amino acid identity with the closest ESBL, namely PER-1 [19]. VEB-1-producing *A. baumannii* were first identified in France, where a single clone was originally identified as the source of a hospital outbreak [20]. Genotyping analysis showed that this VEB-1-producing *A. baumannii* belonged to one of the two major clonal complexes of *A. baumannii*, termed worldwide clone 1 [20]. Further studies showed a nationwide dissemination of VEB-1 in France [21] and its neighbouring country Belgium [7]. In most cases, the *bla*_{VEB-1} gene is identified as a gene cassette in class 1 integrons varying in size and structure [19]. However, in several *A. baumannii* isolates from Argentina, the *bla*_{VEB-1} gene was associated with an *ISCR2* element, which was likely at the origin of the mobilisation of this ESBL gene [22]. VEB-1-producing *A. baumannii* have also been identified in Iran [12]. The *bla*_{VEB-3} variant was reported in a single *A. baumannii* isolate from Taiwan [23] (Table 1).

Since 2010, GES-type ESBLs are increasingly reported from *A. baumannii*. Actually, GES-1 was firstly reported in 2000, being identified from a single *K. pneumoniae* isolate [24]. GES-11, differing from GES-1 by two amino acid substitutions and consequently possessing increased activity towards aztreonam, was first identified in 2009 from an *A. baumannii* isolate from France [25]. GES-11 was then detected in the same species in Belgium, Sweden, Kuwait, Turkey and Tunisia [26–30] and in the Middle East, which might act as a reservoir for multidrug-resistant bacteria [28]. Another

GES variant, namely GES-12, differs from GES-11 by a single Thr237Ala substitution. It has been identified from several isolates in Belgium [26] and possesses increased hydrolytic activity towards ceftazidime [31]. More recently, GES-22, differing from GES-11 by one amino acid substitution, was reported from two *A. baumannii* isolates from Turkey [32]. It was reported that GES-22 possessed a hydrolytic profile similar to that of GES-11 and that the *bla*_{GES-22} gene was located on a class 1 integron inserted into a 75-kb plasmid [32].

On the other hand, the TEM- and SHV-type ESBLs, being widespread among Enterobacteriaceae, have been scarcely identified in *A. baumannii*. The corresponding *bla*_{SHV} and *bla*_{TEM} genes have been identified either on the chromosome (*bla*_{SHV-5}) or on plasmids (*bla*_{SHV-12}, *bla*_{TEM-92}, *bla*_{TEM-116}) [33–35]. Likewise, the genes encoding the CTX-M-type ESBLs, known to be extremely widespread among Enterobacteriaceae, have been rarely identified in *A. baumannii*. CTX-M-2-producing isolates have been identified in Japan and the USA [36,37], and a CTX-M-43-producing isolate has been found in Bolivia [38]. More recently, CTX-M-15-producing *A. baumannii* have been identified in India and Haiti [39,40]. The *bla*_{CTX-M-15} gene was found to be associated with *ISEcp1* in a transposon that integrated into the chromosome of *A. baumannii* [40]. A novel ESBL, named RTG-4, which is the first carbenicillinase to possess ESBL properties, was identified from an *A. baumannii* isolate from France in 2009 [41]. This is an atypical ESBL since it significantly hydrolyses cefepime and cefpirome, but hydrolyses ceftazidime only weakly [41].

Although widespread among Enterobacteriaceae, the rare identification of these ESBLs in *A. baumannii* may be due to limited horizontal gene transfer occurring between these different bacterial genders as a consequence of narrow-spectrum plasmid replication properties.

2.1.1.2. *Pseudomonas aeruginosa*. The PER-1 β-lactamase was the first ESBL identified in *P. aeruginosa* [42]. It was identified in a *P. aeruginosa* isolate from a Turkish patient hospitalised in the Paris area of France in 1991 [42]. National surveys from Turkey then showed that PER-1-producing *P. aeruginosa* isolates are widespread in Turkey [2]. PER-1 has been reported in European countries with no geographical border with Turkey, such as Belgium [43], Italy

[44], Spain [45], Poland [46], Hungary and Serbia [47] and Tunisia [48] as well as in Asian countries [49,50]. In addition, it was identified in Greece and Iran [51,52] (Table 1). Epidemiological surveys have shown that a predominant *P. aeruginosa* sequence type (ST) and single-locus variants, corresponding to international clonal complex CC11, is associated with wide dissemination of PER-1-producing *P. aeruginosa* isolates in Turkey, Belgium and Italy as well as in several Eastern European countries [46,47,51]. The PER-2 β -lactamase, which shares 86% amino acid identity with PER-1 and therefore represents another lineage of the PER-type enzymes, was identified only from a *P. aeruginosa* strain isolated in Bolivia [38] (Table 1).

Another ESBL from *P. aeruginosa* is VEB-1, which was identified from *P. aeruginosa* isolates recovered from patients hospitalised in France but transferred from Thailand [53]. Nosocomial spread of VEB-1-producing *P. aeruginosa* isolates was identified in Thailand [54]. Later, other VEB-like producing isolates were reported from Kuwait and India [55,56], but also in Iran, Bulgaria, the UK and Denmark, highlighting the worldwide dissemination of these VEB-producing strains [12,57–59] (Table 1). Isolates producing VEB-2 or VEB-3 were identified in Thailand and China, with these ESBLs differing from VEB-1 by only a single or two amino acid substitutions, respectively [54,60].

Other ESBLs are the GES enzymes, which have been detected in *P. aeruginosa* (Table 2). The bla_{GES-1} gene was identified from *P. aeruginosa* isolates from France and South America [61–63]. The structurally related ESBLs IBC-2 (differing from GES-1 by a single amino acid residue and then renamed GES-8) and GES-13 were isolated from two *P. aeruginosa* isolates in Greece [64,65]. Another variant, named GES-9, possessing a broad-spectrum hydrolysis profile extended to aztreonam, was identified in a single *P. aeruginosa* isolate from France [66].

The SHV-type ESBLs have been identified in very rare isolates of *P. aeruginosa*, being SHV-2a in France and Tunisia [67,68] and SHV-12 in Thailand [69] and Japan [70] (Table 1). A nosocomial outbreak of SHV-5-producing *P. aeruginosa* was also described in Greece [71]. TEM-type ESBLs have also been rarely reported from *P. aeruginosa*, being TEM-4 [72], TEM-21 [73], TEM-24 [74] and TEM-42 [75]. CTX-M-type ESBLs, with in some cases evidence of their horizontal transfer from Enterobacteriaceae to *P. aeruginosa*, are very rarely identified in *P. aeruginosa*. A single CTX-M-1-producing *P. aeruginosa* isolate has been reported from The Netherlands in 2006 [76], and CTX-M-2- or CTX-M-43-positive *P. aeruginosa* have been identified in South America [38,77,78]. Recently, CTX-M-3, CTX-M-14 and CTX-M-15 were identified from several *P. aeruginosa* isolates from China [50] (Table 1).

In 2005, another ESBL that had weak amino acid identity with other class A ESBLs but had similar biochemical properties was reported; the gene encoding BEL-1 was located in a class 1 integron inserted into the chromosome of a *P. aeruginosa* isolate recovered in a single hospital in Belgium [79]. Later, another study reported the dissemination of BEL-1-producing *P. aeruginosa* isolates in several hospitals located in different geographical areas in Belgium [80]. BEL-2 and BEL-3, each differing from BEL-1 by a single amino acid substitution, were identified in 2010 in a strain recovered in Belgium and Spain, respectively [81,82] (Table 1). Compared with BEL-1, BEL-2 possesses enhanced hydrolytic properties against expanded-spectrum cephalosporins [81].

The ESBL PME-1 is the latest identified ESBL from a clinical *P. aeruginosa* isolate and was recovered in Pennsylvania, USA, in 2008 [83]. This enzyme shares 43% amino acid identity with the closest ESBL CTX-M-9. PME-1 confers a high level of resistance to penicillins, ceftazidime and aztreonam and to a lesser extent cefotaxime, but spares cefepime and the carbapenems. The bla_{PME-1} gene was found to be located on an ca. 9-kb plasmid, flanked on both extremities by two copies of ISCR24 [83].

2.1.2. Carbapenemases

2.1.2.1. *Acinetobacter baumannii*.

Although almost all class A ESBLs do not possess any significant carbapenemase activity, specific GES variants have been shown to possess the ability to compromise the efficacy of carbapenems (Table 2). These are GES enzymes possessing specific residues enlarging their hydrolysis spectrum, and some of them such as GES-5 have been identified in Enterobacteriaceae [84]. The GES-14 variant is one of these GES-type carbapenemases and has been identified in *A. baumannii* in France in 2011 [85], the bla_{GES-14} gene being part of a class 1 integron located on a self-transferable plasmid [85].

Another class A carbapenemase that is commonly identified among Enterobacteriaceae is KPC, with KPC-type enzymes possessing intrinsic high carbapenemase and ESBL activity [86]. These enzymes all confer resistance to all β -lactams, and the corresponding genes are located on mobile genetic elements, enhancing their spread [87]. Despite wide dissemination among enterobacterial species, only a few KPC-type β -lactamases have been identified in *A. baumannii*, being from a series of isolates recovered in Puerto Rico [88]. In that study, ten *A. baumannii* isolates producing KPC-type enzymes were detected, corresponding to KPC-3 (7 isolates) and KPC-2, KPC-4 and KPC-10 in single isolates, respectively [88].

2.1.2.2. *Pseudomonas aeruginosa*.

As highlighted earlier, several GES-type ESBLs exhibit some carbapenemase properties. Actually, the first GES-type carbapenemase was identified from a *P. aeruginosa* isolate, being GES-2, differing from GES-1 by a single amino acid substitution [89]. This isolate was recovered from a patient hospitalised in South Africa and was actually part of an outbreak that occurred in the same hospital [90]. The GES-5 variant possessing significant carbapenemase activity has also been reported from *P. aeruginosa* isolates in China [91], South Africa [92], Brazil [93] and Turkey [94]. These bla_{GES} -type genes are part of class 1 integron structures [92]. Recently, a novel GES variant, GES-18, was identified from a *P. aeruginosa* isolate from Belgium. GES-18 differed from GES-5 by one amino acid substitution and also hydrolysed carbapenems [95].

Although rarely identified, KPC-producing *P. aeruginosa* isolates have been reported, first in Colombia in 2006 [96] and then in Puerto Rico [97,98], Trinidad and Tobago [99], the USA [100] and China [101]. They are increasingly identified in the Americas and the Caribbean region [102–104]. No clear evidence of horizontal transfer of the bla_{KPC} gene from Enterobacteriaceae to non-fermenters has been observed.

2.2. Class B β -lactamases

These β -lactamases, also named metallo- β -lactamases (MBLs), hydrolyse carbapenems and other β -lactams (except monobactams) very efficiently and they are not inhibited by the clinically available β -lactamase inhibitors such as clavulanic acid or tazobactam. However, their activity is inhibited by metal ion chelators [105,106].

2.2.1. *Acinetobacter baumannii*

Carbapenem resistance in this species is most often (if not always) linked to the production of carbapenemases. MBL enzymes are not the most commonly identified carbapenemases in *A. baumannii*; when identified, they are either IMP-like, VIM-like, SIM-1 or NDM-like enzymes [107]. Nine IMP variants have been identified in *A. baumannii*, namely IMP-1 in Italy [108], Japan [109], South Korea [110], India [111], Taiwan [112] and Kuwait [113], IMP-2 in Japan and Italy [109,114], IMP-4 in Hong-Kong [115], Australia and Singapore [116,117], IMP-5 in Portugal [118], IMP-6 in Brazil [119], IMP-8 in China [120], IMP-11 in Japan (accession no. **AB074436**),

Table 2Ambler class A carbapenemases known in *Acinetobacter baumannii* and *Pseudomonas aeruginosa*.

β -Lactamase	Host	Genetic support ^a	Country of isolation	Reference(s)
GES-2	<i>Pseudomonas aeruginosa</i>	P	South Africa	[89,90]
GES-5	<i>Pseudomonas aeruginosa</i>	?	China	[91]
		?	South Africa	[92]
		C	Brazil	[93]
		?	Turkey	[94]
GES-14	<i>Acinetobacter baumannii</i>	P	France	[85]
GES-18	<i>Pseudomonas aeruginosa</i>	C	Belgium	[95]
KPC-2	<i>Acinetobacter baumannii</i> <i>Pseudomonas aeruginosa</i>	?	Puerto Rico	[88]
		C, P	Columbia	[96,102]
		?	Puerto Rico	[97]
		?	Trinidad and Tobago	[99]
		P	USA	[100]
		C	China	[101]
		?	Argentina	[103]
		?	Brazil	[104]
KPC-3	<i>Acinetobacter baumannii</i>	?	Puerto Rico	[88]
KPC-4	<i>Acinetobacter baumannii</i>	?	Puerto Rico	[88]
KPC-5	<i>Pseudomonas aeruginosa</i>	?	Puerto Rico	[98]
KPC-10	<i>Acinetobacter baumannii</i>	?	Puerto Rico	[88]

^a C, chromosome; P, plasmid;?, unknown.

IMP-14 in Thailand [121] and IMP-19 in Japan [122] (Table 3). Noteworthy, VIM-type enzymes that have been widely identified in Enterobacteriaceae have rarely been identified in *A. baumannii*. There are few reports of VIM-1-producers in Greece [123], VIM-2 in South Korea [110] and Kuwait [113], VIM-4 in Italy [124], VIM-6 in India (accession no. **EF645347**) and VIM-11 in Taiwan [23] (Table 3).

The SIM-1 carbapenemase has been reported only in the *A. baumannii* species so far, and only in South Korea, where this resistance trait appears to be widespread [125]. Analysis of the genetic support of the MBL-encoding genes identified in *A. baumannii* shows similar structures, with the *bla*_{IMP}, *bla*_{VIM} and *bla*_{SIM} genes being all embedded in class 1 integron structures [107].

NDM-1 is one of the most recently identified MBLs [126]. Whilst most studies indicate wide dissemination of the *bla*_{NDM-1}-like genes in Enterobacteriaceae, many studies reported on the acquisition of *bla*_{NDM-1}-like genes in *A. baumannii*. Indeed, NDM-1 was first reported in India from Enterobacteriaceae and then in *A. baumannii* [126,127]. Other reports are from different European countries and from China, Japan, Kenya, Brazil, Algeria and Syria [128–137]. An outbreak of NDM-1-producing *A. baumannii*, belonging to ST85, was recently reported in France [138], underscoring the growing concern related to the spread of these isolates in Europe. Identification of several ST85 isolates possessing the *bla*_{NDM-1} gene and originating from North Africa, with no obvious link to the Indian subcontinent, strongly suggests that the source of NDM-producing *A. baumannii* strains could be North Africa [139]. Another variant, NDM-2, was identified in *A. baumannii* strains recovered in Egypt [140], Israel [141] and the UAE [142]. Interestingly, it was evidenced that these NDM-2-producing isolates were clonally related, suggesting that the Middle East as well as the Balkan region and the Indian and China regions might act as reservoirs of NDM-2-producing *Acinetobacter* [143]. In these isolates, the *bla*_{NDM} gene was surrounded by two copies of IS_{Aba125}, thus forming a 10 099-bp composite transposon named Tn₁₂₅ [144]. As opposed to what is observed in Enterobacteriaceae, the IS_{Aba125} element located upstream of *bla*_{NDM} and that plays a role in its expression, is not truncated [144]. Our extensive studies showed that *A. baumannii* was likely the first target of *bla*_{NDM-1} gene acquisition before its transfer to Enterobacteriaceae and *P. aeruginosa* [144]. This represents a new paradigm in antibiotic resistance since it highlights that *Acinetobacter* spp. may be a source of an important resistance trait for Enterobacteriaceae.

2.2.2. *Pseudomonas aeruginosa*

Carbapenem resistance in *P. aeruginosa* is mostly related to porin (OprD) deficiency and more rarely to carbapenemases. Carbapenemases in *P. aeruginosa* are mainly MBLs of the IMP, VIM, SPM and GIM types. IMP-1 was first reported in Enterobacteriaceae and *P. aeruginosa* in Japan and is now globally distributed, suggesting horizontal transfer of *bla*_{IMP-1} between unrelated Gram-negative species [145]. IMP-like enzymes may be divided into several subgroups and the percentage amino acid identity within these groups actually ranges from 90% to 99% [106]. These variants possess very similar hydrolytic activities. Among the 51 known IMP variants, 32 have been reported from *P. aeruginosa* and have been identified throughout the world (Table 3).

Although VIM enzymes share <40% amino acid identity with the IMP-type enzymes, they share the same hydrolytic spectrum [186]. VIM-1 was the first MBL identified in *P. aeruginosa* [187] and has been reported in several European countries (Table 3). However, VIM-2 is now the most widespread MBL in *P. aeruginosa* as a source of multiple outbreaks [106]. Twenty-three of the forty-six VIM variants have been identified in *P. aeruginosa* (Table 3).

β -Lactamase SPM is quite different from VIM and IMP and, accordingly, represents a new subfamily of MBLs. SPM-1 was first isolated in Brazil in 1997 from a *P. aeruginosa* clinical isolate [207], which was highly resistant to all anti-Gram-negative antibiotics except colistin. Dissemination of multidrug-resistant *P. aeruginosa* producing SPM-1 was demonstrated in distinct regions of this country, however they have not disseminated in other countries [208], with the only exception of a single isolate identified in Switzerland from a patient who had previously been hospitalised in Brazil [209]. The *bla*_{SPM-1} gene is either chromosomal or plasmid-encoded. In addition, it is associated with the IS element ISCR4 at the origin of its acquisition and expression and is likely transposed through a rolling-circle replication mechanism [210].

In 2002, a new type of acquired MBL, named GIM-1, was identified in clonally related *P. aeruginosa* isolates from Germany [211,212]. This enzyme also produced by enterobacterial species has only been identified in Germany.

NDM-1-producing *P. aeruginosa* isolates were first reported in 2011, with two strains recovered from Serbia [213]. In 2012, a single NDM-1-producing *P. aeruginosa* belonging to ST235 was isolated in France from a patient previously hospitalised in Serbia [214,215]. More recently, NDM-1-positive *P. aeruginosa* isolates were recovered in India (four isolates), Italy (a single isolate belonging to

Table 3
Ambler class B metallo- β -lactamases known in *Acinetobacter baumannii* and *Pseudomonas aeruginosa*.

β -Lactamase	Host	Genetic environment or support ^a	Country of isolation	Reference(s)
IMP-1	<i>Acinetobacter baumannii</i>	?	Italy	[108]
		1	Japan	[109]
		?	South Korea	[110]
		?	India	[111]
		?	Taiwan	[112]
	<i>Pseudomonas aeruginosa</i>	?	Kuwait	[113]
		1	Japan	[146]
		1	South Korea	[110]
		?	Brazil	[147]
		?	China	[148]
		1	Turkey	[149]
		?	Singapore	[150]
		1	Thailand	[151]
IMP-2	<i>Acinetobacter baumannii</i>	1	Japan	[109]
	<i>Acinetobacter baumannii</i>	1	Italy	[114]
	<i>Pseudomonas aeruginosa</i>	1	Japan	[109]
IMP-4	<i>Acinetobacter baumannii</i>	1	Hong Kong	[115]
		?	Australia	[116]
		1	Singapore	[117]
	<i>Pseudomonas aeruginosa</i>	?	Malaysia	[152]
		1	Australia	[153]
IMP-5	<i>Acinetobacter baumannii</i>	1	Portugal	[118]
	<i>Pseudomonas aeruginosa</i>	1	Portugal	[154]
IMP-6	<i>Acinetobacter baumannii</i>	?	Brazil	[119]
	<i>Pseudomonas aeruginosa</i>	1	South Korea	[155]
	<i>Pseudomonas aeruginosa</i>	?	China	[156]
IMP-7	<i>Pseudomonas aeruginosa</i>	1	Canada	[157]
		?	Malaysia	[158]
		?	Slovakia	[159]
		1	Japan	[160]
		?	Singapore	[150]
		1	Czech Republic	[161]
?	Denmark	[162]		
IMP-8	<i>Acinetobacter baumannii</i>	1	China	[120]
IMP-9	<i>Pseudomonas aeruginosa</i>	1	China	[163]
IMP-10	<i>Pseudomonas aeruginosa</i>	1	Japan	[164]
IMP-11	<i>Acinetobacter baumannii</i>	?	Japan	Accession no. AB074436
	<i>Pseudomonas aeruginosa</i>	?	Japan	Accession no. AB074437
IMP-13	<i>Pseudomonas aeruginosa</i>	1	Austria	[165]
		1	Italy	[166]
		1	France	[167]
		1	Belgium	[168]
IMP-14	<i>Acinetobacter baumannii</i>	1	Thailand	[121]
	<i>Pseudomonas aeruginosa</i>	1	Thailand	[169]
IMP-15	<i>Pseudomonas aeruginosa</i>	1	Mexico	[170]
		1	Spain	[171]
		?	Germany	[172]
IMP-16	<i>Pseudomonas aeruginosa</i>	1	Brazil	[173]
IMP-18	<i>Pseudomonas aeruginosa</i>	?	USA	[174]
		1	Mexico	[175]
		1	Puerto Rico	[97]
IMP-19	<i>Acinetobacter baumannii</i>	?	Japan	[122]
	<i>Pseudomonas aeruginosa</i>	?	Japan	Accession no. AB184876
	<i>Pseudomonas aeruginosa</i>	1	Italy	[176]
IMP-20	<i>Pseudomonas aeruginosa</i>	1	Japan	Accession no. AB196988
IMP-21	<i>Pseudomonas aeruginosa</i>	1	Japan	Accession no. AB204557
IMP-22	<i>Pseudomonas aeruginosa</i>	1	Austria	[165]
		1	Italy	[177]
IMP-25	<i>Pseudomonas aeruginosa</i>	1	China	Accession no. EU352796
IMP-26	<i>Pseudomonas aeruginosa</i>	1	Malaysia	[178]
		?	Singapore	[179]
IMP-29	<i>Pseudomonas aeruginosa</i>	1	France	[180]
IMP-30	<i>Pseudomonas aeruginosa</i>	?	Russia	[181]
IMP-31	<i>Pseudomonas aeruginosa</i>	?	Germany	Accession no. KF148593
IMP-33	<i>Pseudomonas aeruginosa</i>	1	Italy	[182]
IMP-35	<i>Pseudomonas aeruginosa</i>	1	Germany	[183]
IMP-37	<i>Pseudomonas aeruginosa</i>	?	France	Accession no. JX131372
IMP-40	<i>Pseudomonas aeruginosa</i>	?	Japan	Accession no. AB753457

Table 3 (Continued)

β -Lactamase	Host	Genetic environment or support ^a	Country of isolation	Reference(s)
IMP-41	<i>Pseudomonas aeruginosa</i>	?	Japan	Accession no. AB753458
IMP-43	<i>Pseudomonas aeruginosa</i>	1	Japan	[184]
IMP-44	<i>Pseudomonas aeruginosa</i>	1	Japan	[184]
IMP-45	<i>Pseudomonas aeruginosa</i>	1	China	[185]
IMP-48	<i>Pseudomonas aeruginosa</i>	?	USA	Accession no. KM087857
VIM-1	<i>Acinetobacter baumannii</i>	1	Greece	[123]
	<i>Pseudomonas aeruginosa</i>	1	Italy	[187]
		1	France	[188]
		1	Greece	[189]
		?	Germany	[172]
		1	Italy	[176]
VIM-2	<i>Acinetobacter baumannii</i>	1	South Korea	[110]
		?	Kuwait	[113]
	<i>Pseudomonas aeruginosa</i>	1	Tunisia	[190]
		?	Thailand	[169]
		1	Austria	[165]
		1	Mexico	[170]
		?	India	[191]
		?	Kenya	[192]
		1	Hungary	[193]
		1	Malaysia	[194]
		1	South Korea	[106]
		1	Japan	[106]
		1	France	[106]
		1	Greece	[106]
		1	Italy	[106]
		1	Portugal	[106]
		?	Spain	[106]
		1	Croatia	[106]
		1	Poland	[106]
		1	Chile	[106]
		1	Venezuela	[106]
		?	Argentina	[106]
		1	USA	[106]
		?	Belgium	[172]
		?	Germany	[172]
		?	Turkey	[172]
VIM-3	<i>Pseudomonas aeruginosa</i>	?	Taiwan	[195]
VIM-4	<i>Acinetobacter baumannii</i>	?	Italy	[124]
	<i>Pseudomonas aeruginosa</i>	1	Greece	[106]
		?	Sweden	[106]
		1	Poland	[106]
		1	Hungary	[193,196]
		?	France	[172]
VIM-5	<i>Pseudomonas aeruginosa</i>	?	India	[191]
		1	Turkey	[106]
VIM-6	<i>Acinetobacter baumannii</i>	?	India	Accession no. EF645347
	<i>Pseudomonas aeruginosa</i>	1	India	[191]
		1	Indonesia	[197]
		1	South Korea	[197]
		1	Philippines	[197]
VIM-7	<i>Pseudomonas aeruginosa</i>	1	USA	[198]
VIM-8	<i>Pseudomonas aeruginosa</i>	?	Columbia	[199]
VIM-9	<i>Pseudomonas aeruginosa</i>	?	UK	Accession no. AY524988
VIM-10	<i>Pseudomonas aeruginosa</i>	?	UK	[58]
VIM-11	<i>Acinetobacter baumannii</i>	1	Taiwan	[23]
	<i>Pseudomonas aeruginosa</i>	1	India	[191]
		?	Argentina	[200]
		?	Italy	Accession no. AY635904
		1	Malaysia	[194]
VIM-13	<i>Pseudomonas aeruginosa</i>	1	Spain	[201]
VIM-14	<i>Pseudomonas aeruginosa</i>	?	Spain	Accession no. EF055455
		1	Italy	[202]
VIM-15	<i>Pseudomonas aeruginosa</i>	1	Bulgaria	[203]
VIM-16	<i>Pseudomonas aeruginosa</i>	1	Germany	[203]
VIM-17	<i>Pseudomonas aeruginosa</i>	1	Greece	[204]
VIM-18	<i>Pseudomonas aeruginosa</i>	1	India	[191]
VIM-20	<i>Pseudomonas aeruginosa</i>	?	Spain	[205]
VIM-28	<i>Pseudomonas aeruginosa</i>	1	Egypt	[206]
VIM-30	<i>Pseudomonas aeruginosa</i>	1	France	Accession no. JN129451
VIM-36	<i>Pseudomonas aeruginosa</i>	?	Belgium	[172]

Table 3 (Continued)

β -Lactamase	Host	Genetic environment or support ^a	Country of isolation	Reference(s)
VIM-37	<i>Pseudomonas aeruginosa</i>	?	Poland	[172]
VIM-38	<i>Pseudomonas aeruginosa</i>	I	Turkey	[94]
SIM-1	<i>Acinetobacter baumannii</i>	I	South Korea	[125]
NDM-1	<i>Acinetobacter baumannii</i>	C	Czech Republic	[128]
		C	Germany	[129,131]
		C	Belgium	[130]
		C	Slovenia	[131]
		C	France	[131]
		C	Switzerland	[131]
		?	India	[127]
		P	China	[132]
		?	Japan	[133]
		?	Kenya	[134]
		P	Brazil	[135]
		?	Algeria	[136]
		?	Syria	[137]
		C	Serbia	[213]
		C	France	[214,215]
		P	India	[216]
		C	Italy	[217]
?	Egypt	[218]		
?	Slovakia	[219]		
NDM-2	<i>Acinetobacter baumannii</i>	C	Egypt	[140]
		C	Israel	[141]
		?	United Arab Emirates	[142]
SPM-1	<i>Pseudomonas aeruginosa</i>	ISCR4	Brazil	[207]
		ISCR4	Switzerland	[209]
GIM-1	<i>Pseudomonas aeruginosa</i>	I	Germany	[211,212]
FIM-1	<i>Pseudomonas aeruginosa</i>	C	Italy	[221]

^a I, integron present; C, chromosome; P, plasmid; ?, unknown.

ST235), Egypt and Slovakia [216–219]. Interestingly, the emergence of multidrug-resistant VIM-2-producing *P. aeruginosa* in Russia is also linked to a ST235-like dominant clone [220]. Association of ST235-like strains with MBL genes has been reported in several European countries [220], such as in Italy with VIM-1-producers, in Greece, Sweden, Hungary and Belgium with VIM-4-producers, in Spain with VIM-13-producers, and in France with IMP-29-producers [220]. This clone might therefore possess some specific traits enhancing its clonal dissemination.

Recently, a novel MBL named FIM-1, exhibiting its highest similarity (40% amino acid identity) with NDM-type enzymes, was reported in a *P. aeruginosa* isolate from Italy [221]. The *bla*_{FIM-1} gene was chromosomally located and was associated with ISCR19-like elements that were likely involved in its capture and mobilisation [221]; its origin remains unknown.

2.3. Class C β -lactamases

2.3.1. *Acinetobacter baumannii*

Acinetobacter baumannii naturally produces a gene encoding an AmpC-type cephalosporinase. This gene is usually expressed at a basal and low level, therefore the amount of AmpC produced does not have a significant impact on the activity of expanded-spectrum cephalosporins [222]. The presence of a specific IS element IS_{Aba1} (belonging to the IS4 family) upstream of this naturally occurring *ampC* gene provides promoter sequences enhancing its expression, resulting in resistance to broad-spectrum cephalosporins (but sparing carbapenems) [223]. By studying a series of *A. baumannii* strains, a variety of AmpC variants may be identified and these variants have been named ADC-type (*Acinetobacter*-derived cephalosporinase) enzymes [224]. Some ADC variants, such as ADC-33 and ADC-56, possess a slight extended activity towards expanded-spectrum cephalosporins, which allows the classification of these enzymes as extended-spectrum AmpC (ESAC) [225,226]. Indeed, they do hydrolyse ceftazidime more efficiently,

and in addition they hydrolyse fourth-generation cephalosporins such as cefepime, whereas wild-type AmpC enzymes do not. The true clinical significance of these enzymes remains unknown. No acquired AmpC-type-encoding gene has been identified so far in *A. baumannii*.

2.3.2. *Pseudomonas aeruginosa*

A chromosomal gene encoding an AmpC-type cephalosporinase is also intrinsic to *P. aeruginosa*. This *ampC* gene is associated with a LysR-type regulatory gene with which some β -lactam molecules may interact, leading to overexpression of this *ampC* gene [227]. Some β -lactams such as carbapenems are inducers of *ampC* gene expression, although they are not substrates of these cephalosporinases. Selection of mutants overproducing the *ampC* gene is frequently observed in *P. aeruginosa*, leading to acquired resistance to ticarcillin, piperacillin and broad-spectrum cephalosporins (cef-tazidime) [227]. In addition, insertion of the IS1669 element into the LysR regulatory gene (also known as the *ampR* gene) of *ampC* may lead to the overexpression of this enzyme [228]. Apart from these mechanisms leading to increased resistance to expanded-spectrum cephalosporins, very peculiar AmpC-type enzymes of *P. aeruginosa* have been identified possessing a broadened hydrolytic activity towards imipenem [229]. They correspond to naturally occurring, chromosomally encoded AmpC-type β -lactamases possessing an alanine residue at position 105 conferring an additional weak carbapenemase activity [229,230]. The true clinical significance of these enzymes as a source of carbapenem resistance remains to be clarified.

2.4. Class D β -lactamases

Class D β -lactamases, also known as oxacillinases, are β -lactamases grouped in a heterogeneous class of enzymes either with respect to their structural or biochemical properties [231]. These enzymes all hydrolyse amoxicillin and cefalotin and their

Table 4Ambler class D β -lactamases known in *Acinetobacter baumannii* and *Pseudomonas aeruginosa*.

Host	Enzyme subfamily	Additional OXA members	Phenotype	Reference(s)
<i>Acinetobacter baumannii</i>	OXA-23 (ARI-1)	OXA-27, OXA-49	CHDL	[107,236,237]
	OXA-40	OXA-25, OXA-26, OXA-72	CHDL	[232,238–245]
	OXA-58	OXA-96, OXA-97	CHDL	[117,246–248]
	OXA-143		CHDL	[249]
	OXA-235	OXA-236, OXA-237	CHDL	[250]
<i>Pseudomonas aeruginosa</i>	OXA-2	OXA-15, OXA-32, OXA-34, OXA-36, OXA-141, OXA-161	ES-OXA	[231,252–254]
	OXA-10	OXA-11, OXA-13, OXA-14, OXA-16, OXA-17, OXA-19, OXA-28, OXA-129, OXA-142, OXA-145, OXA-147, OXA-183	ES-OXA	[231,255–262]
	OXA-1	OXA-31	ES-OXA	[263]
	OXA-56	OXA-128	ES-OXA	[261]
	OXA-18		ES-OXA	[264]
	OXA-45		ES-OXA	[265]
	OXA-40		CHDL	[267]
	OXA-198		CHDL	[268]

CHDL, carbapenem-hydrolysing class D β -lactamase; ES-OXA, extended-spectrum oxacillinase.

activities are usually not significantly inhibited by clavulanic acid [232]. Some class D β -lactamases hydrolyse expanded-spectrum cephalosporins and a few have been identified in *P. aeruginosa*, but none in *A. baumannii*. Most of these broad-spectrum class D β -lactamases, also called ES-OXA, are point-mutant derivatives of narrow-spectrum β -lactamases [231]. In contrast, carbapenemase activity is also an intrinsic property of many class D β -lactamases, therefore terming them carbapenem-hydrolysing class D β -lactamases (CHDLs) [231].

2.4.1. *Acinetobacter baumannii*

Acinetobacter baumannii possesses naturally occurring class D β -lactamases, known as OXA-51-like enzymes [233]. These enzymes exhibit weak carbapenemase activity and are classified as CHDLs. Noticeably, the corresponding genes are not (or only weakly) expressed in most isolates. However, once overexpressed they may subsequently be involved in reduced susceptibility to carbapenems [234]. Overexpression of these genes encoding OXA-51-like enzymes is often driven by the insertion of an *ISAbA1* element upstream of the *bla*_{OXA-51}-like gene, providing strong promoter sequences.

In addition to these naturally occurring class D β -lactamases, several acquired class D β -lactamases have been identified as a source of carbapenem resistance in *A. baumannii* [107]. These CHDLs confer only reduced susceptibility to carbapenems, but they spare broad-spectrum cephalosporins. Therefore, the high resistance to carbapenems often observed in many *A. baumannii* strains results from the association between a CHDL and other resistance mechanisms, including porin loss and overexpression of efflux systems [107]. Five main groups of acquired CHDLs have been described in *A. baumannii*, corresponding to OXA-23-, OXA-40-, OXA-58-, OXA-143 and OXA-235-like enzymes. The first and most common subgroup of CHDLs is made of OXA-23, OXA-27 and OXA-49 (Table 4) [107]. The *bla*_{OXA-23}-like genes are chromosome- or plasmid-encoded and they are part of transposons, namely Tn2006 and Tn2007 [235]. OXA-23-like enzymes are the most widespread CHDLs in *A. baumannii* worldwide and they have been identified on all continents [107,236]. OXA-23-producing *A. baumannii* are the most common sources of nosocomial outbreaks with carbapenem-resistant *A. baumannii* [231,237]. A second group of acquired CHDLs in *A. baumannii* comprises OXA-25, OXA-26, OXA-40 (formerly known as OXA-24) and OXA-72 (Table 4). OXA-25 has been identified in carbapenem-resistant *A. baumannii* isolates recovered from Spain, and OXA-26 in Belgium [238]. The OXA-40 CHDL was originally identified in a carbapenem-resistant *A. baumannii* isolate in France recovered from a Portuguese patient [239], and then extensively identified in Portugal and Spain, as in other parts of

the world [231,240,241]. The *bla*_{OXA-40}-like genes may be either chromosome- or plasmid-located. OXA-72 has also been identified in different parts of the world (Brazil, Lithuania, Croatia), but predominantly in Asia (China, South Korea, Taiwan, Japan) [231,242–245]. A third group of CHDLs corresponds to OXA-58 and its structurally related enzymes (OXA-96 and OXA-97), first identified from a carbapenem-resistant *A. baumannii* isolate recovered in France [246] in the context of a nosocomial outbreak in a burn unit [247]. This *bla*_{OXA-58} gene has now been reported worldwide [231], being always plasmid-borne and associated with IS elements at the origin of its expression [107]. OXA-96 and OXA-97 are point-mutant variants of OXA-58 sharing the same hydrolytic properties and identified in Singapore and Tunisia, respectively [117,248].

The OXA-143 CHDL was identified in 2009 from a clinical *A. baumannii* isolate that had been recovered in Brazil [249]. It shares 88% amino acid identity with OXA-40, 63% with OXA-23 and 52% with OXA-58. Its substrate profile was similar to those of other CHDLs and its corresponding gene was not integron- or transposon-encoded [249].

Ultimately, a novel subclass of CHDLs has recently been reported from isolates recovered in the USA and Mexico. This subgroup includes OXA-235, OXA-236 and OXA-237 [250], and the corresponding genes have been identified either on chromosomes or plasmids, and bracketed by two copies of *ISAbA1* [250].

2.4.2. *Pseudomonas aeruginosa*

Pseudomonas aeruginosa produces a naturally occurring class D β -lactamase, OXA-50, that does not contribute to the overall β -lactam resistance pattern of *P. aeruginosa*, except for latamoxef [251]. Most of the class D β -lactamases able to hydrolyse expanded-spectrum cephalosporins have been identified in *P. aeruginosa*. There are two main types of expanded-spectrum class D β -lactamases (ES-OXAs). Some are point-mutant derivatives of narrow-spectrum class D β -lactamases, with amino acid substitutions enlarging their spectrum of hydrolysis towards expanded-spectrum cephalosporins. These ES-OXAs mainly derive from the narrow-spectrum β -lactamases OXA-10 (OXA-11, -13, -14, -16, -17, -19, -28, -129, -142, -145, -147 and -183), OXA-2 (OXA-15, -32, -34, -36, -141 and -161) and OXA-1 (OXA-31) (Table 4) [231,252–263]. Other ES-OXAs share only weak amino acid identity with these latter enzymes. OXA-18, which is inhibited by clavulanic acid, was the first identified ES-OXA in a *P. aeruginosa* isolate in Paris from a patient previously hospitalised in Sicily [264]. This enzyme shares <50% amino acid identity with the other class D β -lactamases. OXA-45 is another ES-OXA, identified from a multidrug-resistant Texan *P. aeruginosa* isolate co-expressing

the class B β -lactamase VIM-7 [265]. OXA-45 shares the highest identity with OXA-18 (66%) and, as for OXA-18, its activity is well inhibited by clavulanic acid. Interestingly, the *bla*_{OXA-45} gene, located on a 24-kb plasmid, was not found as a form of a gene cassette but associated with two copies of an ISCR5-like element [266].

Only two acquired class D β -lactamases compromising the efficacy of carbapenems have been reported from *P. aeruginosa*. One is OXA-40, known to be widespread in *A. baumannii*, which was reported in a *P. aeruginosa* isolate in Spain in 2006 [267]. The other is OXA-198 characterised in a *P. aeruginosa* isolate from Belgium [268], sharing <30% amino acid identity with other CHDLs reported in Gram-negative organisms. The *bla*_{OXA-198} gene was harboured by a class 1 integron carried by a non-typeable ca. 46-kb plasmid [268].

3. Broad resistance to aminoglycosides

Aminoglycosides are used in the treatment of a broad range of life-threatening infections. The activity of aminoglycosides depends on binding to a highly conserved motif of 16S rRNA. Mechanisms of aminoglycosides resistance include decreased outer membrane permeability, active efflux and amino acid substitutions in ribosomal proteins, whereas the most common resistance mechanism is enzymatic leading to modification of the drug. Methylation of 16S ribosomal RNA has recently been demonstrated to be another mechanism of resistance encountered in Gram-negative organisms, corresponding to a modification of the antibiotic target [269]. Methylases actually interfere in the binding of these antibiotics to their site of action. These 16S rRNA methylases confer a high level of resistance to clinically useful aminoglycosides such as amikacin, gentamicin and tobramycin [269,270]. The corresponding genes are associated with transposon structures, which are themselves located on transferable plasmids, enhancing their horizontal spread. Isolates producing 16S rRNA methylases are multidrug-resistant, in particular to broad-spectrum β -lactams through the production of ESBLs or MBLs. This multidrug resistance pattern makes treatment of these infections particularly challenging. Ten 16S rRNA methylases have been identified among Gram-negative isolates, namely ArmA, RmtA, RmtB, RmtC, RmtD, RmtD2, RmtE, RmtF, RmtG and NpmA [270–273]. The origin of these genes is likely *Streptomyces* [271] and their prevalence remains unknown depending on the geographical location (possibly more frequent in Asia).

3.1. *Acinetobacter baumannii*

The ArmA enzyme has been reported in many *A. baumannii* worldwide, conferring high-level resistance to all aminoglycosides. Such isolates have been identified in China [274], South Korea [275,276], Vietnam [277], Japan [278], North America [279], Norway [280], Italy [281], Bulgaria [282], Iran [283] and Algeria [136]. The *armA* gene was always found to be located on a functional composite transposon Tn1548 [279]. Despite being quite widespread among *A. baumannii* strains, the *armA* gene possesses a GC content of 30%, which significantly differs from that of the *A. baumannii* core genome estimated at ca. 39%. This highlights the fact that this gene was acquired horizontally from a source that still remains unknown [279]. Noteworthy, the ArmA-encoding gene is often identified among OXA-23-producing *A. baumannii* strains, however both resistance genes are not physically linked on a single plasmid [280–282].

Apart from numerous reports of ArmA-producing *A. baumannii* isolates, the 16S rRNA methylase RmtB has recently been identified in nine *A. baumannii* isolates from Vietnam [277].

3.2. *Pseudomonas aeruginosa*

The first 16S rRNA methylase recovered in *P. aeruginosa* was identified in 2003. It was a clinical isolate from Japan producing RmtA [284,285]. Other RmtA-producing *P. aeruginosa* were then identified in South Korea in 2009 [286]. The *rmtA* gene is located on mobile genetic elements such as transposon Tn5041 [285].

In 2007, the RmtD methylase was firstly reported from a pandrug-resistant *P. aeruginosa* clinical isolate co-producing the MBL SPM-1 in Brazil [287]. RmtD shares 40% amino acid identity with RmtA, and the genetic structures surrounding both corresponding genes shared similar features [287]. Subsequently, another study underscored that co-production of the MBL SPM-1 and the 16S rRNA methylase RmtD was common among imipenem-resistant *P. aeruginosa* isolates recovered in hospitals in São Paulo, Brazil [288]. In these isolates, both the *bla*_{SPM-1} and *rmtD* genes were found to be chromosomally located. The ArmA enzyme was also identified in *P. aeruginosa* co-producing the MBL IMP-1 in South Korea [289].

4. Broad resistance to fluoroquinolones

In Gram-negative organisms, acquisition of resistance to quinolones may be related to chromosomal mutations in genes encoding the topoisomerases or to mutations in the efflux pump regulation systems. In addition, plasmid-mediated quinolone resistance genes (coding for the Qnr proteins) have been identified in Enterobacteriaceae. These acquired Qnr proteins have not been identified in non-fermenters. In *P. aeruginosa* and *A. baumannii*, a single mutation in the *gyrA* gene encoding DNA gyrase is sufficient to confer clinically high-level resistance levels to fluoroquinolones. This is due to the fact that these species intrinsically possess a decreased susceptibility to these antibiotics owing to low permeability or constitutive expression of efflux pumps.

4.1. *Acinetobacter baumannii*

Overexpression of efflux pumps is a source of acquired resistance to fluoroquinolones in this species. Involvement of the *adeABC* operon encoding the AdeA, AdeB and AdeC proteins forming a resistance–nodulation–cell division (RND) efflux system has been demonstrated [290]. In this case, not only fluoroquinolones but also aminoglycosides, tetracyclines, chloramphenicol and trimethoprim are substrates of this efflux system. Therefore, co-selection of this kind of mechanism with non-quinolone antibiotic molecules is possible. Similarly, overexpression of the *adeIJK* operon encoding another RND efflux system of *A. baumannii* has also been shown to interfere with different antibiotics, including the fluoroquinolones [291]. Finally, overproduction of the AbeM efflux system [belonging to the multi-antimicrobial extrusion protein (MATE) family] also contributes to acquired resistance to fluoroquinolones [292].

4.2. *Pseudomonas aeruginosa*

As shown in *A. baumannii*, acquired resistance to fluoroquinolones, apart from being related to mutations in topoisomerase-encoding genes, is mainly related to efflux systems. Their downregulation or upregulation (depending on whether the regulator is positive or negative) contribute significantly to reduced activity of fluoroquinolones. In *P. aeruginosa*, involvement of the MexAB–OprM RND-type system, which expression is constitutive, has been demonstrated, and its wide effect has been highlighted, conferring reduced susceptibility also to chloramphenicol, tetracyclines and β -lactams [293]. Likewise, the *mexCD–oprJ* operon confers extrusion ability with regard to quinolones, penicillins and tetracyclines. Nevertheless, this impact is seen only when the

negative regulatory protein NfxB is altered, indicating that constitutive expression of this efflux system does not contribute to this resistance trait [294]. Another efflux system is MexEF-OprN, which is also not involved in reduced susceptibility unless overexpression is observed [295].

5. Resistance to tigecycline

Tigecycline, a semisynthetic derivative of minocycline, has a peculiar mechanism of action and overcomes the widely distributed *tet* gene-encoded resistance mechanism known to confer resistance to tetracycline. Tigecycline shows good activity towards Gram-negative pathogens that may produce a large array of resistance mechanisms, including ESBLs and carbapenemases [296]. The activity of tigecycline against *A. baumannii* is overall good, and successful results have been reported clinically [296]. Resistance has been noted on several occasions and might be due to upregulation of the AdeABC multidrug efflux pump [297]. Another *A. baumannii*-specific efflux system, named AdelJK, has also been shown to interfere with the efficacy of tigecycline, acting synergistically with AdeABC [291]. Despite the fact that tigecycline has been shown to also be a substrate of the AdeFGH efflux pump [298], only AdeABC and AdelJK efflux pumps appear to be involved in its resistance in clinical isolates [298–300]. Noteworthy, tigecycline resistance levels in *A. baumannii* isolates may increase during therapy with tigecycline in the case of brief exposure to the drug, compromising its efficiency [301]. It was recently reported that a mutation in the *trm* gene, encoding a methyltransferase, was associated with decreased susceptibility to tigecycline in an *A. baumannii* strain [302]. This Trm enzyme could therefore play a role in resistance to tigecycline, however further studies are needed to clarify the possible role of this methyltransferase in the decreased susceptibility to tigecycline [302].

6. Resistance to colistin

During the past decade, we have witnessed a renewal of clinical interest in polymyxins (colistin) owing to two concomitant facts: (i) the emergence of carbapenem-, cephalosporin- and aminoglycoside-resistant Gram-negative isolates; and (ii) the paucity of novel marketed antibiotic molecules. Actually, polymyxins remain most often active against these multidrug-resistant isolates [303]. Emergence of colistin resistance in relation to increased usage is worrisome since polymyxins are the last remaining therapeutic option in many cases. Acquisition of resistance is mostly related to modifications of the lipopolysaccharide (LPS) biosynthesis pathway.

6.1. *Acinetobacter baumannii*

Two mechanisms of resistance to colistin have been described in *A. baumannii*: (i) alterations of the lipid A component of LPS resulting from mutations in the PmrAB two-component system [304,305]; and (ii) complete loss of LPS production resulting from mutations in the *lpxA*, *lpxC* and *lpxD* genes encoding the enzymes that catalyse the first steps in LPS biosynthesis [306]. Resistance to colistin in *A. baumannii* clinical isolates is rarely reported, however evaluation of colistin susceptibility is difficult and often inaccurate, considering that many laboratories do not use the gold standard technique for testing, which is microdilution [307]. The first report of colistin-resistant *A. baumannii* was from South Korea in 2005 [303]. High colistin resistance rates were then reported in other Korean hospitals in 2007 [308], and an outbreak of pandrug-resistant (including colistin) *A. baumannii* strains was reported in Spain in 2009 [309]. More recently, colistin-resistant *A. baumannii*

were recovered in Iran and the USA [310,311]. Noteworthy, development of heteroresistance to colistin in an in vitro model might compromise the use of colistin as a therapeutic option [312].

6.2. *Pseudomonas aeruginosa*

The emergence of colistin-resistant *P. aeruginosa* isolates has also been reported worldwide [313–315]. As observed in *A. baumannii*, resistance to polymyxins is associated with modifications of the lipid A component of LPS. Several two-component regulatory systems, such as PmrAB, PhoPQ, ParRS, CprRS and ColRS, are involved in resistance to polymyxins in *P. aeruginosa* [316–318]. A recent study identified nine genes with amino acid alterations and altered expression levels in colistin-resistant *P. aeruginosa* isolates compared with an isogenic colistin-susceptible isolate [319]. Thus, resistance to colistin is basically mediated by a complicated regulatory network involving a large array of chromosomal genes, which is currently under investigation worldwide, with some side experiments aiming to evaluate the fitness cost of such resistance.

7. Concluding remarks

Increasing rates of bacterial resistance among non-fermenters are threatening the effectiveness of antibiotics used as last-resort therapeutic options. In *A. baumannii* and *P. aeruginosa*, acquisition of resistance traits to these molecules is becoming more and more frequent, leading to multidrug and pandrug resistance. The last years have shown that: (i) most of the broad-spectrum resistance patterns identified in Enterobacteriaceae may be also identified in *P. aeruginosa* and *A. baumannii*; (ii) accumulation of unrelated resistance mechanisms (e.g. to β -lactams and aminoglycosides) is observed daily worldwide; and (iii) *A. baumannii*, although being a weak pathogen compared with *P. aeruginosa*, may play a significant role in spreading broad-spectrum resistance genes to other Gram-negative organisms. International travel and transfer of hospitalised patients further enhances this spread. In parallel, antibiotic selective pressure is also on the rise, in particular for these last-resort antibiotics that have been used only scarcely until recently. Since very few novel and effective antibiotics for the treatment of infections due to multidrug-resistant Gram-negatives isolates are going to be launched in a near future, there is an urgent need to implement strategies that may slow the development of acquired resistance. Use of rapid diagnostic techniques for detection of resistance traits, development of selective media for early recognition of colonised patients, and improvement of antibiotic stewardship may contribute to this containment strategy against antibiotic resistance.

Funding: This work was funded by the University of Fribourg (Fribourg, Switzerland).

Competing interests: None declared.

Ethical approval: Not required.

References

- [1] Weldhagen GF, Poirel L, Nordmann P. Ambler class A extended-spectrum β -lactamases in *Pseudomonas aeruginosa*: novel developments and clinical impact. *Antimicrob Agents Chemother* 2003;47:2385–92.
- [2] Kolayli F, Gacar G, Karadenizli A, Sanic A, Vahaboglu H. PER-1 is still widespread in Turkish hospitals among *Pseudomonas aeruginosa* and *Acinetobacter* spp. *FEMS Microbiol Lett* 2005;15:241–5.
- [3] Yong D, Shin H, Kim S, Lim Y, Yum JH, Lee K, et al. High prevalence of PER-1 extended-spectrum β -lactamase-producing *Acinetobacter* spp. in Korea. *Antimicrob Agents Chemother* 2003;47:1749–51.

- [4] Szabó D, Szentandrassy J, Juhász Z, Katona K, Nagy K, Rókusz L. Imported PER-1 producing *Pseudomonas aeruginosa*, PER-1 producing *Acinetobacter baumannii* and VIM-2-producing *Pseudomonas aeruginosa* strains in Hungary. *Ann Clin Microbiol Antimicrob* 2008;7:12.
- [5] Naas T, Nordmann P, Heidt A. Inter-country transfer or PER-1 extended-spectrum β -lactamase-producing *Acinetobacter baumannii* from Romania. *Int J Antimicrob Agents* 2007;29:226–8.
- [6] Naas T, Kernbaum S, Allali S, Nordmann P. Multidrug-resistant *Acinetobacter baumannii*, Russia. *Emerg Infect Dis* 2007;13:669–71.
- [7] Naas T, Bogaerts P, Bauraing C, Degheldre Y, Glupczynski Y, Nordmann P. Emergence of PER and VEB extended-spectrum β -lactamases in *Acinetobacter baumannii* in Belgium. *J Antimicrob Chemother* 2006;58:178–82.
- [8] Hujer KM, Hujer AM, Hulten EA, Bajaksouzian S, Adams JM, Donskey CJ, et al. Analysis of antibiotic resistance genes in multidrug-resistant *Acinetobacter* sp. isolates from military and civilian patients treated at the Walter Reed Army Medical Center. *Antimicrob Agents Chemother* 2006;50:4114–23.
- [9] Strateva T, Todorova A, Ouzounova-Raykova V, Mitov I. Emergence of a PER-1 extended-spectrum β -lactamase-producing *Acinetobacter baumannii* clinical isolate in Bulgaria. *J Chemother* 2008;20:391–2.
- [10] Litake GM, Ghole VS, Niphadkar KB, Joshi SG. PER-1-type extended-spectrum β -lactamase-producing *Acinetobacter baumannii* clinical isolates from India. *Int J Antimicrob Agents* 2009;34:388–9.
- [11] Zhang JP, Zhu W, Tian SF, Chu YZ, Chen BY. Molecular characteristics and resistant mechanisms of imipenem-resistant *Acinetobacter baumannii* isolates in Shenyang, China. *J Microbiol* 2010;48:689–94.
- [12] Farajnia S, Azhari F, Alikhani MY, Hosseini MK, Peymani A, Sohrabi N. Prevalence of PER and VEB type extended spectrum β -lactamases among multidrug resistant *Acinetobacter baumannii* isolates in North-West of Iran. *Iran J Basic Med Sci* 2013;16:751–5.
- [13] Opazo A, Vali L, Al Obaid K, Dashti AA, Amyes SG. Novel genetic structure harbouring *bla*_{PER-1} in ceftazidime-resistant *Acinetobacter baumannii* isolated from Kuwait. *Int J Antimicrob Agents* 2014;43:383–4.
- [14] Poirel L, Cabanne L, Vahaboglu H, Nordmann P. Genetic environment and expression of the extended-spectrum β -lactamase *bla*_{PER-1} gene in Gram-negative bacteria. *Antimicrob Agents Chemother* 2005;49:1708–13.
- [15] Pasterán F, Rapoport M, Petroni A, Faccione D, Corso A, Galas M, et al. Emergence of PER-2 and VEB-1a in *Acinetobacter baumannii* strains in the Americas. *Antimicrob Agents Chemother* 2006;50:3222–4.
- [16] Al-Hassan L, El Mahallawy H, Amyes SG. First report of *bla*_{PER-3} in *Acinetobacter baumannii*. *Int J Antimicrob Agents* 2013;41:93–4.
- [17] Bonnin RA, Potron A, Poirel L, Lécuyer H, Néri R, Nordmann P. PER-7, an extended-spectrum β -lactamase with increased activity toward broad-spectrum cephalosporins in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2011;55:2424–7.
- [18] Opazo A, Sonnevend A, Lopes B, Hamouda A, Ghazawi A, Pal T, et al. Plasmid-encoded PER-7 β -lactamase responsible for ceftazidime resistance in *Acinetobacter baumannii* isolated in the United Arab Emirates. *J Antimicrob Chemother* 2012;67:1619–22.
- [19] Poirel L, Naas T, Guibert M, Chaibi EB, Labia R, Nordmann P. Molecular and biochemical characterization of VEB-1, a novel class A extended-spectrum β -lactamase encoded by an *Escherichia coli* integron gene. *Antimicrob Agents Chemother* 1999;43:573–81.
- [20] Poirel L, Menuteau O, Agoli N, Cattoen C, Nordmann P. Outbreak of extended-spectrum β -lactamase VEB-1-producing isolates of *Acinetobacter baumannii* in a French hospital. *J Clin Microbiol* 2003;41:3542–7.
- [21] Naas T, Coignard B, Carbonne A, Blanckaert K, Bajolet O, Bernet C, et al. VEB-1 extended-spectrum β -lactamase-producing *Acinetobacter baumannii*, France. *Emerg Infect Dis* 2006;12:1214–22.
- [22] Poirel L, Mugnier PD, Toleman MA, Walsh TR, Rapoport MJ, Petroni A, et al. ISCR2, another vehicle for *bla*_{VEB} gene acquisition. *Antimicrob Agents Chemother* 2009;53:4940–3.
- [23] Huang LY, Chen TL, Lu PL, Tsai CA, Cho WL, Chang FY, et al. Dissemination of multidrug-resistant, class 1 integron-carrying *Acinetobacter baumannii* isolates in Taiwan. *Clin Microbiol Infect* 2008;14:1010–9.
- [24] Poirel L, Le Thomas I, Naas T, Karim A, Nordmann P. Biochemical sequence analyses of GES-1, a novel class A extended-spectrum β -lactamase, and the class 1 integron In52 from *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 2000;44:622–32.
- [25] Moubareck C, Brémont S, Conroy MC, Courvalin P, Lambert T. GES-11, a novel integron-associated GES variant in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2009;53:3579–81.
- [26] Bogaerts P, Naas T, El Garch F, Cuzon G, Deplano A, Delaire T, et al. GES extended-spectrum β -lactamases in *Acinetobacter baumannii* isolates in Belgium. *Antimicrob Agents Chemother* 2010;54:4872–8.
- [27] Karah N, Giske CG, Sundsfjord A, Samuelsen O. A diversity of OXA-carbapenemases and class 1 integrons among carbapenem-resistant *Acinetobacter baumannii* clinical isolates from Sweden belonging to different international clonal lineages. *Microb Drug Resist* 2011;17:545–9.
- [28] Bonnin RA, Rotimi VO, Al Hubail M, Gasiorowski E, Al Sweih N, Nordmann P, et al. Wide dissemination of GES-type carbapenemases in *Acinetobacter baumannii* isolates in Kuwait. *Antimicrob Agents Chemother* 2013;57:183–8.
- [29] Cicek AC, Saral A, Iraz M, Ceylan A, Duzgun AO, Peleg AY, et al. OXA- and GES-type β -lactamases predominate in extensively drug-resistant *Acinetobacter baumannii* isolates from a Turkish university hospital. *Clin Microbiol Infect* 2014;20:410–5.
- [30] Charfi-Kessiss K, Mansour W, Ben Haj Khalifa A, Mastouri M, Nordmann P, Aouni M, et al. Multidrug-resistant *Acinetobacter baumannii* strains carrying the *bla*_{OXA-23} and the *bla*_{GES-11} genes in a neonatology center in Tunisia. *Microb Pathog* 2014;74:20–4.
- [31] Delbrück H, Bogaerts P, Kupper MB, Rezende de Castro R, Bennink S, Glupczynski Y, et al. Kinetic and crystallographic studies of extended-spectrum GES-11, GES-12, and GES-14 β -lactamases. *Antimicrob Agents Chemother* 2012;56:5618–25.
- [32] Castanheira M, Costello SE, Woosley LN, Deshpande LM, Davies TA, Jones RN. Evaluation of clonality and carbapenem resistance mechanisms among *Acinetobacter baumannii*-*Acinetobacter calcoaceticus* complex and Enterobacteriaceae isolates collected in European and Mediterranean countries and detection of two novel β -lactamases, GES-22 and VIM-35. *Antimicrob Agents Chemother* 2014;58:7358–66.
- [33] Naas T, Namdari F, Réglier-Poupet H, Poyart C, Nordmann P. Panresistant extended-spectrum β -lactamase SHV-5-producing *Acinetobacter baumannii* from New York City. *J Antimicrob Chemother* 2007;60:1174–6.
- [34] Endimiani A, Luzzaro F, Migliavacca R, Mantengoli E, Hujer AM, Hujer KM, et al. Spread in an Italian hospital of a clonal *Acinetobacter baumannii* strain producing the TEM-92 extended-spectrum β -lactamase. *Antimicrob Agents Chemother* 2007;51:2211–4.
- [35] Naiemi NA, Duim B, Savelkoul PH, Spanjaard L, de Jonge E, Bart A, et al. Widespread transfer of resistance genes between bacterial species in an intensive care unit: implications for hospital epidemiology. *J Clin Microbiol* 2005;43:4862–4.
- [36] Nagano N, Nagano Y, Cordevant C, Shibata N, Arakawa Y. Nosocomial transmission of CTX-M-2 β -lactamase-producing *Acinetobacter baumannii* in a neurosurgery ward. *J Clin Microbiol* 2004;42:3978–84.
- [37] Adams-Haduch JM, Paterson DL, Sidjabat HE, Pasculle AW, Potoski BA, Muto CA, et al. Genetic basis of multidrug resistance in *Acinetobacter baumannii* clinical isolates at a tertiary medical center in Pennsylvania. *Antimicrob Agents Chemother* 2008;52:3837–43.
- [38] Celenza G, Pellegrini C, Caccamo M, Segatore B, Amicosante G, Perilli M. Spread of *bla*_{CTX-M-type} and *bla*_{PER-2} β -lactamase genes in clinical isolates from Bolivian hospitals. *J Antimicrob Chemother* 2006;57:975–8.
- [39] Shakil S, Khan AU. Detection of CTX-M-15-producing and carbapenem-resistant *Acinetobacter baumannii* strains from urine from an Indian hospital. *J Chemother* 2010;22:324–7.
- [40] Potron A, Munoz-Price LS, Nordmann P, Cleary T, Poirel L. Genetic features of CTX-M-15-producing *Acinetobacter baumannii* from Haiti. *Antimicrob Agents Chemother* 2011;55:5946–8.
- [41] Potron A, Poirel L, Croizé J, Chantepredrix V, Nordmann P. Genetic and biochemical characterization of the first extended-spectrum CARB-type β -lactamase, RTG-4, from *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2009;53:3010–6.
- [42] Nordmann P, Ronco E, Naas T, Dupont C, Michel-Briand Y, Labia R. Characterization of a novel extended-spectrum β -lactamase from *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 1993;37:962–9.
- [43] Claeys G, Verschraegen G, de Baere T, Vaneechoutte M. PER-1 β -lactamase-producing *Pseudomonas aeruginosa* in an intensive care unit. *J Antimicrob Chemother* 2000;45:924–5.
- [44] Luzzaro F, Mantengoli E, Perilli M, Lombardi G, Orlandi V, Orsatti A, et al. Dynamics of a nosocomial outbreak of multidrug-resistant *Pseudomonas aeruginosa* producing the PER-1 extended-spectrum β -lactamase. *J Clin Microbiol* 2001;39:1865–70.
- [45] Tato M, Valverde A, Coque TM, Cantón R. PER-1 multiresistant *Pseudomonas aeruginosa* strain in Spain [in Spanish]. *Enferm Infect Microbiol Clin* 2006;24:472–3.
- [46] Empel J, Filczak K, Mrówka A, Hryniewicz W, Livermore DM, Gniadkowski M. Outbreak of *Pseudomonas aeruginosa* infections with PER-1 extended-spectrum β -lactamase in Warsaw, Poland: further evidence for an international clonal complex. *J Clin Microbiol* 2007;45:2829–34.
- [47] Libisch B, Poirel L, Lepsanovic Z, Mirovic V, Balogh B, Paszti J, et al. Identification of PER-1 extended-spectrum β -lactamase producing *Pseudomonas aeruginosa* clinical isolates of the international clonal complex CC11 from Hungary and Serbia. *FEMS Immunol Med Microbiol* 2008;54:330–8.
- [48] Ktari S, Mnif B, Znazen A, Rekek M, Mezghani S, Mahjoubi-Rhimi F, et al. Diversity of β -lactamases in *Pseudomonas aeruginosa* isolates producing metallo- β -lactamase in two Tunisian hospitals. *Microb Drug Resist* 2011;17:25–30.
- [49] Yamano Y, Nishikawa T, Fujimura T, Yutsudou T, Tsuji M, Miwa H. Occurrence of PER-1 producing clinical isolates of *Pseudomonas aeruginosa* in Japan and their susceptibility to doripenem. *J Antibiot (Tokyo)* 2006;59:791–6.
- [50] Qing Y, Cao KY, Fang ZL, Huang YM, Zhang XF, Tian GB, et al. Outbreak of PER-1 and diversity of β -lactamases among ceftazidime-resistant *Pseudomonas aeruginosa* clinical isolates in China. *J Med Microbiol* 2014;63:386–92.
- [51] Ranellou K, Kadlec K, Poulou A, Voulgaru E, Vrioni G, Schwarz S, et al. Detection of *Pseudomonas aeruginosa* isolates of the international clonal complex 11 carrying the *bla*_{PER-1} extended-spectrum β -lactamase gene in Greece. *J Antimicrob Chemother* 2012;67:357–61.
- [52] Mirsalehian A, Feizabadi M, Nakhjavani FA, Jabalameli F, Goli H, Kalantari N. Detection of VEB-1, OXA-10 and PER-1 genotypes in extended-spectrum β -lactamase-producing *Pseudomonas aeruginosa* strains isolated from burn patients. *Burns* 2009;36:70–4.

- [53] Naas T, Poirel L, Karim A, Nordmann P. Molecular characterization of In50, a class 1 integron encoding the gene for the extended-spectrum β -lactamase VEB-1 in *Pseudomonas aeruginosa*. FEMS Microbiol Lett 1999;176:411–9.
- [54] Girlich D, Naas T, Leelaporn A, Poirel L, Fennewald M, Nordmann P. Nosocomial spread of the integron-located VEB-1-like cassette encoding an extended-spectrum β -lactamase in *Pseudomonas aeruginosa* in Thailand. Clin Infect Dis 2002;34:603–11.
- [55] Poirel L, Rotimi VO, Mokaddas EM, Karim A, Nordmann P. VEB-1-like extended-spectrum β -lactamases in *Pseudomonas aeruginosa*, Kuwait. Emerg Infect Dis 2001;7:468–70.
- [56] Aubert D, Girlich D, Naas T, Nagarajan S, Nordmann P. Functional and structural characterization of the genetic environment of an extended-spectrum β -lactamase bla_{VEB} gene from a *Pseudomonas aeruginosa* isolate obtained in India. Antimicrob Agents Chemother 2004;48:3284–90.
- [57] Strateva T, Ouzounova-Raykova V, Markova B, Todorova A, Marteva-Proevska Y, Mitov I. Widespread detection of VEB-1-type extended-spectrum β -lactamases among nosocomial ceftazidime-resistant *Pseudomonas aeruginosa* isolates in Sofia, Bulgaria. J Chemother 2007;19:140–5.
- [58] Woodford N, Zhang J, Kaufmann ME, Yarde S, Tomas Mdel M, Faris C, et al. Detection of *Pseudomonas aeruginosa* isolates producing VEB-type extended-spectrum β -lactamases in the United Kingdom. J Antimicrob Chemother 2008;62:1265–8.
- [59] Hansen F, Johansen HK, Ostergaard C, Hansen DS, Littauer P, Homl A, et al. Characterization of carbapenem nonsusceptible *Pseudomonas aeruginosa* in Denmark: a nationwide prospective study. Microb Drug Resist 2014;20:22–9.
- [60] Jiang X, Zhang Z, Li M, Zhou D, Ruan F, Lu Y. Detection of extended-spectrum β -lactamases in clinical isolates of *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 2006;50:2990–5.
- [61] Dubois V, Poirel L, Marie C, Arpin C, Nordmann P, Quentin C. Molecular characterization of a novel class 1 integron containing bla_{GES-1} and a fused product of $aac3-lb/aac6-lb'$ gene cassettes in *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 2002;46:638–45.
- [62] Castanheira M, Mendes RE, Walsh TR, Gales AC, Jones RN. Emergence of the extended-spectrum β -lactamase GES-1 in a *Pseudomonas aeruginosa* strain from Brazil: report from the SENTRY Antimicrobial Surveillance Program. Antimicrob Agents Chemother 2004;48:2344–5.
- [63] Pasteran F, Faccone D, Petroni A, Rapoport M, Galas M, Vazquez M, et al. Novel variant (bla_{VIM-11}) of the metallo- β -lactamase bla_{VIM} family in a GES-1 extended-spectrum- β -lactamase-producing *Pseudomonas aeruginosa* clinical isolate in Argentina. Antimicrob Agents Chemother 2005;49:474–5.
- [64] Mavroidi A, Tzelepi E, Tsakris A, Miriagou V, Sofianou D, Tzouveleki LS. An integron-associated β -lactamase (IBC-2) from *Pseudomonas aeruginosa* is a variant of the extended-spectrum β -lactamase IBC-1. J Antimicrob Chemother 2001;48:627–30.
- [65] Kotsakis SD, Papagiannitsis CC, Tzelepi E, Legakis NJ, Miriagou V, Tzouveleki LS. GES-13, a β -lactamase variant possessing Lys-104 and Asn-170 in *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 2010;54:1331–3.
- [66] Poirel L, Brinas L, Fortineau N, Nordmann P. Integron-encoded GES-type extended-spectrum β -lactamase with increased activity toward aztreonam in *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 2005;49:3593–7.
- [67] Naas T, Philippon L, Poirel L, Ronco E, Nordmann P. An SHV-derived extended-spectrum β -lactamase in *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 1999;43:1281–4.
- [68] Mansour W, Dahmen S, Poirel L, Charfi K, Bettaieb D, Boujaafar N, et al. Emergence of SHV-2a extended-spectrum β -lactamases in clinical isolates of *Pseudomonas aeruginosa* in a university hospital in Tunisia. Microb Drug Resist 2009;15:295–301.
- [69] Chanawong A, M'Zali FH, Heritage J, Lulitanond A, Hawkey PM. SHV-12, SHV-5, SHV-2a and VEB-1 extended-spectrum β -lactamases in Gram-negative bacteria isolated in a university hospital in Thailand. J Antimicrob Chemother 2001;48:839–52.
- [70] Uemura S, Yokota S, Mizuno H, Sakawaki E, Sawamoto K, Maekawa K, et al. Acquisition of a transposon encoding extended-spectrum β -lactamase SHV-12 by *Pseudomonas aeruginosa* isolates during the clinical course of a burn patient. Antimicrob Agents Chemother 2010;54:3956–9.
- [71] Poirel L, Lebossi E, Castro M, Fèvre C, Foustoujou M, Nordmann P. Nosocomial outbreak of extended-spectrum β -lactamase SHV-5-producing isolates of *Pseudomonas aeruginosa* in Athens, Greece. Antimicrob Agents Chemother 2004;48:2277–9.
- [72] Poirel L, Ronco E, Naas T, Nordmann P. Extended-spectrum β -lactamase TEM-4 in *Pseudomonas aeruginosa*. Clin Microbiol Infect 1999;5:651–2.
- [73] Dubois V, Arpin C, Noury P, Quentin C. Clinical strain of *Pseudomonas aeruginosa* carrying a bla_{TEM-21} gene located on a chromosomal interrupted TnA type transposon. Antimicrob Agents Chemother 2002;46:3624–6.
- [74] Marchandin H, Jean-Pierre H, De Champs C, Sirot D, Darbas H, Perigault PF, et al. Production of a TEM-24 plasmid-mediated extended-spectrum β -lactamase by a clinical isolate of *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 2000;44:213–6.
- [75] Mugnier P, Dubrous P, Casin I, Arlet G, Collatz E. A TEM-derived extended-spectrum β -lactamase in *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 1996;40:2488–93.
- [76] Al Naiemi N, Duim B, Bart A. A CTX-M extended-spectrum β -lactamase in *Pseudomonas aeruginosa* and *Stenotrophomonas maltophilia*. J Med Microbiol 2006;55:1607–8.
- [77] Picão RC, Poirel L, Gales AC, Nordmann P. Further identification of CTX-M-2 extended-spectrum β -lactamase in *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 2009;53:2225–6.
- [78] Polotto M, Casella T, de Lucca Oliveira MG, Rúbio FG, Nogueira ML, de Almeida MT, et al. Detection of *Pseudomonas aeruginosa* harboring $bla_{CTX-M-2}$, bla_{GES-1} and bla_{GES-5} , bla_{IMP-1} and bla_{PM-1} causing infections in Brazilian tertiary-care hospital. BMC Infect Dis 2012;12:176.
- [79] Poirel L, Brinas L, Verlinde A, Ide L, Nordmann P. BEL-1, a novel clavulanic acid-inhibited extended-spectrum β -lactamase, and the class 1 integron In120 in *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 2005;49:3743–8.
- [80] Bogaerts P, Bauraing C, Deplano A, Glupczynski Y. Emergence and dissemination of BEL-1-producing *Pseudomonas aeruginosa* isolates in Belgium. Antimicrob Agents Chemother 2007;51:1584–5.
- [81] Poirel L, Docquier J, De Luca F, Verlinde A, Ide L, Rossolini GM, et al. BEL-2, an extended-spectrum β -lactamase with increased activity toward expanded-spectrum cephalosporins in *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 2010;54:533–5.
- [82] Juan C, Zamorano L, Pérez JL, Ge Y, Oliver A. Activity of a new antipseudomonal cephalosporin, CXA-101 (FR264205), against carbapenem-resistant and multidrug-resistant *Pseudomonas aeruginosa* clinical strains. Antimicrob Agents Chemother 2010;54:846–51.
- [83] Tian GB, Adams-Haduch JM, Bogdanovich T, Wang HN, Doi Y. PME-1, an extended-spectrum β -lactamase identified in *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 2011;55:2710–3.
- [84] Bae IK, Lee YN, Jeong SH, Hong SG, Lee JH, Lee SH, et al. Genetic and biochemical characterization of GES-5, an extended-spectrum class A β -lactamase from *Klebsiella pneumoniae*. Diagn Microbiol Infect Dis 2007;58:465–8.
- [85] Bonnin RA, Nordmann P, Potron A, Lécuyer H, Zahar JR, Poirel L. Carbapenem-hydrolyzing GES-type extended-spectrum β -lactamase in *Acinetobacter baumannii*. Antimicrob Agents Chemother 2011;55:349–54.
- [86] Nordmann P, Cuzon G, Naas T. The real threat of *Klebsiella pneumoniae* carbapenemase-producing bacteria. Lancet Infect Dis 2009;9:228–36.
- [87] Naas T, Cuzon G, Villegas MV, Lartigue MF, Quinn JP, Nordmann P. Genetic structures at the origin of acquisition of the β -lactamase bla_{KPC} gene. Antimicrob Agents Chemother 2008;52:1257–63.
- [88] Robledo IE, Aquino EE, Santé MI, Santana JL, Otero DM, Leon CF, et al. Detection of KPC in *Acinetobacter* spp. in Puerto Rico. Antimicrob Agents Chemother 2010;54:1354–7.
- [89] Poirel L, Weldhagen GF, Naas T, De Champs C, Dove MG, Nordmann P. GES-2, a class A β -lactamase from *Pseudomonas aeruginosa* with increased hydrolysis of imipenem. Antimicrob Agents Chemother 2001;45:2598–603.
- [90] Poirel L, Weldhagen GF, De Champs C, Nordmann P. A nosocomial outbreak of *Pseudomonas aeruginosa* isolates expressing the extended-spectrum β -lactamase GES-2 in South Africa. J Antimicrob Chemother 2002;49:561–5.
- [91] Wang C, Cai P, Chang D, Mi AA. *Pseudomonas aeruginosa* isolate producing the GES-5 extended-spectrum β -lactamase. J Antimicrob Chemother 2006;57:1261–2.
- [92] Labuschagne J, Weldhagen GF, Ehlers MM, Dove MG. Emergence of class 1 integron-associated GES-5 and GES-5-like extended-spectrum β -lactamases in clinical isolates of *Pseudomonas aeruginosa* in South Africa. Int J Antimicrob Agents 2008;31:527–30.
- [93] Picão RC, Poirel L, Gales AC, Nordmann P. Diversity of β -lactamases produced by ceftazidime-resistant *Pseudomonas aeruginosa* isolates causing bloodstream infections in Brazil. Antimicrob Agents Chemother 2009;53:3908–13.
- [94] Iraz M, Duzgun AO, Cicek AC, Bonnin RA, Ceylan A, Saral A, et al. Characterization of novel VIM carbapenemase, VIM-38, and first detection of GES-5 carbapenem-hydrolyzing β -lactamases in *Pseudomonas aeruginosa* in Turkey. Diagn Microbiol Infect Dis 2014;78:292–4.
- [95] Bebrone C, Bogaerts P, Delbrück H, Benninck S, Kupper MB, Rezende de Castro R, et al. GES-18, a new carbapenem-hydrolyzing GES-type β -lactamase from *Pseudomonas aeruginosa* that contains Ile80 and Ser170 residues. Antimicrob Agents Chemother 2013;57:396–401.
- [96] Villegas MV, Lolans K, Correa A, Kattan JN, Lopez JA, Quinn JP. First identification of *Pseudomonas aeruginosa* isolates producing a KPC-type carbapenem-hydrolyzing β -lactamase. Antimicrob Agents Chemother 2007;51:1553–5.
- [97] Wolter DJ, Khalaf N, Robledo IE, Vazquez GJ, Santé MI, Aquino EE, et al. Surveillance of carbapenem-resistant *Pseudomonas aeruginosa* isolates from Puerto Rican medical center hospitals: dissemination of KPC and IMP-18 β -lactamases. Antimicrob Agents Chemother 2009;53:1660–4.
- [98] Wolter DJ, Kurpiel PM, Woodford N, Palepou MF, Goering RV, Hanson ND. Phenotypic and enzymatic comparative analysis of the novel KPC variant KPC-5 and its evolutionary variants, KPC-2 and KPC-4. Antimicrob Agents Chemother 2009;53:557–662.
- [99] Akpaka PE, Swanston WH, Ihemere HN, Correa A, Torres JA, Tafur JD, et al. Emergence of KPC-producing *Pseudomonas aeruginosa* in Trinidad and Tobago. J Clin Microbiol 2009;47:2670–1.
- [100] Poirel L, Nordmann P, Lagrutta E, Cleary T, Munoz-Price LS. Emergence of KPC-producing *Pseudomonas aeruginosa* in the United States. Antimicrob Agents Chemother 2010;54:3072.
- [101] Ge C, Wei Z, Jiang Y, Shen P, Yu Y, Chen Y. Identification of KPC-2-producing *Pseudomonas aeruginosa* isolates in China. J Antimicrob Chemother 2011;66:1184–6.
- [102] Cuzon G, Naas T, Villegas MV, Correa A, Quinn JP, Nordmann P. Wide dissemination of *Pseudomonas aeruginosa* producing β -lactamase bla_{KPC-2} gene in Colombia. Antimicrob Agents Chemother 2011;55:5350–3.

- [103] Garcia Ramirez D, Nicola F, Zarate S, Relloso S, Smayevsky J, Arduino D. Emergence of *Pseudomonas aeruginosa* with KPC-type carbapenemase in a teaching hospital: an 8-year study. *J Med Microbiol* 2013;62:1565–70.
- [104] Jacome PR, Alves LR, Cabral AB, Lopes AC, Maciel MA. First report of KPC-producing *Pseudomonas aeruginosa* in Brazil. *Antimicrob Agents Chemother* 2012;56:4990.
- [105] Nordmann P, Dortet L, Poirel L. Carbapenem resistance in Enterobacteriaceae: here is the storm! *Trends Mol Med* 2012;18:263–72.
- [106] Walsh TR, Toleman MA, Poirel L, Nordmann P. Metallo- β -lactamases: the quiet before the storm? *Clin Microbiol Rev* 2005;18:306–25.
- [107] Poirel L, Nordmann P. Carbapenem resistance in *Acinetobacter baumannii*: mechanisms and epidemiology. *Clin Microbiol Infect* 2006;12:826–36.
- [108] Cornaglia G, Riccio ML, Mazzariol A, Lauretti L, Fontana R, Rossolini GM. Appearance of IMP-1 metallo- β -lactamases in Europe. *Lancet* 1999;353:899–900.
- [109] Shibata N, Doi Y, Yamane K, Yagi T, Kurokawa H, Shibayama K, et al. PCR typing of genetic determinants for metallo- β -lactamases and integrases carried by Gram-negative bacteria isolated in Japan, with focus on the class 3 integron. *J Clin Microbiol* 2003;41:5407–13.
- [110] Lee K, Lee WG, Uh Y, Ha GY, Cho J, Chong Y. VIM- and IMP-type metallo- β -lactamase-producing *Pseudomonas* spp. and *Acinetobacter* spp. in Korean hospitals. *Emerg Infect Dis* 2003;9:868–71.
- [111] Uma Karthika R, Srinivasa Rao R, Sahoo S, Shashikala P, Kanungo R, Jayachandran S, et al. Phenotypic and genotypic assays for detecting the prevalence of metallo- β -lactamases in clinical isolates of *Acinetobacter baumannii* from a south Indian tertiary care hospital. *J Med Microbiol* 2009;58:430–5.
- [112] Chiu CH, Lee HY, Tseng LY, Chen CL, Chia JH, Su LH, et al. Mechanisms of resistance to ciprofloxacin, ampicillin/sulbactam and imipenem in *Acinetobacter baumannii* clinical isolates in Taiwan. *Int J Antimicrob Agents* 2010;35:382–6.
- [113] Al-Sweih NA, Al-Hubail M, Rotimi VO. Three distinct clones of carbapenem-resistant *Acinetobacter baumannii* with high diversity of carbapenemases isolated from patients in two hospitals in Kuwait. *J Infect Public Health* 2012;5:102–8.
- [114] Riccio ML, Franceschini N, Boschi L, Caravelli B, Cornaglia G, Fontana R, et al. Characterization of the metallo- β -lactamase determinant of *Acinetobacter baumannii* AC-54/97 reveals the existence of *bla*_{IMP} allelic variants carried by gene cassettes of different phylogeny. *Antimicrob Agents Chemother* 2000;44:1229–35.
- [115] Chu YW, Afzal-Shah M, Houang ET, Palepou MI, Lyon DJ, Woodford N, et al. IMP-4, a novel metallo- β -lactamase from nosocomial *Acinetobacter* spp. collected in Hong Kong between 1994 and 1998. *Antimicrob Agents Chemother* 2001;45:710–4.
- [116] Peleg AY, Franklin C, Bell J, Spelman DW. Emergence of IMP-4 metallo- β -lactamase in a clinical isolate from Australia. *J Antimicrob Chemother* 2004;54:699–700.
- [117] Koh TH, Sng LH, Wang GC, Hsu LY, Zhao Y. IMP-4 and OXA β -lactamases in *Acinetobacter baumannii* from Singapore. *J Antimicrob Chemother* 2007;59:627–32.
- [118] Da Silva GJ, Correia M, Vital C, Ribeiro G, Sousa JC, Leitao R, et al. Molecular characterization of *bla*_{IMP-5}, a new integron-borne metallo- β -lactamase gene from an *Acinetobacter baumannii* nosocomial isolate in Portugal. *FEMS Microbiol Lett* 2002;215:33–9.
- [119] Gales AC, Tognim MC, Reis AO, Jones RN, Sader HS. Emergence of an IMP-like metallo-enzyme in an *Acinetobacter baumannii* clinical strain from a Brazilian teaching hospital. *Diagn Microbiol Infect Dis* 2003;45:77–9.
- [120] Wang H, Guo P, Sun H, Wang H, Yang Q, Chen M, et al. Molecular epidemiology of clinical isolates of carbapenem-resistant *Acinetobacter* spp. from Chinese hospitals. *Antimicrob Agents Chemother* 2007;51:4022–8.
- [121] Kansakar P, Dorji D, Chongtrakool P, Mingmongkolchai S, Mokmake B, Dubbs P. Local dissemination of multidrug-resistant *Acinetobacter baumannii* clones in a Thai hospital. *Microb Drug Resist* 2011;17:109–19.
- [122] Yamamoto M, Nagao M, Matsumura Y, Matsushima A, Ito Y, Takakura S, et al. Interspecies dissemination of a novel class 1 integron carrying *bla*_{IMP-19} among *Acinetobacter* species in Japan. *J Antimicrob Chemother* 2011;66:2480–3.
- [123] Tsakris A, Ikonomidis A, Pournaras S, Tzouveleki LS, Sofianou D, Legakis NJ, et al. VIM-1 metallo- β -lactamase in *Acinetobacter baumannii*. *Emerg Infect Dis* 2006;12:981–3.
- [124] Mammina C, Bonura C, Aleo A, Calà C, Caputo G, Cataldo MC, et al. Characterization of *Acinetobacter baumannii* from intensive care units and home care patients in Palermo, Italy. *Clin Microbiol Infect* 2011;17:E12–5.
- [125] Lee K, Yum JH, Yong D, Lee HM, Kim HD, Docquier JD, et al. Novel acquired metallo- β -lactamase gene, *bla*_{SIM-1}, in a class 1 integron from *Acinetobacter baumannii* clinical isolates from Korea. *Antimicrob Agents Chemother* 2005;49:4485–91.
- [126] Yong D, Toleman MA, Giske CG, Cho HS, Sundman K, Lee K, et al. Characterization of a new metallo- β -lactamase gene, *bla*_{NDM-1}, and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother* 2009;53:5046–54.
- [127] Karthikeyan K, Thirunarayan MA, Krishnan P. Coexistence of *bla*_{OXA-23} with *bla*_{NDM-1} and *armA* in clinical isolates of *Acinetobacter baumannii* from India. *J Antimicrob Chemother* 2010;65:2253–4.
- [128] Krizova L, Bonnin RA, Nordmann P, Nemeč A, Poirel L. Characterization of a multidrug-resistant *Acinetobacter baumannii* strain carrying the *bla*_{NDM-1} and *bla*_{OXA-23} carbapenemase genes from the Czech Republic. *J Antimicrob Chemother* 2012;67:1550–2.
- [129] Pfeifer Y, Wilharm G, Zander E, Wichelhaus TA, Göttig S, Hunfeld KP, et al. Molecular characterization of *bla*_{NDM-1} in an *Acinetobacter baumannii* strain isolated in Germany in 2007. *J Antimicrob Chemother* 2011;66:1998–2001.
- [130] Bogaerts P, Rezende de Castro R, Roisin S, Deplano A, Huang TD, Hallin M, et al. Emergence of NDM-1-producing *Acinetobacter baumannii* in Belgium. *J Antimicrob Chemother* 2012;67:1552–3.
- [131] Bonnin RA, Poirel L, Naas T, Pirs M, Seme K, Schrenzel J, et al. Dissemination of New Delhi metallo- β -lactamase-1-producing *Acinetobacter baumannii* in Europe. *Clin Microbiol Infect* 2012;18:E362–5.
- [132] Chen Y, Zhou Z, Jiang Y, Yu Y. Emergence of NDM-1-producing *Acinetobacter baumannii* in China. *J Antimicrob Chemother* 2011;66:1255–9.
- [133] Nakazawa Y, Li R, Tamura T, Hoshina T, Tamura K, Kawano S, et al. A case of NDM-1-producing *Acinetobacter baumannii* transferred from India to Japan. *J Infect Chemother* 2013;19:330–2.
- [134] Revathi G, Siu LK, Lu PL, Huang LY. First report of NDM-1-producing *Acinetobacter baumannii* in East Africa. *Int J Infect Dis* 2013;17:e1255–8.
- [135] Pilonnetto M, Arend L, Vespero EC, Pelisson M, Chagas TP, Carvalho-Assef AP, et al. First report of NDM-1-producing *Acinetobacter baumannii* sequence type 25 in Brazil. *Antimicrob Agents Chemother* 2014;58:7592–4.
- [136] Bakour S, Touati A, Bachiri T, Sahli F, Tiouit D, Naim M, et al. First report of 16S rRNA methylase ArmA-producing *Acinetobacter baumannii* and rapid spread of metallo- β -lactamase NDM-1 in Algerian hospitals. *J Infect Chemother* 2014;20:696–701.
- [137] Rafei R, Dabboussi F, Hamze M, Eveillard M, Lemarié C, Mallat H, et al. First report of *bla*_{NDM-1}-producing *Acinetobacter baumannii* isolated in Lebanon from civilians wounded during the Syrian war. *Int J Infect Dis* 2014;21:21–3.
- [138] Decusser JW, Jansen C, Nordmann P, Emirian A, Bonnin RA, Anais L, et al. Outbreak of NDM-1-producing *Acinetobacter baumannii* in France, January to May 2013. *Euro Surveill* 2013;18, pii: 20547.
- [139] Bonnin RA, Cuzon G, Poirel L, Nordmann P. Multidrug-resistant *Acinetobacter baumannii* clone, France. *Emerg Infect Dis* 2013;19:822–3.
- [140] Kaase M, Nordmann P, Wichelhaus TA, Gatermann SG, Bonnin RA, Poirel L. NDM-2 carbapenemase in *Acinetobacter baumannii* from Egypt. *J Antimicrob Chemother* 2011;66:1260–2.
- [141] Espinal P, Fugazza G, Lopez Y, Kasma M, Lerman Y, Malhotra-Kumar S, et al. Dissemination of an NDM-2-producing *Acinetobacter baumannii* clone in an Israeli rehabilitation center. *Antimicrob Agents Chemother* 2011;55:5396–8.
- [142] Ghazawi A, Sonnevend A, Bonnin RA, Poirel L, Nordmann P, Hashmey R, et al. NDM-2 carbapenemase-producing *Acinetobacter baumannii* in the United Arab Emirates. *Clin Microbiol Infect* 2012;18:E34–6.
- [143] Espinal P, Poirel L, Carmeli Y, Kaase M, Pal T, Nordmann P, et al. Spread of NDM-2-producing *Acinetobacter baumannii* in the Middle East. *J Antimicrob Chemother* 2013;68:1928–30.
- [144] Bonnin RA, Poirel L, Nordmann P. New Delhi metallo- β -lactamase-producing *Acinetobacter baumannii*: a novel paradigm for spreading antibiotic resistance genes. *Future Microbiol* 2014;9:33–41.
- [145] Queenan AM, Bush K. Carbapenemases: the versatile β -lactamases. *Clin Microbiol Rev* 2007;20:440–58.
- [146] Zhao WH, Chen G, Ito R, Hu ZQ. Relevance of resistance levels to carbapenems and integron-borne *bla*_{IMP-1}, *bla*_{IMP-7}, *bla*_{IMP-10} and *bla*_{VIM-2} in clinical isolates of *Pseudomonas aeruginosa*. *J Med Microbiol* 2009;58:1080–5.
- [147] Martins AF, Zavascki AP, Gaspardo PB, Barth AL. Dissemination of *Pseudomonas aeruginosa* producing SPM-1-like and IMP-1-like metallo- β -lactamases in hospitals from southern Brazil. *Infection* 2007;35:457–60.
- [148] Qu TT, Zhang JL, Wang J, Tao J, Yu YS, Chen YG, et al. Evaluation of phenotypic tests for detection of metallo- β -lactamase-producing *Pseudomonas aeruginosa* strains in China. *J Clin Microbiol* 2009;47:1136–42.
- [149] Ozgumus OB, Caylan R, Tosun I, Sandalli C, Aydin K, Koksai I. Molecular epidemiology of clinical *Pseudomonas aeruginosa* isolates carrying IMP-1 metallo- β -lactamase gene in a university hospital in Turkey. *Microb Drug Resist* 2007;13:191–8.
- [150] Koh TH, Wang GC, Sng LH. Clonal spread of IMP-1-producing *Pseudomonas aeruginosa* in two hospitals in Singapore. *J Clin Microbiol* 2004;42:5378–80.
- [151] Boonkerd N, Pibalpakdi P, Tiloklums M, Niumsup PR. Class 1 integron containing metallo β -lactamase gene *bla*_{IMP-1} in carbapenem-resistant *Pseudomonas aeruginosa* in Thailand. *J Infect Chemother* 2009;15:257–61.
- [152] Bert F, Vanjak D, Leflon-Guibout V, Mrejen S, Delpierre S, Redondo A, et al. IMP-4-producing *Pseudomonas aeruginosa* in a French patient repatriated from Malaysia: impact of early detection and control measures. *Clin Infect Dis* 2007;44:764–5.
- [153] Peleg AY, Franklin C, Bell JM, Spelman DW. Dissemination of the metallo- β -lactamase gene *bla*_{IMP-4} among Gram-negative pathogens in a clinical setting in Australia. *Clin Infect Dis* 2005;41:1549–56.
- [154] Brízio A, Conceição T, Pimentel M, Da Silva G, Duarte A. High-level expression of IMP-5 carbapenemase owing to point mutation in the –35 promoter region of class 1 integron among *Pseudomonas aeruginosa* clinical isolates. *Int J Antimicrob Agents* 2006;27:27–31.
- [155] Ryoo NH, Lee K, Lim JB, Lee YH, Bae IK, Jeong SH. Outbreak by meropenem-resistant *Pseudomonas aeruginosa* producing IMP-6 metallo- β -lactamase in a Korean hospital. *Diagn Microbiol Infect Dis* 2009;63:115–7.
- [156] Chen Y, Sun M, Wang M, Lu Y, Yan Z. Dissemination of IMP-6-producing *Pseudomonas aeruginosa* ST244 in multiple cities in China. *Eur J Clin Microbiol Infect Dis* 2014;33:1181–7.
- [157] Gibb AP, Tribuddharat C, Moore RA, Louie TJ, Krulicki W, Livermore DM, et al. Nosocomial outbreak of carbapenem-resistant *Pseudomonas aeruginosa* with a new *bla*_{IMP} allele, *bla*_{IMP-7}. *Antimicrob Agents Chemother* 2002;46:255–8.

- [158] Ho SE, Subramaniam G, Palasubramaniam S, Navaratnam P. Carbapenem-resistant *Pseudomonas aeruginosa* in Malaysia producing IMP-7 β -lactamase. *Antimicrob Agents Chemother* 2002;46:3286–7.
- [159] Ohlasova D, Kmet V, Niks M. First report of the carbapenem-resistant *Pseudomonas aeruginosa* producing IMP-7 metallo- β -lactamase in Slovakia. *Int J Antimicrob Agents* 2007;30:370–1.
- [160] Kouda S, Kuwahara R, Ohara M, Shigeta M, Fujiwara T, Komatsuzawa H, et al. First isolation of *bla*_{IMP-7} in a *Pseudomonas aeruginosa* in Japan. *J Infect Chemother* 2007;13:276–7.
- [161] Hrabák J, Fridrichová M, Stolbová M, Bergerova T, Zemlickova H, Urbaskova P. First identification of metallo- β -lactamase-producing *Pseudomonas aeruginosa* in the Czech Republic. *Euro Surveill* 2009;14:pii:19102.
- [162] Hammerum AM, Jakobsen L, Hansen F, Stegger M, Sorensen LA, Andersen PS, et al. Characterisation of an IMP-7-producing ST357 *Pseudomonas aeruginosa* isolate detected in Denmark using whole genome sequencing. *Int J Antimicrob Agents* 2015;45:200–1.
- [163] Xiong J, Hynes MF, Ye H, Chen H, Yang Y, M'zali F, et al. *bla*_{IMP-9} and its association with large plasmids carried by *Pseudomonas aeruginosa* isolates from the People's Republic of China. *Antimicrob Agents Chemother* 2006;50:355–8.
- [164] Iyobe S, Kusadokoro H, Takahashi A, Yomoda S, Okubo T, Nakamura A, et al. Detection of a variant metallo- β -lactamase, IMP-10, from two unrelated strains of *Pseudomonas aeruginosa* and an *Alcaligenes xylosoxidans* strain. *Antimicrob Agents Chemother* 2002;46:2014–6.
- [165] Duljasz W, Gniadkowski M, Sitter S, Wojna A, Jebelean C. First organisms with acquired metallo- β -lactamases (IMP-13, IMP-22, and VIM-2) reported in Austria. *Antimicrob Agents Chemother* 2009;53:2221–2.
- [166] Pagani L, Colino C, Migliavacca R, Labonia M, Docquier JD, Nucleo A, et al. Nosocomial outbreak caused by multidrug-resistant *Pseudomonas aeruginosa* producing IMP-13 metallo- β -lactamase. *J Clin Microbiol* 2005;43:3824–8.
- [167] Fournier D, Jeannot K, Robert-Nicoud M, Muller E, Cholley P, van der Mee-Marquet N, et al. Spread of the *bla*_{IMP-13} gene in French *Pseudomonas aeruginosa* through sequence types ST621, ST308 and ST111. *Int J Antimicrob Agents* 2012;40:571–3.
- [168] Naas T, Bogaerts P, Kostyanev T, Cuzon G, Huang TD, Ozsu S, et al. Silent spread of IMP-13-producing *Pseudomonas aeruginosa* belonging to sequence type 621 in Belgium. *J Antimicrob Chemother* 2011;66:2178–9.
- [169] Piyakul C, Tiyawisutrisri R, Boonbumrung K. Emergence of metallo- β -lactamase IMP-14 and VIM-2 in *Pseudomonas aeruginosa* clinical isolates from a tertiary-level hospital in Thailand. *Epidemiol Infect* 2012;140:539–41.
- [170] Quinones-Falconi F, Galicia-Velasco M, Marchiario P, Mussi MA, Ballerini V, Vila AJ. Emergence of *Pseudomonas aeruginosa* strains producing metallo- β -lactamases of the IMP-15 and VIM-2 types in Mexico. *Clin Microbiol Infect* 2010;16:126–31.
- [171] Gilarranz R, Juan C, Castillo-Vera J, Chamizo FJ, Artiles F, Alamo I, et al. First detection in Europe of the metallo- β -lactamase IMP-15 in clinical strains of *Pseudomonas putida* and *Pseudomonas aeruginosa*. *Clin Microbiol Infect* 2013;19:E424–7.
- [172] Castanheira M, Deshpande LM, Costello A, Davies TA, Jones TN. Epidemiology and carbapenem resistance mechanisms of carbapenem-non-susceptible *Pseudomonas aeruginosa* collected during 2009–11 in 14 European and Mediterranean countries. *J Antimicrob Chemother* 2014;69:1804–14.
- [173] Mendes RE, Toleman MA, Ribeiro J, Sader HS, Jones RN, Walsh TR. Integron carrying a novel metallo- β -lactamase gene, *bla*_{IMP-16}, and a fused form of aminoglycoside-resistant gene *aac(6')-30/aac(6')-Ib'*: report from the SEN-TRY Antimicrobial Surveillance Program. *Antimicrob Agents Chemother* 2004;48:4693–702.
- [174] Hanson ND, Hossain A, Buck L, Moland ES, Thomson KS. First occurrence of a *Pseudomonas aeruginosa* isolate in the United States producing an IMP metallo- β -lactamase, IMP-18. *Antimicrob Agents Chemother* 2006;50:2272–3.
- [175] Garza-Ramos U, Tinoco P, Silva-Sanchez J, Morfin-Otero R, Rodriguez-Noriega E, Leon-Garnica G, et al. Metallo- β -lactamase IMP-18 is located in a class 1 integron (In96) in a clinical isolate of *Pseudomonas aeruginosa* from Mexico. *Int J Antimicrob Agents* 2008;31:78–80.
- [176] Pollini S, Antonelli A, Venturelli C, Maradei S, Veggetti A, Bracco S, et al. Acquisition of plasmid-borne *bla*_{IMP-19} gene by a VIM-1-positive *Pseudomonas aeruginosa* of the sequence type 235 epidemic lineage. *J Antimicrob Chemother* 2013;68:722–4.
- [177] Pellegrini C, Mercuri PS, Celenza G, Galleni M, Segatore B, Sacchetti E, et al. Identification of *bla*_{IMP-22} in *Pseudomonas* spp. in urban wastewater and nosocomial environments: biochemical characterization of a new IMP metallo-enzyme variant and its genetic location. *J Antimicrob Chemother* 2009;63:901–8.
- [178] Kim MJ, Bae IK, Jeong SH, Kim SH, Song JH, Choi JY. Dissemination of metallo- β -lactamase-producing *Pseudomonas aeruginosa* of sequence type 235 in Asian countries. *J Antimicrob Chemother* 2013;68:2820–4.
- [179] Koh TH, Khoo CT, Tan TT, Arshad MA, Ang LP, Lau LJ, et al. Multilocus sequence types of carbapenem-resistant *Pseudomonas aeruginosa* in Singapore carrying metallo- β -lactamase genes, including the novel *bla*_{IMP-26} gene. *J Clin Microbiol* 2010;48:2563–4.
- [180] Jeannot K, Poirel L, Robert-Nicoud M, Cholley P, Nordmann P, Plésiat P. IMP-29, a novel IMP-type metallo- β -lactamase in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2012;56:2187–90.
- [181] Pegg KM, Liu EM, Lacuran AE, Oelschlaeger P. Biochemical characterization of IMP-30, a metallo- β -lactamase with enhanced activity toward ceftazidime. *Antimicrob Agents Chemother* 2013;57:5122–6.
- [182] Deshpande LM, Davies TA, Blandino G, Nicoletti G, Jones RN, Castanheira M. IMP-33, a new IMP variant detected in *Pseudomonas aeruginosa* from Sicily. *Antimicrob Agents Chemother* 2013;57:6401–3.
- [183] Pournaras S, Köck R, Mossialos D, Mellmann A, Sakellaris V, Stathopoulos C, et al. Detection of a phylogenetically distinct IMP-type metallo- β -lactamase, IMP-35, in a CC235 *Pseudomonas aeruginosa* from the Dutch–German border region (Euregio). *J Antimicrob Chemother* 2013;68:1271–6.
- [184] Tada T, Miyoshi-Akiyama T, Shimada K, Shimojima M, Kirikae T. IMP-43 and IMP-44 metallo- β -lactamases with increased carbapenemase activities in multidrug-resistant *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2013;57:4427–32.
- [185] Wang Y, Wang X, Schwarz S, Zhang R, Lei L, Liu X, et al. IMP-45-producing multidrug-resistant *Pseudomonas aeruginosa* of canine origin. *J Antimicrob Chemother* 2014;69:2579–81.
- [186] Nordmann P, Poirel L. Emerging carbapenemases in Gram-negative aerobes. *Clin Microbiol Infect* 2002;8:321–31.
- [187] Lauretti L, Riccio ML, Mazzariol A, Cornaglia G, Amicosante G, Fontana R, et al. Cloning and characterization of *bla*_{VIM-4}, a new integron-borne metallo- β -lactamase gene from a *Pseudomonas aeruginosa* clinical isolate. *Antimicrob Agents Chemother* 1999;43:1584–90.
- [188] Corvec S, Poirel L, Decousser JW, Allouch PY, Drugeon H, Nordmann P. Emergence of carbapenem-hydrolysing metallo- β -lactamase VIM-1 in *Pseudomonas aeruginosa* isolates in France. *Clin Microbiol Infect* 2006;12:941–2.
- [189] Tsakris A, Pournaras S, Woodford N, Palepou MF, Babini GS, Douboyas J, et al. Outbreak of infections caused by *Pseudomonas aeruginosa* producing VIM-1 carbapenemase in Greece. *J Clin Microbiol* 2000;38:1290–2.
- [190] Mansour W, Poirel L, Bettaieb D, Bouallegue O, Boujaafar N, Nordmann P. Metallo- β -lactamase-producing *Pseudomonas aeruginosa* isolates in Tunisia. *Diagn Microbiol Infect Dis* 2009;64:458–61.
- [191] Castanheira M, Bell JM, Turnidge JD, Mathai D, Jones RN. Carbapenem resistance among *Pseudomonas aeruginosa* strains from India: evidence for nationwide endemicity of multiple metallo- β -lactamase clones (VIM-2, -5, -6, and -11 and the newly characterized VIM-18). *Antimicrob Agents Chemother* 2009;53:1225–7.
- [192] Pitout JD, Revathi G, Chow BL, Kabera B, Kariuki S, Nordmann P, et al. Metallo- β -lactamase-producing *Pseudomonas aeruginosa* isolated from a large tertiary centre in Kenya. *Clin Microbiol Infect* 2008;14:755–9.
- [193] Libisch B, Watine J, Balogh B, Gacs M, Muzslay M, Szabo G, et al. Molecular typing indicates an important role for two international clonal complexes in dissemination of VIM-producing *Pseudomonas aeruginosa* clinical isolates in Hungary. *Res Microbiol* 2008;159:162–8.
- [194] Khosravi Y, Tee Tay S, Vadivelu J. Metallo- β -lactamase-producing imipenem-resistant *Pseudomonas aeruginosa* clinical isolates in a university teaching hospital in Malaysia: detection of IMP-7 and first identification of IMP-4, VIM-2, and VIM-11. *Diagn Microbiol Infect Dis* 2010;67:294–6.
- [195] Yan JJ, Hsueh PR, Ko WC, Luh KT, Tsai SH, Wu HM, et al. Metallo- β -lactamases in clinical *Pseudomonas* isolates in Taiwan and identification of VIM-3, a novel variant of the VIM-2 enzyme. *Antimicrob Agents Chemother* 2001;45:2224–8.
- [196] Libisch B, Gacs M, Csiszár K, Muzslay M, Rókusz L, Füzi M. Isolation of an integron-borne *bla*_{VIM-4} type metallo- β -lactamase gene from a carbapenem-resistant *Pseudomonas aeruginosa* clinical isolate in Hungary. *Antimicrob Agents Chemother* 2004;48:3576–8.
- [197] Castanheira M, Bell JM, Turnidge JD, Mendes RE, Jones RN. Dissemination and genetic context analysis of *bla*_{VIM-6} among *Pseudomonas aeruginosa* isolates in Asian-Pacific nations. *Clin Microbiol Infect* 2009;16:186–9.
- [198] Toleman MA, Rolston K, Jones RN, Walsh TR. *bla*_{VIM-7}, an evolutionarily distinct metallo- β -lactamase gene in a *Pseudomonas aeruginosa* isolate from the United States. *Antimicrob Agents Chemother* 2004;48:329–32.
- [199] Crespo MP, Woodford N, Sinclair A, Kaufmann ME, Turton J, Glover J, et al. Outbreak of carbapenem-resistant *Pseudomonas aeruginosa* producing VIM-8, a novel metallo- β -lactamase, in a tertiary care center in Cali, Colombia. *J Clin Microbiol* 2004;42:5094–101.
- [200] Marchiario P, Tomatis PE, Mussi MA, Pasteran F, Viale AM, Limansky AS, et al. Biochemical characterization of metallo- β -lactamase VIM-11 from a *Pseudomonas aeruginosa* clinical strain. *Antimicrob Agents Chemother* 2008;52:2250–2.
- [201] Juan C, Beceiro A, Gutiérrez O, Alberti S, Garau M, Perez JL, et al. Characterization of the new metallo- β -lactamase VIM-13 and its integron-borne gene from a *Pseudomonas aeruginosa* clinical isolate in Spain. *Antimicrob Agents Chemother* 2008;52:3589–96.
- [202] Mazzariol A, Mamma C, Koncan R, Di Gaetano V, Di Carlo P, Cipolla D, et al. A novel VIM-type metallo- β -lactamase (VIM-14) in a *Pseudomonas aeruginosa* clinical isolate from a neonatal intensive care unit. *Clin Microbiol Infect* 2011;17:722–4.
- [203] Schneider I, Keuleyan E, Raschofer R, Markovska R, Queenan AM, Bauernfeind A. VIM-15 and VIM-16, two new VIM-2-like metallo- β -lactamases in *Pseudomonas aeruginosa* isolates from Bulgaria and Germany. *Antimicrob Agents Chemother* 2008;52:2977–9.
- [204] Siarkou VI, Vitti D, Protonotariou E, Ikonomidis A, Sofianou D. Molecular epidemiology of outbreak-related *Pseudomonas aeruginosa* strains carrying the novel variant *bla*_{VIM-17} metallo- β -lactamase gene. *Antimicrob Agents Chemother* 2009;53:1325–30.
- [205] Riera E, Cabot G, Mulet X, Garcia-Castillo M, del Campo Juan C, et al. *Pseudomonas aeruginosa* carbapenem resistance mechanisms in Spain: impact

- on the activity of imipenem, meropenem and doripenem. *J Antimicrob Chemother* 2011;66:2022–7.
- [206] El-Mahdy TS. Identification of a novel metallo- β -lactamase VIM-28 located within unusual arrangement of class 1 integron structure in *Pseudomonas aeruginosa* from Egypt. *Jpn J Infect Dis* 2014;67:382–4.
- [207] Poirel L, Magalhaes M, Lopes M, Nordmann P. Molecular analysis of metallo- β -lactamase gene *bla*_{SPM-1}-surrounding sequences from disseminated *Pseudomonas aeruginosa* isolates in Recife, Brazil. *Antimicrob Agents Chemother* 2004;48:1406–9.
- [208] Gales AC, Menezes LC, Silbert S, Sader HS. Dissemination in distinct Brazilian regions of an epidemic carbapenem-resistant *Pseudomonas aeruginosa* producing SPM metallo- β -lactamase. *J Antimicrob Chemother* 2003;52:699–702.
- [209] Salabi AE, Toleman MA, Weeks J, Bruderer T, Frei R, Walsh TR. First report of the metallo- β -lactamase SPM-1 in Europe. *Antimicrob Agents Chemother* 2010;54:582.
- [210] Toleman MA, Bennett PM, Walsh TR. ISCR elements: novel gene-capturing systems of the 21st century? *Microbiol Mol Biol Rev* 2006;70:296–316.
- [211] Castanheira M, Toleman MA, Jones RN, Schmidt FJ, Walsh TR. Molecular characterization of a β -lactamase gene, *bla*_{GIM-1}, encoding a new subclass of metallo- β -lactamase. *Antimicrob Agents Chemother* 2004;48:4654–61.
- [212] Wendel AF, Brodner AH, Wydra S, Ressina S, Henrich B, Pfeffer K, et al. Genetic characterization and emergence of the metallo- β -lactamase GIM-1 in *Pseudomonas* spp. and Enterobacteriaceae during a long-term outbreak. *Antimicrob Agents Chemother* 2013;57:5162–5.
- [213] Jovicic B, Lepsanovic Z, Suljagic V, Rackov G, Begovic J, Topisirovic L, et al. Emergence of NDM-1 metallo- β -lactamase in *Pseudomonas aeruginosa* clinical isolates from Serbia. *Antimicrob Agents Chemother* 2011;55:3929–31.
- [214] Flateau C, Janvier F, Delacour H, Males S, Ficko C, Andriamanantena D, et al. Recurrent pyelonephritis due to NDM-1 metallo- β -lactamase producing *Pseudomonas aeruginosa* in a patient returning from Serbia, France, 2012. *Euro Surveill* 2012;17:pii:20311.
- [215] Janvier F, Jeannot K, Tessé S, Robert-Nicoud M, Delacour H, Rapp C, et al. Molecular characterization of *bla*_{NDM-1} in a sequence type 235 *Pseudomonas aeruginosa* isolate from France. *Antimicrob Agents Chemother* 2013;57:3408–11.
- [216] Khajuria A, Praharaj AK, Kumar M, Grover N. Emergence of NDM-1 in the clinical isolates of *Pseudomonas aeruginosa* in India. *J Clin Diagn Res* 2013;7:1328–31.
- [217] Carattoli A, Fortini D, Galetti R, Garcia-Fernandez A, Nardi G, Orazi D, et al. Isolation of NDM-1-producing *Pseudomonas aeruginosa* sequence type ST235 from a stem cell transplant patient in Italy, May 2013. *Euro Surveill* 2013;18:pii: 20633.
- [218] Zafer MM, Amin M, El Mahallawy H, Ashour MS, Al Agamy M. First report of NDM-1-producing *Pseudomonas aeruginosa* in Egypt. *Int J Infect Dis* 2014;29:80–1.
- [219] Kulkova N, Babalova M, Sokolova J, Krcmery V. First report of New Delhi metallo- β -lactamase-1-producing strains in Slovakia. *Microb Drug Resist* 2015;21:117–20.
- [220] Edelstein MV, Skleenova EN, Shevchenko OV, D'souza JW, Tapalski DV, Azizov IS, et al. Spread of extensively resistant VIM-2-positive ST235 *Pseudomonas aeruginosa* in Belarus, Kazakhstan, and Russia: a longitudinal epidemiological and clinical study. *Lancet Infect Dis* 2013;13:867–76.
- [221] Pollini S, Maradei S, Pecile P, Olivo G, Luzzaro F, Docquier JD, et al. FIM-1, a new acquired metallo- β -lactamase from a *Pseudomonas aeruginosa* clinical isolate from Italy. *Antimicrob Agents Chemother* 2013;57:410–6.
- [222] Bou G, Martínez-Beltrán J. Cloning, nucleotide sequencing, and analysis of the gene encoding an AmpC β -lactamase in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2000;44:428–32.
- [223] Héritier C, Poirel L, Nordmann P. Cephalosporinase over-expression resulting from insertion of ISAb1 in *Acinetobacter baumannii*. *Clin Microbiol Infect* 2006;12:123–30.
- [224] Hujer KM, Hamza NS, Hujer AM, Perez F, Helfand MS, Bethel CR, et al. Identification of a new allelic variant of the *Acinetobacter baumannii* cephalosporinase, ADC-7 β -lactamase: defining a unique family of class C enzymes. *Antimicrob Agents Chemother* 2005;49:2941–8.
- [225] Rodríguez-Martínez JM, Nordmann P, Ronco E, Poirel L. Extended-spectrum cephalosporinase in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2010;54:3484–8.
- [226] Tian GB, Adams-Haduch JM, Taracila M, Bonomo RA, Wang HN, Doi Y. Extended-spectrum AmpC cephalosporinase in *Acinetobacter baumannii*: ADC-56 confers resistance to cefepime. *Antimicrob Agents Chemother* 2011;55:4922–5.
- [227] Bonomo RA, Szabo D. Mechanisms of multidrug resistance in *Acinetobacter* species and *Pseudomonas aeruginosa*. *Clin Infect Dis* 2006;43:49–56.
- [228] Bagge N, Ciofu O, Hentzer M, Campbell JJ, Givskov M, Høiby N. Constitutive high expression of chromosomal β -lactamase in *Pseudomonas aeruginosa* caused by a new insertion sequence (IS1669) located in *ampD*. *Antimicrob Agents Chemother* 2002;46:3406–11.
- [229] Rodríguez-Martínez JM, Poirel L, Nordmann P. Extended-spectrum cephalosporinases in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2009;53:1766–71.
- [230] Rodríguez-Martínez JM, Poirel L, Nordmann P. Molecular epidemiology and mechanisms of carbapenem resistance in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2009;53:4783–8.
- [231] Poirel L, Naas T, Nordmann P. Diversity, epidemiology and genetics of class D β -lactamases. *Antimicrob Agents Chemother* 2010;54:24–38.
- [232] Poirel L, Pitout JD, Nordmann P. Carbapenemases: molecular diversity and clinical consequences. *Future Microbiol* 2007;2:501–12.
- [233] Héritier C, Poirel L, Fournier PE, Claverie JM, Raoult D, Nordmann P. Characterization of the naturally occurring oxacillinase of *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2005;49:4174–9.
- [234] Turton JF, Ward ME, Woodford N, Kaufmann ME, Pike R, Livermore DM, et al. The role of ISAb1 in expression of OXA carbapenemase genes in *Acinetobacter baumannii*. *FEMS Microbiol Lett* 2006;258:72–7.
- [235] Corvec S, Poirel L, Naas T, Drugeon H, Nordmann P. Genetics and expression of the carbapenem-hydrolyzing oxacillinase gene *bla*_{OXA-23} in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2007;51:1530–3.
- [236] Olaitan AO, Berrazeg M, Fagade OE, Adelowo OO, Alli JA, Rolain JM. Emergence of multidrug-resistant *Acinetobacter baumannii* producing OXA-23 carbapenemase, Nigeria. *Int J Infect Dis* 2013;17:e469–70.
- [237] Merino M, Poza M, Roca I, Barba MJ, Sousa MD, Vila J, et al. Nosocomial outbreak of a multiresistant *Acinetobacter baumannii* expressing OXA-23 carbapenemase in Spain. *Microb Drug Resist* 2014;20:259–63.
- [238] Afzal-Shah M, Woodford N, Livermore DM. Characterization of OXA-25, OXA-26, and OXA-27, molecular class D β -lactamases associated with carbapenem resistance in clinical isolates of *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2001;45:583–8.
- [239] Héritier C, Poirel L, Aubert D, Nordmann P. Genetic and functional analysis of the chromosome-encoded carbapenem-hydrolyzing oxacillinase OXA-40 of *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2003;47:268–73.
- [240] Lolans K, Rice TW, Munoz-Price LS, Quinn JP. Multicity outbreak of carbapenem-resistant *Acinetobacter baumannii* isolates producing the carbapenemase OXA-40. *Antimicrob Agents Chemother* 2006;50:2941–5.
- [241] Da Silva GJ, Quintera S, Bértolo E, Sousa JC, Gallego L, Duarte A, et al. Long-term dissemination of an OXA-40 carbapenemase-producing *Acinetobacter baumannii* clone in the Iberian Peninsula. *J Antimicrob Chemother* 2004;54:255–8.
- [242] Tada T, Miyoshi-Akiyama T, Shimada K, Shimojima M, Kirikae T. Dissemination of 16S rRNA methylase ArmA-producing *Acinetobacter baumannii* and emergence of OXA-72 carbapenemase coproducers in Japan. *Antimicrob Agents Chemother* 2014;58:2916–20.
- [243] de Sá Cavalcanti FL, Almeida AC, Vilela MA, de Moraes Junior MA, de Moraes MM, Leal-Balbino TC. Emergence of extensively drug-resistant OXA-72-producing *Acinetobacter baumannii* in Recife, Brazil: risk of clonal dissemination? *Diagn Microbiol Infect Dis* 2013;77:250–1.
- [244] Povilonis J, Seputiene V, Krasauskas R, Juskaite R, Miskinyte M, Suziedelis K, et al. Spread of carbapenem-resistant *Acinetobacter baumannii* carrying a plasmid with two genes encoding OXA-72 carbapenemase in Lithuanian hospitals. *J Antimicrob Chemother* 2013;68:1000–6.
- [245] Goic-Barisic I, Towner KJ, Kovacic A, Sisko-Kraljevic K, Tonkic M, Novak A, et al. Outbreak in Croatia caused by a new carbapenem-resistant clone of *Acinetobacter baumannii* producing OXA-72 carbapenemase. *J Hosp Infect* 2011;77:368–9.
- [246] Poirel L, Marqué S, Héritier C, Segonds C, Chabanon G, Nordmann P. OXA-58, a novel class D β -lactamase involved in resistance to carbapenems in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2005;49:202–8.
- [247] Héritier C, Dubouix A, Poirel L, Marty N, Nordmann P. A nosocomial outbreak of *Acinetobacter baumannii* isolates expressing the carbapenem-hydrolyzing oxacillinase OXA-58. *J Antimicrob Chemother* 2005;55:115–8.
- [248] Poirel L, Mansour W, Bouallegue O, Nordmann P. Carbapenem-resistant *Acinetobacter baumannii* isolates from Tunisia producing the OXA-58-like carbapenem-hydrolyzing oxacillinase OXA-97. *Antimicrob Agents Chemother* 2008;52:1613–7.
- [249] Higgins PG, Poirel L, Lehmann M, Nordmann P, Seifert H. OXA-143, a novel carbapenem-hydrolyzing class D β -lactamase in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2009;53:5035–8.
- [250] Higgins PG, Pérez-Llarena FJ, Zander E, Fernández A, Bou G, Seifert H. OXA-235, a novel class D β -lactamase involved in resistance to carbapenems in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2013;57:2121–6.
- [251] Girlich D, Naas T, Nordmann P. Biochemical characterization of the naturally occurring oxacillinase OXA-50 of *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2004;48:2043–8.
- [252] Danel F, Hall LM, Gur D, Livermore DM. OXA-15, an extended-spectrum variant of OXA-2 β -lactamase, isolated from a *Pseudomonas aeruginosa* strain. *Antimicrob Agents Chemother* 1997;41:785–90.
- [253] Poirel L, Gerome P, De Champs C, Stephanazzi J, Naas T, Nordmann P. Integron-located *oxa-32* gene cassette encoding an extended-spectrum variant of OXA-2 β -lactamase from *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2002;46:566–9.
- [254] Juan C, Mulet X, Zamorano L, Albertí S, Pérez JL, Oliver A. Detection of the novel extended-spectrum β -lactamase OXA-161 from a plasmid-located integron in *Pseudomonas aeruginosa* clinical isolates from Spain. *Antimicrob Agents Chemother* 2009;53:5288–90.
- [255] Hall LM, Livermore DM, Gur D, Akova M, Akalin HE. OXA-11, an extended-spectrum variant of OXA-10 (PSE-2) β -lactamase from *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 1993;37:1637–44.
- [256] Danel F, Hall LM, Gur D, Livermore DM. OXA-14, another extended-spectrum variant of OXA-10 (PSE-2) β -lactamase from *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 1995;39:1881–4.

- [257] Danel F, Hall LM, Gur D, Livermore DM. OXA-16, a further extended-spectrum variant of OXA-10 β -lactamase, from two *Pseudomonas aeruginosa* isolates. *Antimicrob Agents Chemother* 1998;42:3117–22.
- [258] Danel F, Hall LM, Duke B, Gur D, Livermore DM. OXA-17, a further extended-spectrum variant of OXA-10 β -lactamase, isolated from *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 1999;43:1362–6.
- [259] Mugnier P, Casin I, Bouthors AT, Collatz E. Novel OXA-10-derived extended-spectrum β -lactamases selected in vivo or in vitro. *Antimicrob Agents Chemother* 1998;42:3113–6.
- [260] Poirel L, Girlich D, Naas T, Nordmann P. OXA-28, an extended-spectrum variant of OXA-10 β -lactamase from *Pseudomonas aeruginosa* and its plasmid- and integron-located gene. *Antimicrob Agents Chemother* 2001;45:447–53.
- [261] Liu W, Liu X, Liao J, Zhang Y, Liang X. Identification of *bla*_{OXA-128} and *bla*_{OXA-129}, two novel OXA-type extended-spectrum- β -lactamases in *Pseudomonas aeruginosa*, in Hunan Province, China. *J Basic Microbiol* 2010;50(Suppl. 1):S116–9.
- [262] Hocquet D, Colomb M, Dehecq B, Belmonte O, Courvalin P, Plésiat P, et al. Ceftazidime-hydrolyzing β -lactamase OXA-145 with impaired hydrolysis of penicillins in *Pseudomonas aeruginosa*. *J Antimicrob Chemother* 2011;66:1745–50.
- [263] Aubert D, Poirel L, Chevalier J, Leotard S, Pages JM, Nordmann P. Oxacillinase-mediated resistance to cefepime and susceptibility to ceftazidime in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2001;45:1615–20.
- [264] Philippon LN, Naas T, Bouthors AT, Barakett V, Nordmann P. OXA-18, a class D clavulanic acid-inhibited extended-spectrum β -lactamase from *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 1997;41:2188–95.
- [265] Toleman MA, Rolston K, Jones RN, Walsh TR. Molecular and biochemical characterization of OXA-45, an extended-spectrum class 2d' β -lactamase in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2003;47:2859–63.
- [266] Li H, Walsh TR, Toleman MA. Molecular analysis of the sequences surrounding *bla*_{OXA-45} reveals acquisition of this gene by *Pseudomonas aeruginosa* via a novel ISCR element, ISCR5. *Antimicrob Agents Chemother* 2009;53:1248–51.
- [267] Sevillano E, Gallego L, Garcia-Lobo JM. First detection of the OXA-40 carbapenemase in *Pseudomonas aeruginosa* isolates, located on a plasmid also found in *Acinetobacter baumannii*. *Pathol Biol* 2009;57:493–5.
- [268] El Garch F, Bogaerts P, Bebrone C, Galleni M, Glupczynski Y. OXA-198, an acquired carbapenem-hydrolyzing class D β -lactamase from *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2011;55:4828–33.
- [269] Doi Y, Arakawa Y. 16S ribosomal RNA methylation: emerging resistance mechanism against aminoglycosides. *Clin Infect Dis* 2007;45:88–94.
- [270] Doi Y, Wachino J, Arakawa Y. Nomenclature of plasmid-mediated 16S rRNA methylases responsible for panaminoglycoside resistance. *Antimicrob Agents Chemother* 2008;52:2287–8.
- [271] Wachino J, Arakawa Y. Exogenously acquired 16S rRNA methyltransferases found in aminoglycoside-resistant pathogenic Gram-negative bacteria: an update. *Drug Resist Updat* 2012;15:133–48.
- [272] Galimand M, Courvalin P, Lambert T, RmtF, a new member of the aminoglycoside resistance 16S rRNA N7 G1405 methyltransferase family. *Antimicrob Agents Chemother* 2012;56:3960–2.
- [273] Bueno MF, Francisco GR, O'Hara JA, de Oliveira Garcia D, Doi Y. Coproduction of 16S rRNA methyltransferase RmtD or RmtG with KPC-2 and CTX-M group extended-spectrum β -lactamases in *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 2013;57:2397–400.
- [274] Zhou H, Du XX, Yang Q, Zhou JY, Yu YS, Li LJ. Study on carbapenemase and 16S rRNA methylase of imipenem-resistant *Acinetobacter baumannii*. *Zhonghua Liu Xing Bing Xue Za Zhi* 2009;30:269–72 [in Chinese].
- [275] Cho YJ, Moon DC, Jin JS, Choi CH, Lee YC, Lee JC. Genetic basis of resistance to aminoglycosides in *Acinetobacter* spp. and spread of *armA* in *Acinetobacter baumannii* sequence group 1 in Korean hospitals. *Diagn Microbiol Infect Dis* 2009;64:185–90.
- [276] Hong SB, Shin KS, Ha J, Han K. Co-existence of *bla*_{OXA-23} and *armA* in multidrug-resistant *Acinetobacter baumannii* isolated from a hospital in South Korea. *J Med Microbiol* 2013;62:836–44.
- [277] Tada T, Miyoshi-Akiyama T, Kato Y, Ohmagari N, Takeshita N, Hung NV, et al. Emergence of 16S rRNA methylase-producing *Acinetobacter baumannii* and *Pseudomonas aeruginosa* isolates in hospitals in Vietnam. *BMC Infect Dis* 2013;13:251–6.
- [278] Yamada Y, Suwabe A. Diverse carbapenem-resistance mechanisms in 16S rRNA methylase-producing *Acinetobacter baumannii*. *J Med Microbiol* 2013;62:618–22.
- [279] Doi Y, Adams JM, Yamane K, Paterson DL. Identification of 16S rRNA methylase-producing *Acinetobacter baumannii* clinical strains in North America. *Antimicrob Agents Chemother* 2007;51:4209–10.
- [280] Karah N, Haldorsen B, Hermansen NO, Tveten Y, Ragnhildstveit E, Skutlaberg DH, et al. Emergence of OXA-carbapenemase- and 16S rRNA methylase-producing international clones of *Acinetobacter baumannii* in Norway. *J Med Microbiol* 2011;60:515–21.
- [281] Brigante G, Migliavacca R, Bramati S, Motta E, Nucleo E, Nanenti M, et al. Emergence and spread of a multidrug-resistant *Acinetobacter baumannii* clone producing both the carbapenemase OXA-23 and the 16S rRNA methylase ArmA. *J Med Microbiol* 2012;61:653–61.
- [282] Strateva T, Markova B, Marteva-Proevska Y, Ivanova D, Mitov I. Widespread dissemination of multidrug-resistant *Acinetobacter baumannii* producing OXA-23 carbapenemase and ArmA 16S ribosomal RNA methylase in a Bulgarian university hospital. *Braz J Infect Dis* 2012;16:307–10.
- [283] Aghazadeh M, Rezaee MA, Nahaei MR, Mahdian R, Pajand O, Saffari F, et al. Dissemination of aminoglycoside-modifying enzymes and 16S rRNA methylases among *Acinetobacter baumannii* and *Pseudomonas aeruginosa* isolates. *Microb Drug Resist* 2013;19:282–8.
- [284] Yokoyama K, Doi Y, Yamane K, Kurokawa H, Shibata N, Shibayama K, et al. Acquisition of 16S rRNA methylase gene in *Pseudomonas aeruginosa*. *Lancet* 2003;362:1888–93.
- [285] Yamane K, Doi Y, Yokoyama K, Yagi T, Kurokawa H, Shibata N, et al. Genetic environments of the *rmtA* gene in *Pseudomonas aeruginosa* clinical isolates. *Antimicrob Agents Chemother* 2004;48:2069–74.
- [286] Jin JS, Kwon KT, Moon DC, Lee JC. Emergence of 16S rRNA methylase *rmtA* in colistin-only-sensitive *Pseudomonas aeruginosa* in South Korea. *Int J Antimicrob Agents* 2009;33:490–1.
- [287] Doi Y, de Oliveira Garcia D, Adams J, Paterson DL. Coproduction of novel 16S rRNA methylase RmtD and metallo- β -lactamase SPM-1 in a panresistant *Pseudomonas aeruginosa* isolate from Brazil. *Antimicrob Agents Chemother* 2007;51:852–6.
- [288] Doi Y, Ghilardi AC, Adams J, de Oliveira Garcia D, Paterson DL. High prevalence of metallo- β -lactamase and 16S rRNA methylase coproduction among imipenem-resistant *Pseudomonas aeruginosa* isolates in Brazil. *Antimicrob Agents Chemother* 2007;51:3388–90.
- [289] Gurung M, Moon DC, Tamang MD, Kim J, Lee YC, Seol SY, et al. Emergence of 16S rRNA methylase gene *armA* and carriage of *bla*_{IMP-1} in *Pseudomonas aeruginosa* isolates from South Korea. *Diagn Microbiol Infect Dis* 2010;68:468–70.
- [290] Magnet S, Courvalin P, Lambert T. Resistance-nodulation-cell division-type efflux pump involved in aminoglycoside resistance in *Acinetobacter baumannii* strain BM4454. *Antimicrob Agents Chemother* 2001;45:3375–80.
- [291] Damier-Piolle L, Magnet S, Brémont S, Lambert T, Courvalin P, AdelJK, a resistance-nodulation-cell division pump effluxing multiple antibiotics in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2008;52:557–62.
- [292] Su XZ, Chen J, Mizushima T, Kuroda T, Abe M, Tsuchiya T. An H⁺-coupled *Acinetobacter baumannii* multidrug efflux pump belonging to the MATE family of transporters. *Antimicrob Agents Chemother* 2005;49:4362–4.
- [293] Li XZ, Nikaido H, Poole K. Role of *mexA-mexB-oprM* in antibiotic efflux in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 1995;39:1948–53.
- [294] Poole K, Gotoh N, Tsujimoto H, Zhao Q, Wada A, Yamasaki T, et al. Overexpression of the *mexC-mexD-oprJ* efflux operon in *nfxB*-type multidrug-resistant strains of *Pseudomonas aeruginosa*. *Mol Microbiol* 1996;21:713–24.
- [295] Llanes C, Köhler T, Patry I, Dehecq B, van Delden C, Plésiat P. Role of the MexEF-OprN efflux system in low-level resistance of *Pseudomonas aeruginosa* to ciprofloxacin. *Antimicrob Agents Chemother* 2011;55:5676–84.
- [296] Hawkey P, Finch R. Tigecycline: in-vitro performance as a predictor of clinical efficacy. *Clin Microbiol Infect* 2007;13:354–62.
- [297] Ruzin A, Keeney D, Bradford PA. AdeABC multidrug efflux pump is associated with decreased susceptibility to tigecycline in *Acinetobacter calcoaceticus-Acinetobacter baumannii* complex. *J Antimicrob Chemother* 2007;59:1001–4.
- [298] Coyne S, Rosenfeld N, Lambert T, Courvalin P, Perichon B. Overexpression of resistance-nodulation-cell division pump AdeFGH confers multidrug resistance in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2010;54:4389–93.
- [299] Coyne S, Courvalin P, Périchon B. Efflux-mediated antibiotic resistance in *Acinetobacter* spp. *Antimicrob Agents Chemother* 2011;55:947–53.
- [300] Rumbo C, Gato E, López M, Ruiz de Alegría C, Fernández-Cuenca F, Martínez-Martínez L, et al. Contribution of efflux pumps, porins, and β -lactamases to multidrug resistance in clinical isolates of *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2013;57:5247–57.
- [301] Reid GE, Grim SA, Aldeza CA, Janda WM, Clark NM. Rapid development of *Acinetobacter baumannii* resistance to tigecycline. *Pharmacotherapy* 2007;27:1198–201.
- [302] Chen Q, Li X, Zhou H, Jiang Y, Chen Y, Hua X, et al. Decreased susceptibility to tigecycline in *Acinetobacter baumannii* mediated by a mutation in *trm* encoding SAM-dependent methyltransferase. *J Antimicrob Chemother* 2014;69:72–6.
- [303] Li J, Rayner CR, Nation RL, Owen RJ, Spelman D, Tan KE, et al. Heteroresistance to colistin in multidrug-resistant *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2006;50:2946–50.
- [304] Adams MD, Nickel GC, Bajaksouzian S, Lavender H, Murthy AR, Jacobs MR, et al. Resistance to colistin in *Acinetobacter baumannii* associated with mutations in the PmrAB two-component system. *Antimicrob Agents Chemother* 2009;53:3628–34.
- [305] Beceiro A, Llobet E, Aranda J, Bengoechea JA, Doumith M, Hornsey M, et al. Phosphoethanolamine modification of lipid A in colistin-resistant variants of *Acinetobacter baumannii* mediated by the *pmrAB* two-component regulatory system. *Antimicrob Agents Chemother* 2011;55:3370–9.
- [306] Moffatt JH, Harper M, Harrison P, Hale JD, Vinogradov E, Seemann T, et al. Colistin resistance in *Acinetobacter baumannii* is mediated by complete loss of lipopolysaccharide production. *Antimicrob Agents Chemother* 2010;54:4971–7.
- [307] Hindler JA, Humphries RM. Colistin MIC variability by method for contemporary clinical isolates of multidrug-resistant Gram-negative bacilli. *J Clin Microbiol* 2013;51:1678–84.

- [308] Ko KS, Suh JY, Kwon KT, Jung SI, Park KH, Kang CI, et al. High rates of resistance to colistin and polymyxin B in subgroups of *Acinetobacter baumannii* isolates from Korea. *J Antimicrob Chemother* 2007;60:1163–7.
- [309] Valencia R, Arroyo LA, Conde M, Aldana JM, Torres MJ, Fernandez-Cuenca F, et al. Nosocomial outbreak of infection with pan-drug-resistant *Acinetobacter baumannii* in a tertiary care university hospital. *Infect Control Hosp Epidemiol* 2009;30:257–63.
- [310] Bahador A, Taheri M, Pourakbari B, Hashemizadeh Z, Rostami H, Mansoori N, et al. Emergence of rifampicin, tigecycline, and colistin-resistant *Acinetobacter baumannii* in Iran; spreading of MDR strains of novel international clone variants. *Microb Drug Resist* 2013;19:397–406.
- [311] Lesho E, Yoon EJ, McGann P, Snesrud E, Kwak Y, Milillo M, et al. Emergence of colistin-resistance in extremely drug-resistant *Acinetobacter baumannii* containing a novel *pmrCAB* operon during colistin therapy of wound infections. *J Infect Dis* 2013;208:1142–51.
- [312] Tan CH, Li J, Nation RL. Activity of colistin against heteroresistant *Acinetobacter baumannii* and emergence of resistance in an in vitro pharmacokinetic/pharmacodynamic model. *Antimicrob Agents Chemother* 2007;51:3413–5.
- [313] Denton M, Kerr K, Mooney L, Keer V, Rajgopal A, Brownlee K, et al. Transmission of colistin-resistant *Pseudomonas aeruginosa* between patients attending a pediatric cystic fibrosis center. *Pediatr Pulmonol* 2002;34:257–61.
- [314] Landman D, Bratu S, Alam M, Quale J. Citywide emergence of *Pseudomonas aeruginosa* strains with reduced susceptibility to polymyxin B. *J Antimicrob Chemother* 2005;55:954–7.
- [315] Lee JY, Song JH, Ko KS. Identification of nonclonal *Pseudomonas aeruginosa* isolates with reduced colistin susceptibility in Korea. *Microb Drug Resist* 2011;17:299–304.
- [316] Moskowitz SM, Ernst RK, Miller SI. PmrAB, a two-component regulatory system of *Pseudomonas aeruginosa* that modulates resistance to cationic antimicrobial peptides and addition of aminoarabinose to lipid A. *J Bacteriol* 2004;186:575–9.
- [317] Barrow K, Kwon DH. Alterations in two-component regulatory systems of *phoPQ* and *pmrAB* are associated with polymyxin B resistance in clinical isolates of *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2009;53:5150–4.
- [318] Gutu AD, Sgambati N, Strasbourger P, Brannon MK, Jacobs MA, Haugen E, et al. Polymyxin resistance of *Pseudomonas aeruginosa phoQ* mutants is dependent on additional two-component regulatory systems. *Antimicrob Agents Chemother* 2013;57:2204–15.
- [319] Lee JY, Na IY, Park YK, Ko KS. Genomic variations between colistin-susceptible and -resistant *Pseudomonas aeruginosa* clinical isolates and their effects on colistin resistance. *J Antimicrob Chemother* 2014;69:1248–56.