

**Mononuclear silver(I) complexes with 1,7-phenanthroline as potent
inhibitors of *Candida* growth**

Nada D. Savić^{a,#}, Sandra Vojnović^{b,#}, Biljana Đ. Glišić^a, Aurélien Crochet^c, Aleksandar Pavić^b, Goran V. Janjić^d, Marina Pekmezović^e, Igor M. Opsenica^f, Katharina M. Fromm^{c,*}, Jasmina Nikodinovic-Runic^{b,*}, Miloš I. Djuran^{g,*}

^a*University of Kragujevac, Faculty of Science, Department of Chemistry, R. Domanovića 12, 34000 Kragujevac, Serbia*

^b*Institute of Molecular Genetics and Genetic Engineering, University of Belgrade, Vojvode Stepe 444a, 11000 Belgrade*

^c*Department of Chemistry, University of Fribourg, Chemin du Musée 9, CH-1700 Fribourg, Switzerland*

^d*Institute of Chemistry, Metallurgy and Technology, University of Belgrade, Njegoševa 12, 11000 Belgrade, Serbia*

^e*Department of Microbial Pathogenicity Mechanisms, Hans Knöll Institute, Jena, Germany*

^f*University of Belgrade-Faculty of Chemistry, Studentski trg 16, 11158 Belgrade, Serbia*

^g*Serbian Academy of Sciences and Arts, Knez Mihailova 35, 11000 Belgrade, Serbia*

[#]N.D.S. and S.V. contributed equally.

*Corresponding authors. Tel.: +41 26 300 8732 (K. M. Fromm); Tel.: +381 11 397 6034; fax: +381 11 397 5808 (J. Nikodinovic-Runic); Tel.: +381 34 300 251; fax: +381 34 335 040 (M. I. Djuran).

E-mail address: katharina.fromm@unifr.ch (K. M. Fromm);

jasmina.nikodinovic@gmail.com (J. Nikodinovic-Runic); djuran@kg.ac.rs (M. I. Djuran).

Abstract

Mononuclear silver(I) complexes with 1,7-phenanthroline (1,7-phen), [Ag(NO₃-O,O')(1,7-phen-N7)₂] (**1**) and [Ag(1,7-phen-N7)₂]X, X = ClO₄⁻ (**2**), CF₃SO₃⁻ (**3**), BF₄⁻ (**4**) and SbF₆⁻ (**5**) were synthesized and structurally characterized by NMR (¹H and ¹³C), IR and UV-Vis spectroscopy and ESI mass spectrometry. The crystal structures of **1**, **3** and **4** were determined by single-crystal X-ray diffraction analysis. In all these complexes, 1,7-phen coordinates to the Ag(I) ion in a monodentate fashion *via* the less sterically hindered N7 nitrogen atom. The investigation of the solution stability of **1** – **5** in DMSO revealed that they are sufficiently stable in this solvent at room temperature. Complexes **1** – **5** showed selectivity towards *Candida* spp. in comparison to bacteria, effectively inhibiting the growth of four different *Candida* species with minimal inhibitory concentrations (MIC) between 1.2 and 11.3 μM. Based on the lowest MIC values and the lowest cytotoxicity against healthy human fibroblasts with selectivity index of more than 30, the antifungal potential was examined in detail for the complex **1**. It had the ability to attenuate *C. albicans* virulence and to reduce epithelial cell damage in the cell infection model. Induction of reactive oxygen species (ROS) response has been detected in *C. albicans*, with fungal DNA being one of the possible target biomolecules. The toxicity profile of **1** in the zebrafish model (*Danio rerio*) revealed improved safety and activity in comparison to that of clinically utilized silver(I) sulfadiazine.

Keywords: Silver(I) complexes; Antimicrobial activity; Cytotoxicity; *Candida*; *Danio rerio*

TABLE OF CONTENTS

¹ H NMR spectrum of 1	S5
¹³ C NMR spectrum of 1	S6
¹ H NMR spectrum of 2	S7
¹³ C NMR spectrum of 2	S8
¹ H NMR spectrum of 3	S9
¹³ C NMR spectrum of 3	S10
¹ H NMR spectrum of 4	S11
¹³ C NMR spectrum of 4	S12
¹ H NMR spectrum of 5	S13
¹³ C NMR spectrum of 5	S14

Fig. S1. Molecular structure of the silver(I) complex **3**. Non-coordinating triflate anion in **3** is omitted for clarity. Displacement ellipsoids are drawn at 50% probability level and H atoms are represented by spheres of arbitrary size. S15

Fig. S2. Silver(I) complex **3** stability over time measured by ¹H NMR spectroscopy. ¹H NMR spectrum was measured immediately (**a**) and 48 h (**b**) after complex dissolution in DMSO-*d*₆. S16

Fig. S3. Air/light stability of silver(I) complexes **1** – **5**. S17

Fig. S4. The most stable binding of the tested compounds to base pairs of DNA and the corresponding binding energies, as assessed by molecular docking. S18

Fig. S5. The heartbeating rate of zebrafish embryos at 114 hpf upon different concentrations of complex **1**, 1,7-phen and silver(I) sulfadiazine (AgSD). Statistically significant differences between data of untreated control and treatments were denoted with asterisk (*P < 0.5, **P < 0.01; ***P < 0.001; S19

Student t' test).

Table S1

Selected bond distances (Å) and valence angles (°) in the silver(I) complexes **1**, **3** S20
and **4**.


Table S2

Details of the crystal structure determinations of the silver(I) complexes **1**, **3** and S21
4.


Table S3


Lethal and teratogenic effects observed in zebrafish (*Danio rerio*) embryos at S22
different hours post fertilization (hpf).



^1H NMR (400 MHz, $\text{DMF-}d_7$)




$[\text{Ag}(\text{NO}_3\text{-O,O})(1,7\text{-phen-N}7)_2]$ (**1**)





^{13}C NMR (101 MHz, $\text{DMF-}d_7$)


^1H NMR (400 MHz, $\text{DMF-}d_7$)**[Ag(1,7-phen- $N7$) $_2$] ClO_4 (2)**

^{13}C NMR (101 MHz, $\text{DMF-}d_7$)


¹H NMR (400 MHz, DMF-*d*₇)



^{13}C NMR (101 MHz, DMF- d_7)




^1H NMR (400 MHz, $\text{DMF-}d_7$)




S11


^{13}C NMR (101 MHz, $\text{DMF-}d_7$)




^1H NMR (400 MHz, $\text{DMF-}d_7$)



$[\text{Ag}(1,7\text{-phen-}N7)_2]\text{SbF}_6$ (**5**)



^{13}C NMR (101 MHz, $\text{DMF-}d_7$)




Fig. S1. Molecular structure of the silver(I) complex **3**. Non-coordinating triflate anion in **3** is omitted for clarity. Displacement ellipsoids are drawn at 50% probability level and H atoms are represented by spheres of arbitrary size.




Fig. S2. Silver(I) complex **3** stability over time measured by ^1H NMR spectroscopy. ^1H NMR spectrum was measured immediately (**a**) and 48 h (**b**) after complex dissolution in $\text{DMSO}-d_6$.




Fig. S3. Air/light stability of silver(I) complexes **1 – 5**.




Fig. S4. The most stable binding of the tested compounds to base pairs of DNA and the corresponding binding energies, as assessed by molecular docking.




Fig. S5. The heartbeating rate of zebrafish embryos at 114 hpf upon different concentrations of complex **1**, 1,7-phen and silver(I) sulfadiazine (AgSD). Statistically significant differences between data of untreated control and treatments were denoted with asterisk (* $P < 0.5$, ** $P < 0.01$; *** $P < 0.001$; Student t' test).

Table S1Selected bond distances (Å) and valence angles (°) in the silver(I) complexes **1**, **3** and **4**.

	1	3	4
Ag—N1	2.203(4)	2.137(10)	2.172(7)
Ag—N3	2.206(4)	2.146(9)	2.191(7)
Ag—O2	2.630(4)	/	/
Ag—O3	2.652(4)	/	/
N1—Ag—N3	154.78(15)	174.2(3)	166.6(3)
N1—Ag—O3	95.33(15)	/	/
N3—Ag—O2	93.83(15)	/	/
O2—Ag—O3	47.85(12)	/	/
N1—Ag—O2	110.83(15)	/	/
N3—Ag—O3	105.53(15)	/	/
N5—O3—Ag	95.1(3)	/	/
N5—O2—Ag	95.8(3)	/	/
C1—N1—Ag	120.6(3)	118.4(8)	121.5(6)
C5—N1—Ag	121.8(3)	121.7(8)	119.6(5)
C13—N3—Ag	119.5(3)	119.2(8)	119.1(6)
C17—N3—Ag	122.7(3)	120.1(7)	120.1(5)

Table S2Details of the crystal structure determinations of the silver(I) complexes **1**, **3** and **4**.

	1	3	4
Empirical formula	C ₂₄ H ₁₆ AgN ₅ O ₃	C ₂₅ H ₁₆ AgF ₃ N ₄ O ₃ S	C ₂₄ H ₁₆ AgBF ₄ N ₄
CCDC number	CCDC-1829501	CCDC-1829502	CCDC-1829503
Formula weight (g.mol ⁻¹)	530.29	617.35	555.09
Crystal system, space group	triclinic, <i>P</i> $\bar{1}$	triclinic, <i>P</i> $\bar{1}$	triclinic, <i>P</i> $\bar{1}$
<i>a</i> (Å)	7.1629(7)	6.9800(5)	7.1759(8)
<i>b</i> (Å)	10.5142(9)	13.6380(9)	10.9209(12)
<i>c</i> (Å)	14.5205(12)	13.9579(9)	14.5568(15)
α (°)	78.215(7)	109.122(5)	83.120(9)
β (°)	81.410(7)	103.327(5)	86.671(9)
γ (°)	73.611(7)	103.352(5)	71.989(9)
<i>V</i> (Å ³)	1022.11(16)	1151.72(14)	1076.8(2)
<i>F</i> ₀₀₀	532	616	552
<i>Z</i>	2	2	2
X-radiation, λ / Å	Mo- <i>K</i> α 0.71073	Mo- <i>K</i> α 0.71073	Mo- <i>K</i> α 0.71073
data collect. temperatur. /K	298(2)	298(2)	250(2)
Calculated density (Mg m ⁻³)	1.723	1.780	1.712
Absorption coefficient (mm ⁻¹)	1.026	1.028	0.991
Crystal size (mm)	0.400 × 0.100 × 0.020	0.29 x 0.13 x 0.04	0.210 × 0.130 × 0.030
θ range (°)	1.4 to 25.2	1.6 to 25.1	1.4 to 25.3
index ranges <i>h</i> , <i>k</i> , <i>l</i>	-8 ... 8, -12 ... 12, -17 ... 17	-7 ... 8, -16... 16, -16... 16	-8 ... 8, -13 ... 13, -4 ... 17
No. of collected and independent reflections	13116, 3640	14162, 4106	3842, 3842
<i>R</i> _{int}	0.1022	0.1034	
Data / restraints / parameters	3640 / 0 / 298	4106/151/334	3842 / 0 / 308
Goodness-on-fit on <i>F</i> ²	0.990	0.993	0.915
Final <i>R</i> indices	0.0481, 0.1111	0.0868, 0.2212	0.0699, 0.1669
[<i>F</i> _o > 4σ(<i>F</i> _o)] <i>R</i> (<i>F</i>), <i>wR</i> (<i>F</i> ²)			
Final <i>R</i> indices (all data) <i>R</i> (<i>F</i>), <i>wR</i> (<i>F</i> ²)	0.0741, 0.1237	0.1638, 0.2838	0.1335, 0.2031
Difference density: max, min (e Å ⁻³)	1.06, -0.62	2.03, -0.66	0.68, -0.65
Twinned Data Refinement	/	/	0.674(4), 0.326(4)

Table S3

Lethal and teratogenic effects observed in zebrafish (*Danio rerio*) embryos at different hours post fertilization (hpf).

Category	Developmental endpoints	Exposure time (hpf)			
		24	48	72	96/114
Lethal effect	Egg coagulation ^a	●	●	●	●
	No somite formation	●	●	●	●
	Tail not detached	●	●	●	●
	No heartbeat		●	●	●
Teratogenic effect	Malformation of head	●	●	●	●
	Malformation of eyes ^b	●	●	●	●
	Malformation of sacculi/otoliths ^c	●	●	●	●
	Malformation of chorda	●	●	●	●
	Malformation of tail ^d	●	●	●	●
	Scoliosis	●	●	●	●
	Heartbeat frequency		●	●	●
	Blood circulation		●	●	●
	Pericardial edema	●	●	●	●
	Yolk edema	●	●	●	●
	Yolk absorption	●	●	●	●
	Growth retardation ^e	●	●	●	●

^aNo clear organs structure is recognized.

^bMalformation of eyes was recorded for the retardation in eye development and abnormality in shape and size.

^cPresence of no, one or more than two otoliths per sacculus, as well as reduction and enlargement of otoliths and/or sacculi (otic vesicles).

^dTail malformation was recorded when the tail was bent, twisted or shorter than to control embryos as assessed by optical comparison.

^eGrowth retardation was recorded by comparing with the control embryos in development or size (before hatching, at 24 and 48 hpf) or in a body length (after hatching, at and onwards 72 hpf) using by optical comparison using an inverted microscope (CKX41; Olympus, Tokyo, Japan).