

Spatial and Temporal Analysis of Meso- and Microplastic Pollution in the Ligurian and Tyrrhenian Seas

Jessica Caldwell^a, Livius Muff^a, Christopher Kim Pham^b, Alke Petri-Fink^a, Barbara Rothen-Rutishauser^a, Roman Lehner^{a,c}

^aAdolphe Merkle Institute, Université de Fribourg, Chemin des Verdiers 4, 1700 Fribourg, Switzerland

^bInstitute of Marine Research, University of the Azores, Rua da Mãe de Deus 13A, 9700, Ponta Delgada, Região Autónoma dos Açores, Portugal

^cSail and Explore Association, Kirchweg 42, 8755 Ennenda, Switzerland

Supporting Information

Table SI1: A summary of the abundance, mass, and sampling location for each collected sample.

Sample Number	Coordinates at start	Coordinates at end	Mesoplastic Abundance (Items/Km ²)	Microplastic Abundance (Items/Km ²)	Total Abundance (Items/Km ²)	Sample Mass (G/KM ²)
1	N41° 30.807 E9° 23.543	N41° 32.345 E9° 24.538	5,181	4,145	9,326	11.40
2	N41° 41.03 E9° 25.683	N41° 42.602 E9° 25.867	21,336	49,972	71,308	193.15
3	N41° 47.029 E9° 25.98	N41° 48.772 E9° 26.125	8,114	38,540	46,653	286.51
4	N42° 0.919 E9° 34.24	N42° 02.711 E9° 34.952	11,848	58,768	70,616	223.22
5	N42° 22.66 E9° 38.477	N42° 24.191 E9° 38.428	0	6,940	6,940	17.93
6	N42° 41.911	N42° 42.087	3,902	26,829	30,731	73.66

E9° 31.037 E9° 33.487

7	N42° 43.933	N42° 43.934	9,123	19,767	28,890	97.31
	E9° 48.993	E9° 51.364				

8	N42° 43.247	N42° 43.279	26,189	41,154	67,344	213.79
	E10° 8.546	E10° 10.793				

9	N42° 45.057	N42° 47.101	1,488	3,719	5,206	16.73
	E10° 26.478	E10° 28.138				

10	N42° 51.899	N42° 53.01	2,242	9,716	11,958	26.91
	E10° 27.489	E10° 26.93				

11	N42° 54.958	N42° 55.722	6,281	26,382	32,663	54.02
	E10° 12.005	E10° 10.392				

12	N43° 0.8	N43° 0.919	147,120	3,666,898	3,814,018	3,298.40
	E10° 0.378	E9° 58.646				

13	N43° 19.873	N43° 21.325	18,075	127,060	145,136	363.64
	E10° 9.522	E10° 10.64				

14	N43° 39.39	N43° 40.505	9,989	21,643	31,633	86.57
	E10° 10.55	E10° 8.978				

15	N43° 52.613	N43° 53.989	0	1,286	1,286	0.64
	E10° 2.3	E10° 2.23				

16	N43° 59.414	N43° 58.736	22,711	71,682	94,393	287.44
	E9° 46.464	E9° 45.013				

17	N44° 6.355	N44° 6.78	30,922	70,127	101,049	797.90
	E9° 42.985	E9° 40.839				

18	N44° 16.073	N44° 17.049	8,596	81,948	90,544	158.17
	E9° 20.603	E9° 18.935				
19	N44° 17.382	N44° 17.324	86,326	272,842	359,168	1,431.63
	E9° 11.408	E9° 9.453				
20	N44° 19.953	N44° 20.643	19,314	79,128	98,442	244.86
	E9° 4.568	E9° 2.837				




Figure S11: A map of the sample locations with their numeric label. Labels were assigned based upon the order of collection (i.e. sample 1 was the first collected).




Figure S12: Ratios for the particle sizes present within each shape category. The color legend used for all pie charts in the figure is displayed in lower left corner of the whole sample box.




Figure S13: A comparison of a pristine polymer spectra (black) with spectra collected from representative particles from the sample (red). The polyethylene reference is compared to the spectra of a representative pellet (A). The polystyrene reference is compared to the spectra of a representative foam (B). The polyamide reference is compared to the spectra of representative fragment (C). The polypropylene reference spectrum is compared to an additional representative fragment (D). Representative particles are shown as insets in the lower left corner of their spectra.