

Published in "Discrete Applied Mathematics 308 (2022) 111-129"  
which should be cited to refer to this work.  
<https://doi.org/10.1016/j.dam.2019.10.008>

## On some special classes of contact $B_0$ -VPG graphs

*Flavia Bonomo-Braberman, María Pía Mazzoleni, Mariano Leonardo Rean, Bernard Ries*

**ABSTRACT:** A graph  $G$  is a  $B_0$ -VPG graph if one can associate a horizontal or vertical path on a rectangular grid with each vertex such that two vertices are adjacent if and only if the corresponding paths intersect in at least one grid-point. A graph  $G$  is a *contact  $B_0$ -VPG graph* if it is a  $B_0$ -VPG graph admitting a representation with no one-point paths, no two paths crossing, and no two paths sharing an edge of the grid. In this paper, we present a minimal forbidden induced subgraph characterisation of contact  $B_0$ -VPG graphs within four special graph classes: chordal graphs, tree-cographs,  $P_4$ -tidy graphs and  $P_5$ -free graphs. Moreover, we present a polynomial-time algorithm for recognising chordal contact  $B_0$ -VPG graphs.

### Introduction

Columbic et al. introduced in [2] the concept of *vertex intersection graphs of paths in a grid* (referred to as *VPG graphs*). An undirected graph  $G = (V, E)$  is called a VPG graph if one can associate a path in a rectangular grid with each vertex such that two vertices are adjacent if and only if the corresponding paths intersect in at least one grid-point. In the seminal paper on VPG graphs it was shown that this class is equivalent to the earlier defined class of string graphs [14].

Under the perspective of paths in grids, a particular attention was paid to the case where the paths have a limited number of bends. An undirected graph  $G = (V, E)$  is then called a  $B_k$ -VPG graph, for some integer  $k \geq 0$ , if one can associate a path with at most  $k$  bends in a rectangular grid with each vertex such that two vertices are adjacent if and only if the corresponding paths intersect in at least one grid-point. Recognition of VPG graphs is NP-complete by the equivalence with string graphs. Moreover  $B_k$ -VPG recognition is NP-complete for all  $k$  [6].

Since their introduction,  $B_k$ -VPG graphs have been studied by many researchers and the community of people working on these graph classes or related ones is still growing (see for instance [1,2,5,7-9,15,18]).

In this paper, we are interested in a subclass of  $B_k$ -VPG graphs called *contact  $B_k$ -VPG*. A *contact  $B_k$ -VPG representation* of  $G$  is a VPG representation in which each path has length at least one, at most  $k$  bends, and intersecting paths neither cross each other nor share an edge of the grid. A graph is a *contact  $B_k$ -VPG graph* if it has a contact  $B_k$ -VPG representation. Here, we will focus on the special case when  $k = 0$ , i.e. each path is a horizontal or vertical path in the grid.

Contact graphs in general (graphs where vertices represent geometric objects which are allowed to touch but not to cross each other, a natural model arising from real physical objects) have been considered in the past (see for instance [10,11,19,20]). In particular, for intersection models of lines in the plane, it is often the case that three lines intersecting at a same point is not allowed, but we do not impose such a restriction.

As for many graph classes having not many known full characterisations (for example, a complete list of minimal forbidden induced subgraphs is not known), their characterisation within well studied graphs classes or with respect to graph parameters was investigated. In the case of contact  $B_k$ -VPG graphs, it was shown in [12] that every planar bipartite graph is a contact  $B_0$ -VPG graph. Later, in [7], the authors show that every triangle-free planar graph is a contact  $B_1$ -VPG graph. In a recent paper (see [13]), contact  $B_k$ -VPG graphs have been investigated from a structural point of view and it was for instance shown that they do not contain cliques of size 7 and they always contain a vertex of degree at most 6. Moreover, it was shown that they are 6-colourable. Regarding contact  $B_0$ -VPG graphs, it was shown that they are 4-colourable. Furthermore, 3-colouring and the recognition problem were shown to be NP-complete.

In this paper, our goal is to get a better understanding and knowledge of the underlying structure of contact  $B_0$ -VPG graphs. Even though classical graph problems may be difficult to solve in this graph class (see for instance [13]), a better knowledge of its structural properties may lead to good approximation algorithms for these problems. We will consider the following four special graph classes: chordal graphs, tree-cographs,  $P_4$ -tidy graphs and  $P_5$ -free graphs, and we will characterise those graphs from these families that are contact  $B_0$ -VPG. Moreover, we will present a polynomial-time algorithm for recognising chordal contact  $B_0$ -VPG graphs based on our characterisation. For the other graph classes considered here, the characterisation immediately yields a polynomial-time recognition algorithm.

A preliminary version of this paper appears in [4].

## Preliminaries

For concepts and notations not defined here we refer the reader to [3]. All graphs in this paper are simple (i.e., without loops or multiple edges). Let  $G = (V, E)$  be a graph. If  $u, v \in V$  and  $uv \notin E$ ,  $uv$  is called a *nonedge* of  $G$ . We write  $G - v$  for the subgraph obtained by deleting a vertex  $v$  and all the edges incident to  $v$ . Similarly, we write  $G - e$  for the subgraph obtained by deleting an edge  $e$  without deleting its endpoints.

For each vertex  $v$  of  $G$ ,  $N_G(v)$  denotes the *neighbourhood* of  $v$  in  $G$  and  $N_G[v]$  denotes the *closed neighbourhood*, i.e.  $N_G(v) \cup \{v\}$ . For a set  $A \subseteq V$ , we denote by  $N(A)$  the set of vertices having a neighbour in  $A$ , and by  $N[A]$  the set of vertices belonging to  $A$  or having a neighbour in  $A$ . Two vertices  $v$  and  $w$  of  $G$  are *false twins* (resp. *true twins*) if  $N_G(v) = N_G(w)$  (resp.  $N_G[v] = N_G[w]$ ).

Given a subset  $A \subseteq V$ ,  $G[A]$  stands for *the subgraph of  $G$  induced by  $A$* , and  $G \setminus A$  denotes the *induced subgraph  $G[V \setminus A]$* . We say that a vertex  $v \in V \setminus A$  is *complete to  $A$*  if  $v$  is adjacent to every vertex of  $A$ , and that  $v$  is *anticomplete to  $A$*  if  $v$  has no neighbour in  $A$ . Similarly, we say that two disjoint sets  $A, B \subset V$  are *complete* (resp. *anticomplete*) to each other if every vertex in  $A$  is complete (resp. anticomplete) to  $B$ .

A *clique* is a set of pairwise adjacent vertices. A vertex  $v$  is *simplicial*, if  $N_G(v)$  is a clique. A *stable set* is a set of vertices no two of which are adjacent. A *complete graph* is a graph such that all its vertices are adjacent to each other, i.e. a graph induced by a clique. The *complete graph on  $n$  vertices* is denoted by  $K_n$ . In particular,  $K_3$  is called a *triangle*.  $K_4^-$  stands for the graph obtained from  $K_4$  by deleting exactly one edge.

The *complement graph* of  $G = (V, E)$  is the graph  $\bar{G} = (V, \bar{E})$  such that  $\bar{E} = \{uv \mid uv \notin E\}$ . Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs. The *disjoint union of  $G_1$  and  $G_2$* , denoted by  $G_1 \cup G_2$ , is the graph whose vertex set is  $V_1 \cup V_2$  and whose edge set is  $E_1 \cup E_2$ . The *join of  $G_1$  and  $G_2$* , denoted by  $G_1 \vee G_2$ , is the graph obtained by first taking the disjoint union of  $G_1$  and  $G_2$  and then making  $V_1$  and  $V_2$  complete to each other. Notice that  $\overline{G_1 \cup G_2} = \bar{G}_1 \vee \bar{G}_2$ .

Given a graph  $H$ , we say that  $G$  *contains no induced  $H$* , if  $G$  contains no induced subgraph isomorphic to  $H$ . If  $\mathcal{H}$  is a family of graphs,  $G$  is said to be  *$\mathcal{H}$ -free* if  $G$  contains no induced subgraph isomorphic to some graph belonging to  $\mathcal{H}$ .

Let  $\mathcal{G}$  be a class of graphs. A graph belonging to  $\mathcal{G}$  is called a  *$\mathcal{G}$ -graph*. If  $G \in \mathcal{G}$  implies that every induced subgraph of  $G$  is a  $\mathcal{G}$ -graph,  $\mathcal{G}$  is said to be *hereditary*. If  $\mathcal{G}$  is a hereditary class, a graph  $H$  is a *minimal forbidden induced subgraph of  $\mathcal{G}$* , or more briefly, *minimally non- $\mathcal{G}$* , if  $H$  does not belong to  $\mathcal{G}$  but every proper induced subgraph of  $H$  is a  $\mathcal{G}$ -graph.

A *path* is a sequence of vertices  $v_1, \dots, v_k$  such that  $v_i$  is adjacent to  $v_{i+1}$ , for  $i = 1, \dots, k-1$ . The vertices  $v_2, \dots, v_{k-1}$  are called *internal vertices* of the path. If there is no edge  $v_i v_j$  such that  $|i-j| \geq 2$ , the path is said to be *chordless* or *induced*. A *cycle  $C$*  is a sequence of vertices  $v_1, \dots, v_k$  such that  $v_i$  is adjacent to  $v_{i+1}$  for  $i = 1, \dots, k$ , where indices are taken modulo  $k$ . If there is no edge  $v_i v_j$  such that  $|i-j| \geq 2$ ,  $C$  is said to be *chordless* or *induced*. The induced path (resp. induced cycle) on  $n$  vertices is denoted  $P_n$  (resp.  $C_n$ ). A graph is called *chordal* if it does not contain any chordless cycle of length at least four. A *block graph* is a chordal graph which is  $K_4^-$ -free.

A graph is *bipartite*, if its vertex set can be partitioned into two stable sets. If, in addition, the two stable sets are complete to each other, the graph is called *complete bipartite*.  $K_{n,m}$  stands for the complete bipartite graph whose vertex set can be partitioned into two stable sets  $V_1, V_2$  such that  $|V_1| = n$  and  $|V_2| = m$ .

A graph  $G$  is *connected*, if for each pair of vertices  $u, v$  there exists a path from  $u$  to  $v$ . A *tree* is a connected graph with no induced cycle. Given a connected graph  $G = (V, E)$ , the *distance between two vertices  $u, v \in V$* , denoted by  $d_G(u, v)$ , is the number of edges of a shortest path from  $u$  to  $v$ . The *diameter* of  $G$  is the maximum distance between two vertices.

An undirected graph  $G = (V, E)$  is called a  *$B_k$ -VPG graph*, for some integer  $k \geq 0$ , if one can associate a path with at most  $k$  *bends* (a bend is a 90 degrees turn of a path at a grid-point) on a rectangular grid with each vertex such that




Fig. 1. A graph  $G$  and a contact  $B_0$ -VPG representation of it.




Fig. 2. The graph  $H_0$ .

two vertices are adjacent if and only if the corresponding paths intersect in at least one grid-point. Such a representation is called a  $B_k$ -VPG representation. The horizontal grid lines will be referred to as *rows* and denoted by  $x_0, x_1, \dots$  and the vertical grid lines will be referred to as *columns* and denoted by  $y_0, y_1, \dots$ . We are interested in a subclass of  $B_0$ -VPG graphs called contact  $B_0$ -VPG. A contact  $B_0$ -VPG representation  $\mathcal{R}(G)$  of  $G$  is a  $B_0$ -VPG representation in which each path in the representation is either a horizontal path or a vertical path on the grid, with length at least one (the length is the number of grid-points minus one), such that two vertices are adjacent if and only if the corresponding paths intersect in at least one grid-point without crossing each other and without sharing an edge of the grid. A graph is a contact  $B_0$ -VPG graph if it has a contact  $B_0$ -VPG representation. For every vertex  $v$ , we denote by  $P_v$  the corresponding path in  $\mathcal{R}(G)$  (see Fig. 1). Consider a clique  $K$  in  $G$ . A path  $P_v$  representing a vertex  $v \in K$  is called a *path of the clique*  $K$ .

Let us start with an easy but very helpful lemma.

**Lemma 1.** *Let  $G$  be a contact  $B_0$ -VPG graph. Then the size of a biggest clique in  $G$  is at most 4, i.e.  $G$  is  $K_5$ -free.*

**Proof.** Given two adjacent vertices in  $G$ , the intersection of their paths in any contact  $B_0$ -VPG representation is exactly one grid point. Moreover, it is easy to see that all paths corresponding to vertices in a clique of  $G$  must intersect in the same grid point. Assume there is a clique  $K$  of size 5 in  $G$  and let  $P$  be the point of intersection of the corresponding paths in the grid. At least two of the paths must be in the same row or the same column, and contain at least one grid edge intersecting  $P$  (a path cannot be only a grid point), a contradiction.  $\square$

**Remark 2.** Let  $G$  be a  $K_4^-$ -free graph containing an induced cycle  $C$  of at least 4 vertices. Then no vertex is adjacent to 3 consecutive vertices of  $C$ .

Let  $G$  be a contact  $B_0$ -VPG graph, and  $K$  be a clique in  $G$ . A vertex  $v$  is called an *end* in a contact  $B_0$ -VPG representation of  $K$  if the grid point representing the intersection of the paths of  $K$  corresponds to an endpoint of  $P_v$ .

**Remark 3.** Let  $G$  be a contact  $B_0$ -VPG graph, and  $K$  be a clique in  $G$  of size four. Then, every vertex in  $K$  is an end in any contact  $B_0$ -VPG representation of  $K$ .

**Lemma 4.** *In any contact  $B_0$ -VPG representation of  $C_4$ , the union of the paths representing vertices in  $C$  must enclose a rectangle of the grid.*

**Proof.** Consider a  $B_0$ -VPG representation of  $C_4$ . At least two vertices, say  $a$  and  $b$ , in  $C$  have the same direction. We can assume that  $P_a$  and  $P_b$  are both vertical. If  $a$  and  $b$  are adjacent, then the corresponding paths intersect in a row  $x_i$  of the grid. One of them, say  $P_a$ , is above  $x_i$  and the other is below  $x_i$ . Let  $c$  be the vertex adjacent to  $a$  and non adjacent to  $b$ . Clearly, the path  $P_c$  representing  $c$  must be also above  $x_i$ . Similarly, the path representing the vertex  $d$  adjacent to  $b$  and non adjacent to  $a$  must be below  $x_i$ . But then it is impossible for  $P_c$  and  $P_d$  to intersect. Therefore,  $a$  and  $b$  are non adjacent. Now, it is clear that  $P_c$  and  $P_d$  must be both horizontal, otherwise we could repeat the previous argument. If  $P_a$  and  $P_b$  lie in columns  $y_i$  and  $y_j$ , then  $P_c$  and  $P_d$  must contain all points of the grid between  $y_i$  and  $y_j$  in their respective columns, say  $x_k$  and  $x_l$ . Then, these paths enclose the rectangle limited by rows  $y_i, y_j$  and columns  $x_k, x_l$ .  $\square$

In what follows, we give a set of graphs that are not contact  $B_0$ -VPG graphs. We will use this result later to obtain our characterisations. Let  $H_0$  denote the graph composed of three  $K_4$ 's that share a common vertex and such that there are no further edges (see Fig. 2).

**Lemma 5.** *If  $G$  is a contact  $B_0$ -VPG graph, then  $G$  is  $\{K_5, K_{3,3}, H_0, K_4^-\}$ -free.*

**Proof.** Let  $G$  be a contact  $B_0$ -VPG graph. It immediately follows from Lemma 1 that  $G$  is  $K_5$ -free.

Now consider the graph  $K_{3,3}$ . Let  $C$  be a cycle of length four in  $K_{3,3}$  induced by the vertices  $a, b, c, d$ . If  $K_{3,3}$  is contact  $B_0$ -VPG, then, by Lemma 4, in any contact  $B_0$ -VPG representation of  $C$ , the union of the paths representing vertices in  $C$  must enclose a rectangle of the grid. Assume that  $P_a, P_c$  are horizontal paths, and  $P_b, P_d$  are vertical paths. Now, consider vertices  $e$  and  $f$  in  $K_{3,3}$  with  $e$  being adjacent to  $a$  and  $c$ , and  $f$  being adjacent to  $b$  and  $d$ . Each of the paths  $P_e, P_f$  must intersect opposite paths of the rectangle. Clearly,  $P_e$  must be a vertical path and  $P_f$  must be a horizontal. If  $P_e$  is contained inside the rectangle, then it is impossible for  $P_f$  to intersect  $P_b, P_d$  while being inside the rectangle without crossing  $P_e$ . So  $P_f$  must be outside the rectangle, but then it cannot intersect  $P_e$ . If  $P_e$  lies outside the rectangle, then of course  $P_f$  has to lie outside the rectangle as well, otherwise it cannot intersect  $P_e$ . But now it cannot intersect both  $P_b, P_d$  without crossing at least one of them. So we conclude that  $K_{3,3}$  is not  $B_0$ -VPG.

Now let  $v, w$  be two adjacent vertices in  $G$ . Then, in any contact  $B_0$ -VPG representation of  $G$ ,  $P_v$  and  $P_w$  intersect at a grid-point  $P$ . Clearly, every common neighbour of  $v$  and  $w$  must also contain  $P$ . Hence,  $v$  and  $w$  cannot have two common neighbours that are non-adjacent. So,  $G$  is  $K_4^-$ -free.

Finally, consider the graph  $H_0$  which consists of three cliques of size four, say  $A, B$  and  $C$ , with a common vertex  $x$ . Suppose that  $H_0$  is contact  $B_0$ -VPG. Then, it follows from Remark 3 that every vertex in  $H_0$  is an end in any contact  $B_0$ -VPG representation of  $H_0$ . In particular, vertex  $x$  is an end in any contact  $B_0$ -VPG representation of  $A, B$  and  $C$ . In other words, the grid-point representing the intersection of the paths of each of these three cliques corresponds to an endpoint of  $P_x$ . Since these cliques have only vertex  $x$  in common, these grid-points are all distinct. But this is a contradiction, since  $P_x$  has only two endpoints. So we conclude that  $H_0$  is not contact  $B_0$ -VPG, and hence the result follows.  $\square$

## Chordal graphs

In this section, we will consider chordal graphs and characterise those that are contact  $B_0$ -VPG. First, let us point out the following corollary.

**Corollary 6.** *A chordal contact  $B_0$ -VPG graph is a block graph.*

This follows directly from Lemma 5 and the definition of block graphs.

The following lemma states an important property of minimal chordal non contact  $B_0$ -VPG graphs that contain neither  $K_5$  nor  $K_4^-$ .

**Lemma 7.** *Let  $G$  be a  $\{K_5, K_4^-\}$ -free graph. If  $G$  is a minimal non contact  $B_0$ -VPG graph, then every simplicial vertex of  $G$  has degree exactly three.*

**Proof.** Since  $G$  is  $K_5$ -free, every clique in  $G$  has size at most four. Therefore, every simplicial vertex has degree at most three. Let  $v$  be a simplicial vertex of  $G$ . Assume first that  $v$  has degree one and consider a contact  $B_0$ -VPG representation of  $G - v$  (which exists since  $G$  is minimal non contact  $B_0$ -VPG). Let  $w$  be the unique neighbour of  $v$  in  $G$ . Without loss of generality, we may assume that the path  $P_w$  lies on some row of the grid. Now clearly, we can add one extra column to the grid between any two consecutive vertices of the grid belonging to  $P_w$  and adapt all paths without changing the intersections (if the new column is added between column  $y_i$  and  $y_{i+1}$ , we extend all paths containing a grid-edge with endpoints in column  $y_i$  and  $y_{i+1}$  in such a way that they contain the new edges in the same row and between column  $y_i$  and  $y_{i+2}$  of the new grid, and any other path remains the same). But then we may add a path representing  $v$  on this column which only intersects  $P_w$  (adding a row to the grid and adapting the paths again, if necessary) and thus, we obtain a contact  $B_0$ -VPG representation of  $G$ , a contradiction. So suppose now that  $v$  has degree two, and again consider a contact  $B_0$ -VPG representation of  $G - v$ . Let  $w_1, w_2$  be the two neighbours of  $v$  in  $G$ . Then,  $w_1, w_2$  do not have any other common neighbour since  $G$  is  $K_4^-$ -free. Let  $P$  be the grid-point corresponding to the intersection of the paths  $P_{w_1}$  and  $P_{w_2}$ . Since these paths do not cross and since  $w_1, w_2$  do not have any other common neighbour (except  $v$ ), there is at least one grid-edge having  $P$  as one of its endpoints and which is not used by any path of the representation. But then we may add a path representing  $v$  by using only this particular grid-edge (or adding a row/column to the grid that subdivides this edge and adapting the paths, if the other endpoint of the grid-edge belongs to a path in the representation). Thus, we obtain a contact  $B_0$ -VPG representation of  $G$ , a contradiction. We conclude therefore that  $v$  has degree exactly three.  $\square$

Let  $v$  be a vertex of a contact  $B_0$ -VPG graph  $G$ . An endpoint of its corresponding path  $P_v$  is *free* in a representation of  $G$ , if  $P_v$  does not intersect any other path at that endpoint;  $v$  is called *internal* if no representation of  $G$  with a free endpoint of  $P_v$  exists. If in a representation of  $G$  a path  $P_v$  intersects a path  $P_w$  but not at an endpoint of  $P_w$ ,  $v$  is called a *middle neighbour* of  $w$ .

In the following two lemmas we associate the fact of being or not an internal vertex of  $G$  with the contact  $B_0$ -VPG representation of  $G$ .




Fig. 3. How to split  $P_u$  into two paths.

**Lemma 8.** *Let  $G$  be a chordal contact  $B_0$ -VPG graph and let  $v$  be a non internal vertex in  $G$ . Then, there exists a contact  $B_0$ -VPG representation of  $G$  in which all the paths representing vertices in  $G - v$  lie to the left of a free endpoint of  $P_v$  (by considering  $P_v$  as a horizontal path).*

**Proof.** We will do a proof by induction on the number of vertices of  $G$ . If there is only one vertex in  $G$  the result is trivial. Suppose  $G$  is a graph with at least two vertices. Consider a contact  $B_0$ -VPG representation of  $G$ . Without loss of generality, we may assume that  $P_v$  lies on a row  $x_i$  between columns  $y_j, y_k, j < k$ , and its right endpoint is free. Such a representation exists, since  $v$  is not internal.

If  $v$  is a middle neighbour of another vertex, say  $u$ , we do the following. Assume  $P_u$  lies on column  $y_j$  between rows  $x_\ell$  and  $x_t, \ell < t$ . We split  $P_u$  into two paths,  $P_{u_1}, P_{u_2}$ , such that  $P_{u_1}$  goes from row  $x_i$  to row  $x_t$  and  $P_{u_2}$  goes from row  $x_\ell$  to row  $x_i$  (see Fig. 3). We denote the corresponding graph by  $G^*$ . If  $v$  is not a middle vertex of another vertex, then we simply set  $G^* = G$ .

**Claim.** *The graph  $G^*$  is chordal.*

In the second case, it is trivial. In the first case, suppose  $G^*$  contains a chordless cycle  $C$  of length at least 4. Since  $G$  is chordal,  $C$  contains at least one of  $u_1, u_2$ . Suppose first it contains both  $u_1$  and  $u_2$ . As they are adjacent in  $G^*$ , and contracting them into the vertex  $u$  yields an induced subgraph of  $G$ , it follows that  $C$  has length 4. As in the proof of Lemma 4, it can be seen that the paths corresponding to two consecutive vertices in a  $C_4$  cannot be both vertical. So, suppose that  $C$  contains only one of  $u_1, u_2$ , say  $u_1$ . Since  $G$  is chordal,  $u_2$  has to be adjacent to every vertex of  $C \setminus N_{G^*}[u_1]$ . Since  $u_1$  and  $u_2$  cannot have two non-adjacent common neighbours, at least one of the neighbours of  $u_1$  in  $C$  is not adjacent to  $u_2$ . Thus, its corresponding path either lies on column  $y_j$  having its lower endpoint in row  $x_t$  or lies on some row between  $x_{i+1}$  and  $x_t$ . In either case, this vertex cannot have a common neighbour with  $u_2$ , a contradiction.  $\diamond$

Now, for every vertex  $w$  in  $N_{G^*}(v)$ , consider the connected component  $C_w$  of  $G^* - (N_{G^*}[v] - w)$  containing  $w$ . Notice that  $C_w$  is also chordal contact  $B_0$ -VPG and  $w$  is non internal in  $G^* - (N_{G^*}[v] - w)$ . Furthermore, if there are two distinct vertices  $w$  and  $w'$  in  $N_{G^*}(v)$ , then  $C_w$  and  $C_{w'}$  are disjoint. By contradiction, suppose that a vertex  $x$  is in the intersection of  $C_w$  and  $C_{w'}$ . Then, there is a path  $\alpha_1$  between  $w$  and  $x$ , and a path  $\alpha_2$  between  $x$  and  $w'$ . First, suppose  $w$  and  $w'$  are non adjacent. Joining both paths we can extract a new induced path  $\alpha_3$  between  $w$  and  $w'$  which necessarily has length  $\geq 3$ . But then, adding  $v$  to  $\alpha_3$  forms an induced cycle with length  $\geq 4$ , a contradiction. On the other hand, if  $w$  and  $w'$  are adjacent, first remove the edge  $w$  and  $w'$ . Joining the paths  $\alpha_1$  and  $\alpha_2$  we can extract an induced path  $\alpha_3$  between  $w$  and  $w'$ , which necessarily has length  $\geq 4$ , since  $G$  is  $K_4^-$ -free (see Lemma 5) and, therefore, any vertex adjacent to both  $w$  and  $w'$  must be also adjacent to  $v$ , implying that it does not belong to  $C_w$ . Adding the edge between  $w$  and  $w'$  again, we obtain an induced cycle with length  $\geq 4$ , a contradiction.

Finally, considering the case in which  $G^* = G$ , it is clear that  $C_w$  has at least one vertex less than  $G$ , namely  $v$ ; otherwise, if  $u$  was split, the size of  $G^*$  is one more than the size of  $G$ , but then at least two vertices are removed in  $C_w$ , namely  $v$  and one between  $u_1$  and  $u_2$  (since there is only one vertex in  $N_{G^*}[v]$  that we are not removing).

Then, by induction, there exists a contact  $B_0$ -VPG representation of  $C_w$ , for each such  $w$ , with all the paths lying to the left of one free endpoint of  $P_w$ . Now, we replace the initial representation of  $C_w$  by the new one (the one where all the paths lie to the left of one free endpoint of  $P_w$ ) by rotating it such that  $P_w$  has its free endpoint on the grid-point corresponding to the intersection of  $P_w$  and  $P_v$ , and belongs to the same side as in the old representation. Notice that we may need to extend the path  $P_v$  to the right before doing the replacement of these new representations to assure that they do not overlap. Therefore, by extending if necessary the path  $P_v$  a little more to the right, we obtain a contact  $B_0$ -VPG representation of  $G^*$  in which all the paths lie to the left of one free endpoint of  $P_v$  (see Fig. 4). In case  $P_u$  was split into  $P_{u_1}$  and  $P_{u_2}$ , we now glue these two paths together again.  $\square$

**Lemma 9.** *Let  $G$  be a chordal contact  $B_0$ -VPG graph. A vertex  $v$  in  $G$  is internal if and only if in every contact  $B_0$ -VPG representation of  $G$ , each endpoint of the path  $P_v$  either corresponds to the intersection of a representation of  $K_4$  or intersects a path  $P_w$ , which represents an internal vertex  $w$ , but not at an endpoint of  $P_w$ .*




Fig. 4. Figure illustrating Lemma 8.




Fig. 5. Figure illustrating Lemma 9.

**Proof.** The if part is trivial. Assume now that  $v$  is an internal vertex of  $G$  and consider an arbitrary contact  $B_0$ -VPG representation of  $G$ . Let  $P$  be an endpoint of the path  $P_v$  and  $K$  the maximal clique corresponding to all the paths containing the point  $P$ . Notice that clearly  $v$  is an end in  $K$  by definition of  $K$ . First, suppose there is a vertex  $w$  in  $K$  which is not an end. Then, it follows from Remark 3 that the size of  $K$  is at most three. Without loss of generality, we may assume that  $P_v$  lies on some row and  $P_w$  on some column. If  $w$  is an internal vertex, we are done. So we may assume now that  $w$  is not an internal vertex in  $G$ . Consider  $G \setminus (K \setminus \{w\})$ , and let  $C_w$  be the connected component of  $G \setminus (K \setminus \{w\})$  containing  $w$ . Notice that  $w$  is not an internal vertex in  $C_w$  either. By Lemma 8, there exists a contact  $B_0$ -VPG representation of  $C_w$  with all the paths lying to the left of a free endpoint of  $P_w$ . Now, replace the old representation of  $C_w$  by the new one such that  $P$  corresponds to the free endpoint of  $P_w$  in the representation of  $C_w$  (it might be necessary to refine – by adding rows and/or columns – the grid to ensure that there are no unwanted intersections) and  $P_w$  uses the same column as before. Finally, if  $K$  had size three, say it contains some vertex  $u$  in addition to  $v$  and  $w$ , then we proceed as follows. Similar to the above, there exists a contact  $B_0$ -VPG representation of  $C_u$ , the connected component of  $G \setminus (K \setminus \{u\})$  containing  $u$ , with all the paths lying to the left of a free endpoint of  $P_u$ , since  $u$  is clearly not internal in  $C_u$ . We then replace the old representation of  $C_u$  by the new one such that the endpoint of  $P_u$  that intersected  $P_w$  previously corresponds to the grid-point  $P$  and  $P_u$  lies on the same column as  $P_w$  (again, we may have to refine the grid). This clearly gives us a contact  $B_0$ -VPG representation of  $G$ . But now we may extend  $P_v$  such that it strictly contains the grid-point  $P$  and thus,  $P_v$  has a free endpoint, a contradiction (see Fig. 5). So  $w$  must be an internal vertex.


Now, assume that all vertices in  $K$  are ends. If  $|K| = 4$ , we are done. So we may assume that  $|K| \leq 3$ . Hence, there is at least one grid-edge containing  $P$ , which is not used by any paths of the representation. Without loss of generality, we may assume that this grid-edge belongs to some row  $x_i$ . If  $P_v$  is horizontal, we may extend it such that it strictly contains  $P$ . But then  $v$  is not internal anymore, a contradiction. If  $P_v$  is vertical, then we may extend  $P_w$ , where  $w \in K$  is such that  $P_w$  is a horizontal path. But now we are again in the first case discussed above.  $\square$

In other words, Lemma 9 tells us that a vertex  $v$  is an internal vertex in a chordal contact  $B_0$ -VPG graph if and only if we are in one of the following situations:

- $v$  is the intersection of two cliques of size four (we say that  $v$  is of type 1);
- $v$  belongs to exactly one clique of size four and in every contact  $B_0$ -VPG representation,  $v$  is a middle neighbour of some internal vertex (we say that  $v$  is of type 2);
- $v$  does not belong to any clique of size four and in every contact  $B_0$ -VPG representation,  $v$  is a middle neighbour of two internal vertices (we say that  $v$  is of type 3).

Notice that two internal vertices of type 1 cannot be adjacent (except when they belong to a same  $K_4$ ). Furthermore, an internal vertex of type 1 cannot be the middle-neighbour of some other vertex.

Let  $\mathcal{T}$  be the family of graphs containing  $H_0$  (see Fig. 2) as well as all graphs that can be partitioned into a nontrivial tree  $T_0$  of maximum degree at most three and the disjoint union of triangles, in such a way that each triangle is complete



**Fig. 6.** An example of a graph in  $\mathcal{T}$ .

to a vertex  $v$  of  $T_0$  and anticomplete to  $T_0 - \{v\}$ , every leaf  $v$  of  $T_0$  is complete to exactly two triangles, every vertex of degree two in  $T_0$  is complete to exactly one triangle, and vertices of degree three in  $T_0$  have no neighbours outside  $T_0$  (see Fig. 6).

Notice that all graphs in  $\mathcal{T}$  are chordal. We denote by  $B(T)$  the base tree of  $T$  in  $\mathcal{T}$ .

**Lemma 10.** *The graphs in  $\mathcal{T}$  are not contact  $B_0$ -VPG.*

**Proof.** By Lemma 5, the graph  $H_0$  is not contact  $B_0$ -VPG. Consider now a graph  $T \in \mathcal{T}$ ,  $T \neq H_0$ . Suppose that  $T$  is contact  $B_0$ -VPG. Consider an arbitrary contact  $B_0$ -VPG representation of  $T$ . Consider the base tree  $B(T)$  and direct an edge  $uv$  of it from  $u$  to  $v$  if the path  $P_v$  contains an endpoint of the path  $P_u$  (this way some edges might be directed both ways). If a vertex  $v$  has degree  $d_B(v)$  in  $B(T)$ , then by definition of the family  $\mathcal{T}$ ,  $v$  belongs to  $3 - d_B(v)$   $K_4$ 's in  $T$ . Notice that  $P_v$  spends one endpoint in each of these  $K_4$ 's. Thus, any vertex  $v$  in  $B(T)$  has at most  $2 - (3 - d_B(v)) = d_B(v) - 1$  outgoing edges. This implies that the sum of out-degrees in  $B(T)$  is at most  $\sum_{v \in B(T)} (d_B(v) - 1) = n - 2$ , where  $n$  is the number of vertices in  $B(T)$ . But this is clearly impossible since there are  $n - 1$  edges in  $B(T)$  and all edges are directed.  $\square$

We will show now how to construct new graphs in  $\mathcal{T}$  from others.

**Lemma 11.**

- (i) Given  $T \in \mathcal{T}$  and  $v \in B(T)$  such that  $v$  belongs to at least one  $K_4$ , say  $K$ , then the graph  $T'$  constructed by removing the other vertices in  $K$  (different from  $v$ ) and adding one vertex  $w$  to  $B(T)$ , belonging to two copies of  $K_4$  (sharing vertex  $w$ ), and adjacent to  $v$ , belongs to  $\mathcal{T}$ .
- (ii) Given  $T_1, T_2 \in \mathcal{T}$ ,  $v_1 \in B(T_1)$  and  $v_2 \in B(T_2)$  such that  $v_1$  and  $v_2$  belong to at least one  $K_4$  each, say  $K_1$  and  $K_2$ , then the graph  $T'$  constructed by removing the other vertices in  $K_1$  and  $K_2$  (different from  $v_1$  and  $v_2$ ) and adding one vertex  $w$  to  $B(T_1) \cup B(T_2)$ , belonging to a  $K_4$ , and adjacent to both  $v_1$  and  $v_2$ , belongs to  $\mathcal{T}$ .

**Proof.**

- (i) In this case we have  $B(T') = B(T) \cup \{w\}$ . It is clear that every vertex in  $B(T')$  has degree 3 or less, since we only changed the degree of  $v$ , which is one less, and the degree of  $w$  is one (only adjacent to  $v$  in  $B(T')$ ). Moreover,  $w$  is a leaf in  $B(T')$  and, by construction, it belongs to two copies of  $K_4$  (sharing vertex  $w$ ). Finally, notice that  $v$  has degree 1 or 2 in  $B(T)$  since vertices of degree 3 in  $B(T)$  does not belong to any  $K_4$ . If  $v$  is a leaf in  $B(T)$ , then  $v$  is a degree 2 vertex in  $B(T')$  and, since we removed the other vertices in  $K$ , it belongs to only one  $K_4$  in  $T'$ . Otherwise,  $v$  has degree 2 in  $B(T)$  and therefore, it has degree 3 in  $B(T')$  and does not belong to any  $K_4$  in  $T'$ . Thus,  $T' \in \mathcal{T}$ .
- (ii) In this case we have  $B(T') = B(T_1) \cup B(T_2) \cup \{w\}$ . The proof follows in the same manner as the previous item.  $\square$

For the next lemma we need to consider an orientation of some edges related to a contact  $B_0$ -VPG representation of  $G$ , given by the following rule. If  $v, w \in G$  and  $v$  is a middle neighbour of  $w$ , then we give the orientation from  $v$  to  $w$ . Let  $C_v$  be the reachable vertices starting from  $v$ , including  $v$ . Notice that if  $v$  is internal,  $C_v = \{v\}$  if and only if  $v$  is of type 1. Also notice that  $C_v$  is independent of the representation for internal vertices. As a consequence of the previous lemma, we can prove the following.

**Lemma 12.** *Let  $G$  be a chordal contact  $B_0$ -VPG graph. If a vertex  $v$  in  $G$  is internal, the graph  $G'$  constructed by adding a  $K_4$ , say  $K$ , containing  $v$  to  $G$  contains an induced subgraph  $T \in \mathcal{T}$ . Moreover,  $B(T) = C_v$ .*

**Proof.** We will prove this by induction in the number of vertices in  $C_v$ . By Lemma 9,  $v$  must be of type 1, 2 or 3. As noted before, the base case is when  $v$  is of type 1. But then  $v$  is the intersection of three cliques of size 4 in  $G'$ , namely  $K$  and the two cliques in which  $v$  is an end; and thus,  $G'$  contains  $T = H_0$ . Therefore  $B(T) = \{v\} = C_v$ .

Now, if  $v$  is of type 2,  $v$  is a middle neighbour of exactly one other internal vertex  $w$ . Therefore  $C_v = C_w \cup \{v\}$ . Define  $G_w$  as the induced subgraph of the connected component of  $G - v$  containing  $w$ . Notice that  $w$  is still internal in  $G_w$  since  $v$  is a middle neighbour of  $w$  in  $G$ . Then, adding a  $K' = K_4$  containing  $w$  to  $G_w$  we obtain a  $T_1 \in \mathcal{T}$  induced in  $G_w$  (and, therefore, also induced in  $G$ ) with  $B(T_1) = C_w$ , by inductive hypothesis applied to  $w$  in  $G_w$ . By Lemma 11 (i), we can construct  $T \in \mathcal{T}$  by removing the other vertices in  $K'$  (different from  $w$ ) and adding the vertex  $v$  (in  $G$ ) to  $B(T_1)$ , which belongs to two copies of  $K_4$  (one is  $K$  and the other is the one in which  $v$  is an end), and is adjacent to  $w$ . Then,  $T$  is an induced subgraph of  $G$  and we have  $B(T) = C_w \cup \{v\} = C_v$ . Finally, if  $v$  is of type 3,  $v$  is a middle neighbour of exactly two other internal vertices  $w_1$  and  $w_2$ . The proof continues in the same manner as before, applying inductive hypothesis to the corresponding  $G_{w_1}$  and  $G_{w_2}$  and then using the second item of Lemma 11.  $\square$

Using Lemmas 7–12, we are able to prove the following theorem, which provides a minimal forbidden induced subgraph characterisation of chordal contact  $B_0$ -VPG graphs.

**Theorem 13.** *Let  $G$  be a chordal graph. Let  $\mathcal{F} = \mathcal{T} \cup \{K_5, K_4^-\}$ . Then,  $G$  is a contact  $B_0$ -VPG graph if and only if  $G$  is  $\mathcal{F}$ -free.*

**Proof.** Suppose that  $G$  is a chordal contact  $B_0$ -VPG graph. It follows from Lemmas 5 and 10 that  $G$  is  $\mathcal{T}$ -free and contains neither a  $K_4^-$  nor a  $K_5$ .

Conversely, suppose now that  $G$  is chordal and  $\mathcal{F}$ -free. By contradiction, suppose that  $G$  is not contact  $B_0$ -VPG and assume furthermore that  $G$  is a minimal non contact  $B_0$ -VPG graph. Let  $v$  be a simplicial vertex of  $G$  ( $v$  exists since  $G$  is chordal). By Lemma 7, it follows that  $v$  has degree three. Consider a contact  $B_0$ -VPG representation of  $G - v$  and let  $K = \{v_1, v_2, v_3\}$  be the set of neighbours of  $v$  in  $G$ . Since  $G$  is  $K_4^-$ -free, it follows that any two neighbours of  $v$  cannot have a common neighbour which is not in  $K$ . First suppose that all the vertices in  $K$  are ends in the representation of  $G - v$ . Thus, there exists a grid-edge not used by any path and which has one endpoint corresponding to the intersection of the paths  $P_{v_1}, P_{v_2}, P_{v_3}$ . But now we may add the path  $P_v$  using exactly this grid-edge (we may have to add a row/column to the grid that subdivides this grid-edge and adapt the paths, if the other endpoint of the grid-edge belongs to a path in the representation). Hence, we obtain a contact  $B_0$ -VPG representation of  $G$ , a contradiction.

Thus, we may assume now that there exists a vertex in  $K$  which is not an end, say  $v_1$ . Notice that  $v_1$  must be an internal vertex. If not, there is a contact  $B_0$ -VPG representation of  $G - v$  in which  $v_1$  has a free end. Then, using similar arguments as in the proof of Lemma 9, we may obtain a representation of  $G - v$  in which all vertices of  $K$  are ends. As described previously, we can add  $P_v$  to obtain a contact  $B_0$ -VPG representation of  $G$ , a contradiction. Now, consider the graph  $G - K$ . This graph is clearly chordal contact  $B_0$ -VPG as being an induced subgraph of  $G - v$ . Then, by Lemma 12, adding the clique  $K \cup \{v\}$  (containing the internal vertex  $v$ ) to  $G - K$  (which gives the graph  $G$ ) contains an induced subgraph  $T \in \mathcal{T}$ , a contradiction.  $\square$

Interval graphs form a subclass of chordal graphs. They are defined as being chordal graphs not containing any asteroidal triple, i.e. not containing three pairwise non-adjacent vertices such that there exists a path between any two of them avoiding the neighbourhood of the third one. Clearly, any graph in  $\mathcal{T}$  for which the base tree has maximum degree three contains an asteroidal triple. On the other hand,  $H_0$  and every graph in  $\mathcal{T}$  obtained from a base tree of maximum degree at most two are clearly interval graphs. Denote by  $\mathcal{T}'$  the family consisting of  $H_0$  and the graphs of  $\mathcal{T}$  whose base tree has maximum degree at most two. We obtain the following corollary which provides a minimal forbidden induced subgraph characterisation of contact  $B_0$ -VPG graphs restricted to interval graphs.

**Corollary 14.** *Let  $G$  be an interval graph and  $\mathcal{F}' = \mathcal{T}' \cup \{K_5, K_4^-\}$ . Then,  $G$  is a contact  $B_0$ -VPG graph if and only if  $G$  is  $\mathcal{F}'$ -free.*


### Recognition algorithm

In this section, we will provide a polynomial-time recognition algorithm for chordal contact  $B_0$ -VPG graphs which is based on the characterisation given in Section 3. This algorithm takes a chordal graph as input and returns YES if the graph is contact  $B_0$ -VPG and, if not, it returns NO as well as a forbidden induced subgraph. The main loop (step 7) will try to find a graph  $T \in \mathcal{T}$ ,  $T \neq H_0$ . For this purpose, some vertices will be marked and some edges will be directed and coloured. At the beginning all vertices are unmarked and all edges are undirected and uncoloured. We will first give the pseudo-code of our algorithm and then explain the different steps.

**Input:** a chordal graph  $G = (V, E)$ ;

**Output:** YES, if  $G$  is contact  $B_0$ -VPG; NO and a forbidden induced subgraph, if  $G$  is not contact  $B_0$ -VPG.

1. list all maximal cliques in  $G$ ;
2. if some edge belongs to two maximal cliques, return NO and  $K_4^-$ ;
3. if a maximal clique contains at least five vertices, return NO and  $K_5$ ;
4. label the vertices such that  $l(v) =$  number of  $K_4$ 's that  $v$  belongs to;
5. if for some vertex  $v$ ,  $l(v) \geq 3$ , return NO and  $H_0$ ;
6. if  $l(v) \leq 1 \forall v \in V \setminus \{w\}$  and  $l(w) \leq 2$ , return YES;



**Fig. 7.** An example of a possible running of the algorithm. The vertices marked in the algorithm are numbered in the order of the marking process. The vertex labelled  $R$  corresponds to the root of the tree in the forbidden structure, given in step 8 (whose other vertices are marked as 3 and 4).

7. while there exists an unmarked vertex  $v$  with  $2 - l(v)$  outgoing arcs incident to it, do
  - 7.1 mark  $v$  as internal;
  - 7.2 direct the edges that are currently undirected, uncoloured, not belonging to a  $K_4$ , and incident to  $v$  towards  $v$ ;
  - 7.3 for any two incoming arcs  $wv, w'v$  such that  $ww' \in E$ , colour  $ww'$ ;
8. if there exists some vertex  $v$  with more than  $2 - l(v)$  outgoing arcs, return NO and find  $T \in \mathcal{T}$  by running BFS starting with  $v$ , following the outgoing arcs, and adding for each vertex the corresponding  $K_4$ 's that it belongs to; else return YES.

An example of a possible running of the algorithm is illustrated in Fig. 7.

Steps 1–5 can clearly be done in polynomial time (see for example [16] for listing all maximal cliques in a chordal graph). Furthermore, it is obvious to see how to find the forbidden induced subgraph in steps 2, 3 and 5. Notice that if the algorithm has not returned NO after step 5, we know that  $G$  is  $\{K_4^-, K_5, H_0\}$ -free. So we are left with checking whether  $G$  contains some graph  $T \in \mathcal{T}$ ,  $T \neq H_0$ . Since each graph  $T \in \mathcal{T}$  contains at least two vertices belonging to two  $K_4$ 's, it follows that if at most one vertex has label 2,  $G$  is  $\mathcal{T}$ -free (step 6), and thus we conclude by Theorem 13 that  $G$  is contact  $B_0$ -VPG.

During step 7, we detect those vertices in  $G$  that, in case  $G$  is contact  $B_0$ -VPG, must be internal vertices (and mark them as such) and those vertices  $w$  that are middle neighbours of internal vertices  $v$  (we direct the edges  $wv$  from  $w$  to  $v$ ). Furthermore, we colour those edges whose endpoints are middle neighbours of a same internal vertex.

Consider a vertex  $v$  with  $2 - l(v)$  outgoing arcs. If a vertex  $v$  has  $l(v) = 2$ , then, in case  $G$  is contact  $B_0$ -VPG,  $v$  must be an internal vertex (see Lemma 9). This implies that any neighbour of  $v$ , which does not belong to a same  $K_4$  as  $v$ , must be a middle neighbour of  $v$ . If  $l(v) = 1$ , this means that  $v$  belongs to one  $K_4$  and is a middle neighbour of some internal vertex. Thus, by Lemma 9 we know that  $v$  is internal. Similarly, if  $l(v) = 0$ , this means that  $v$  is a middle neighbour of two distinct internal vertices. Again, by Lemma 9 we conclude that  $v$  is internal. Clearly, step 7 can be run in polynomial time.

So we are left with step 8, i.e., we need to show that  $G$  is contact  $B_0$ -VPG if and only if there exists no vertex with more than  $2 - l(v)$  outgoing arcs. First notice that only vertices marked as internal have incoming arcs. Furthermore, notice that every maximal clique of size three containing an internal vertex has two directed edges of the form  $wv, w'v$  and the third edge is coloured, where  $v$  is the first of the three vertices that was marked as internal. This is because the graph is  $K_4^-$ -free and the edges of a  $K_4$  are neither directed nor coloured.

**Lemma 15.** *Every vertex marked as internal in step 7 has either label 2 or is the root of a directed induced tree (directed from the root to the leaves) where the root  $w$  has degree  $2 - l(w)$  and every other vertex  $v$  has degree  $3 - l(v)$  in that tree, namely one incoming arc and  $2 - l(v)$  outgoing arcs.*

**Proof.** By induction in the number of iterations in step 7. In the first iteration, no edge has been directed. Therefore, any vertex marked as internal must have label 2, having zero outgoing edges. Now assume the result is true for any vertex marked before the  $n$ th iteration. Let  $v$  be the vertex marked in the  $n$ th iteration. If  $l(v) = 2$  we are done. Suppose  $l(v) = 1$ . Then, there is an outgoing edge from  $v$  to a vertex  $w$ . Since only vertices marked as internal have incoming arcs,  $w$  must be internal. Now, by inductive hypothesis ( $w$  was marked in a previous iteration), the result is true for  $w$ . If  $l(w) = 2$ ,  $v$  is the root of the tree consisting of the two vertices  $v$  and  $w$ , where  $v$  has degree  $2 - l(v) = 1$  and  $w$  has degree  $3 - l(w) = 1$  (one incoming arc). Otherwise,  $w$  is the root of a tree  $T'$  satisfying the hypothesis of the lemma, but then the tree  $T$  constructed from  $T'$  by adding  $v$  with an outgoing edge to  $w$  also clearly satisfies the hypothesis. In a similar manner can be constructed the tree in the case  $l(v) = 0$ . Finally, let us show that the tree is necessarily induced. Suppose there is an edge not in the tree that joins two vertices of the tree. Since the graph is a block graph, the vertices

in the resulting cycle induce a clique, so in particular there is a triangle formed by two edges of the tree and an edge not in the tree. But, as observed above, in every triangle of  $G$  having two directed edges, the edges point to the same vertex (and the third edge is coloured, not directed). Since no vertex in the tree has in-degree more than one, this is impossible.  $\square$

Based on the lemma, it is clear now that if a vertex has more than  $2 - l(v)$  outgoing arcs, then that vertex is the root of a directed induced tree (directed from the root to the leaves), where every vertex  $v$  has degree  $3 - l(v)$ , i.e., a tree that is the base tree  $B(T)$  of a graph  $T \in \mathcal{T}$ . Indeed, notice that every vertex  $v$  in a base tree has degree  $3 - l(v)$ . The fact that tree is induced can be proved the same way as above. This base tree can be found by a breadth-first search from a vertex having out-degree at least  $3 - l(v)$ , using the directed edges. Thanks to the labels, representing the number of  $K_4$ 's a vertex belongs to, it is then possible to extend the  $B(T)$  to an induced subgraph  $T \in \mathcal{T}$ . This can clearly be implemented to run in polynomial time.

To finish the proof that our algorithm is correct, it remains to show that if  $G$  contains an induced subgraph in  $\mathcal{T}$ , then the algorithm will find a vertex with at least  $3 - l(v)$  outgoing arcs. This, along with [Theorem 13](#), says that if the algorithm outputs YES then the graph is contact  $B_0$ -VPG (given that the detection of  $K_5$ ,  $K_4^-$  and  $H_0$  is clear). Recall that we know that  $G$  is a block graph after step 2. Notice that if a block of size 2 in a graph of  $\mathcal{T}$  is replaced by a block of size 4, we obtain either  $H_0$  or a smaller graph in  $\mathcal{T}$  as an induced subgraph. Moreover, adding an edge to a graph of  $\mathcal{T}$  in such a way that now contains a triangle, then we obtain a smaller induced graph in  $\mathcal{T}$ . Let  $G$  be a block graph with no induced  $K_5$  or  $H_0$ . By the remark above, if  $G$  contains a graph in  $\mathcal{T}$  as induced subgraph, then  $G$  contains one, say  $T$ , such that no edge of the base tree  $B(T)$  is contained in a  $K_4$  in  $G$ , and no triangle of  $G$  contains two edges of  $B(T)$ . So, all the edges of  $B(T)$  are candidates to be directed or coloured.

In fact, by step 7 of the algorithm, every vertex of  $B(T)$  is eventually marked as internal, and every edge incident with it is either directed or coloured, unless the algorithm ends with answer NO before. Notice that by the remark about the maximal cliques of size three and the fact that no triangle of  $G$  contains two edges of  $B(T)$ , if an edge  $vw$  of  $B(T)$  is coloured, then both  $v$  and  $w$  have an outgoing arc not belonging to  $B(T)$ . So, in order to obtain a lower bound on the out-degrees of the vertices of  $B(T)$  in  $G$ , we can consider only the arcs of  $B(T)$  and we can consider the coloured edges as bidirected edges. With an argument similar to the one in the proof of [Lemma 10](#), at least one vertex has out-degree at least  $3 - l(v)$ .

## Tree-cographs

In this section, we present a minimal forbidden induced subgraph characterisation for contact  $B_0$ -VPG graphs within the class of tree-cographs.

*Tree-cographs* [[24](#)] are a generalisation of cographs, i.e.  $P_4$ -free graphs. They are defined recursively as follows: trees are tree-cographs; the disjoint union of tree-cographs is a tree-cograph; and the complement of a tree-cograph is also a tree-cograph.

It follows from the definition that every tree-cograph is either a tree, or the complement of a tree, or the disjoint union of tree-cographs, or the join of tree-cographs. Let us start with the following two trivial facts.

**Fact 16.** *Every tree is a contact  $B_0$ -VPG graph.*

**Fact 17.** *The disjoint union of contact  $B_0$ -VPG graphs is contact  $B_0$ -VPG.*

Now let us consider the complement of trees. We obtain the following.

**Lemma 18.** *Let  $T$  be a tree. Then  $\bar{T}$  is contact  $B_0$ -VPG if and only if it is  $\{K_5, K_4^-\}$ -free.*

**Proof.** If  $\bar{T}$  is contact  $B_0$ -VPG, then it follows from [Lemma 5](#) that  $\bar{T}$  is  $\{K_5, K_4^-\}$ -free.


Suppose now that  $\bar{T}$  is  $\{K_5, K_4^-\}$ -free, then  $T$  has stability number at most 4. In particular, it has at most four leaves. Since it does not have co- $(K_4-e)$ 's either, we conclude that  $T$  is either a star with at most 4 leaves, a  $P_4$  or a  $P_5$ . Hence,  $\bar{T}$  is either a  $K_4 \cup K_1$ , a  $P_4$  or  $\bar{P}_5$ . Clearly, all these graphs are contact  $B_0$ -VPG.  $\square$

Using the previous results, we are able to obtain the following characterisation of tree-cographs that are contact  $B_0$ -VPG.

**Theorem 19.** *Let  $G$  be a tree-cograph. Then  $G$  is contact  $B_0$ -VPG if and only if  $G$  is  $\{K_5, K_{3,3}, H_0, K_4^-\}$ -free.*

**Proof.** If  $G$  is contact  $B_0$ -VPG, then it follows from [Lemma 5](#) that  $G$  is  $\{K_5, K_{3,3}, H_0, K_4^-\}$ -free.

Suppose now that  $G$  is a  $\{K_5, K_{3,3}, H_0, K_4^-\}$ -free tree cograph on  $n$  vertices. We will do a proof by induction on the number of vertices of  $G$ . Let us assume the theorem holds for graphs of less than  $n$  vertices. If  $G$  is a tree, the complement of a tree or the disjoint union of tree-cographs, then the result holds by [Facts 16, 17](#), [Lemma 18](#) and the induction hypothesis. So we may assume now that  $G$  is the join of two tree-cographs, say  $G_1, G_2$ .



**Fig. 8.** A graph  $G$  with  $G_1$  with a most two triangles and  $G_2 = K_1$ , and a contact  $B_0$ -VPG representation of  $G$ .

Since  $G$  is  $K_4^-$ -free, both  $G_1$  and  $G_2$  are  $P_3$ -free, i.e., they are the disjoint union of cliques. Furthermore, since  $G$  is  $K_5$ -free, it follows that  $\omega(G_1) + \omega(G_2) \leq 4$  and, in particular, none of  $G_1, G_2$  contains a  $K_4$ .

First suppose that one of  $G_1, G_2$ , say  $G_1$ , contains a triangle. Then  $G_2$  contains no  $K_2$ . But since  $G$  is  $K_4^-$ -free,  $G_2$  contains no  $2K_1$  either. So  $G_2$  is the trivial graph. Now, since  $G$  is  $H_0$ -free,  $G_1$  contains at most two triangles. But then  $G$  is clearly contact  $B_0$ -VPG. We show in Fig. 8 how to represent the join of the trivial graph and a graph consisting in the disjoint union of at most two triangles, an arbitrary number of edges and isolated vertices as a contact  $B_0$ -VPG graph.

Next suppose that  $\omega(G_1) = \omega(G_2) = 2$ . Since  $G$  is  $K_4^-$ -free, neither  $G_1$  nor  $G_2$  contains  $2K_1$ . So  $G = K_4$ , and hence it is contact  $B_0$ -VPG.

Suppose now  $\omega(G_1) = 2$  and  $\omega(G_2) = 1$ . Since  $G$  is  $K_4^-$ -free,  $G_2$  contains no  $2K_1$ , so  $G_2$  is the trivial graph and hence clearly contact  $B_0$ -VPG.

Finally, consider the case when  $\omega(G_1) = \omega(G_2) = 1$ . Since  $G$  is  $K_{3,3}$ -free, it follows that  $G$  is either the star  $K_{1,n-1}$  or the complete bipartite graph  $K_{2,n-2}$ . Thus again,  $G$  is clearly contact  $B_0$ -VPG.  $\square$

From the proofs of the previous results, the following fact can be deduced.

**Corollary 20.** Every contact  $B_0$ -VPG tree-cograph is the disjoint union of trees,  $\overline{P_5}$ 's, and contact  $B_0$ -VPG cographs.

### $P_4$ -Tidy graphs


Let  $G$  be a graph and let  $A$  be a vertex set that induces a  $P_4$  in  $G$ . A vertex  $v$  of  $G$  is said to be a *partner* of  $A$  if  $G[A \cup \{v\}]$  contains at least two induced  $P_4$ 's. The graph  $G$  is called  $P_4$ -tidy, if each vertex set  $A$  inducing a  $P_4$  in  $G$  has at most one partner [17]. The class of  $P_4$ -tidy graphs is an extension of the class of cographs, i.e.  $P_4$ -free graphs, and it contains many other graph classes defined by bounding the number of  $P_4$ 's according to different criteria; e.g.,  $P_4$ -sparse graphs [21],  $P_4$ -lite graphs [22], and  $P_4$ -extendible graphs [23].

A *spider* [21] is a graph whose vertex set can be partitioned into three sets  $S, C$ , and  $R$ , where  $S = \{s_1, \dots, s_k\}$  ( $k \geq 2$ ) is a stable set;  $C = \{c_1, \dots, c_k\}$  is a clique;  $s_i$  is adjacent to  $c_j$  if and only if  $i = j$  (a *thin spider*), or  $s_i$  is adjacent to  $c_j$  if and only if  $i \neq j$  (a *thick spider*);  $R$  is allowed to be empty and if it is not, then all the vertices in  $R$  are adjacent to all the vertices in  $C$  and non-adjacent to all the vertices in  $S$ . The triple  $(S, C, R)$  is called the *spider partition*. By  $\text{thin}_k(H)$  and  $\text{thick}_k(H)$  we respectively denote the thin spider and the thick spider with  $|C| = |S| = k$  and  $H$  the subgraph induced by  $R$ . If  $R$  is an empty set we denote them by  $\text{thin}_k$  and  $\text{thick}_k$ , respectively. Clearly, the complement of a thin spider is a thick spider, and vice versa. A *fat spider* is obtained from a spider by adding a true or false twin of a vertex  $v \in S \cup C$ . The following theorem characterises  $P_4$ -tidy graphs.

**Theorem 21 ([17]).** Let  $G$  be a  $P_4$ -tidy graph with at least two vertices. Then, exactly one of the following conditions holds:


1.  $G$  is disconnected.
2.  $\overline{G}$  is disconnected.
3.  $G$  is isomorphic to  $P_5, \overline{P_5}, C_5$ , a spider, or a fat spider.

This allows us to obtain the following characterisation of contact  $B_0$ -VPG  $P_4$ -tidy graphs.



**Fig. 9.** Representation of a thin spider  $(C, S, R)$  with  $|C| = 4$  and  $R$  empty.

Adding a true twin to a vertex of  $S$ . Adding a false twin to a vertex of  $S$ . Adding a false twin to a vertex of  $C$ ,  $|C| = 2$ .



**Fig. 10.**  $G$  is a fat spider arising from the thin spider  $(C, S, R)$ .

**Theorem 22.** Let  $G$  be a  $P_4$ -tidy graph. Then  $G$  is contact  $B_0$ -VPG if and only if  $G$  is  $\{K_5, K_{3,3}, H_0, K_4^-\}$ -free.

**Proof.** If  $G$  is a contact  $B_0$ -VPG graph, then it follows from Lemma 5 that  $G$  is  $\{K_5, K_{3,3}, H_0, K_4^-\}$ -free.

Suppose that  $G$  is a  $\{K_5, K_{3,3}, H_0, K_4^-\}$ -free  $P_4$ -tidy graph on  $n$  vertices. We will do a proof by induction on the number of vertices of  $G$ . Let us assume the theorem holds for graphs of less than  $n$  vertices. It follows from Theorem 21 that  $G$  is (i) either disconnected; (ii) or  $\bar{G}$  is disconnected; (iii) or  $G$  is isomorphic to  $P_5, \bar{P}_5, C_5$ , a spider, or a fat spider.

If  $G$  is disconnected,  $G$  is the union of  $P_4$ -tidy graphs. Thus the result holds by Fact 17 and the induction hypothesis.

If  $\bar{G}$  is disconnected, it follows that  $G$  is the join of two  $P_4$ -tidy graphs, say  $G_1, G_2$ . Then we do exactly the same case analysis as in the proof of Theorem 19.

Now suppose that  $G$  is a spider with partition  $(C, S, R)$ . Since  $G$  is  $K_4^-$ -free,  $G$  is necessarily a thin spider. Furthermore, since  $G$  is  $K_5$ -free, we have  $|C| \leq 4$ . If  $|C| = 4$ , then  $R$  must be empty. If  $|C| = 3$ , then  $|R| \leq 1$  because  $G$  is  $\{K_5, K_4^-\}$ -free. If  $|C| = 2$ , then, for the same reasons,  $|R| \leq 2$  and if  $|R| = 2$ , then  $R$  induces  $K_2$ . Notice that for all these cases, the graph obtained is an induced subgraph of the graph corresponding to the case  $|C| = 4$  and  $R = \emptyset$ . We provide a contact  $B_0$ -VPG representation of that case in Fig. 9.

Suppose now that  $G$  is a fat spider arising from the thin spider with partition  $(C, S, R)$ . Since  $G$  is  $K_4^-$ -free, it does not arise from adding a true twin to a vertex of  $C$ . For the same reason, if  $|C| \geq 3$ ,  $G$  does not arise from adding a false twin to a vertex of  $C$ , and if  $|C| = 2$ , we may add a false twin of a vertex of  $C$  only if  $R$  is empty. We provide a contact  $B_0$ -VPG representation for each of these remaining cases in Fig. 10.

Finally, it is easy to see that  $P_5, \bar{P}_5$ , and  $C_5$  are all contact  $B_0$ -VPG graphs.  $\square$

For  $P_4$ -tidy graphs a linear time recognition algorithm is known [17]. Using the decomposition properties of the class, the characterisation of the possible cases in the proof of [Theorem 19](#) for graphs with disconnected complement, and the possible cases in the proof of [Theorem 22](#) for spiders and fat spiders, we can obtain a linear-time algorithm to determine whether a  $P_4$ -tidy graph is contact  $B_0$ -VPG. Moreover, we can output a minimal forbidden induced subgraph in the case the answer is no.

### $P_5$ -Free contact $B_0$ -VPG graphs

In this section, we will present a characterisation of  $P_5$ -free contact  $B_0$ -VPG graphs. Notice that every  $P_k$ -free graph, with  $1 \leq k \leq 2$ , is clearly contact  $B_0$ -VPG. Moreover, a  $P_3$ -free graph  $G$  is a disjoint union of cliques, therefore  $G$  is contact  $B_0$ -VPG if and only if  $G$  is  $K_5$ -free.

Concerning  $P_4$ -free graphs, we have the following corollary of [Theorem 19](#) or [Theorem 22](#), since  $P_4$ -free graphs form a subclass of tree-cographs and  $P_4$ -tidy graphs.

**Theorem 23.** *Let  $G$  be a  $P_4$ -free graph. Then  $G$  is contact  $B_0$ -VPG if and only if  $G$  is  $\{K_5, K_{3,3}, H_0, K_4^-\}$ -free.*

Thus, the next graph class to consider is the class of  $P_5$ -free graphs. As we will see, the characterisation of  $P_5$ -free contact  $B_0$ -VPG graphs is much more complex than the characterisation of  $P_k$ -free graphs,  $k \leq 4$ . Consider a  $P_5$ -free graph  $G$ . If  $G$  is chordal, we obtain a characterisation using [Theorem 13](#). Hence, we may assume that  $G$  is non chordal. Since  $G$  is  $P_5$ -free it follows that  $G$  contains an induced cycle of length  $\ell \in \{4, 5\}$ . In what follows, we will first analyse the case when  $G$  contains an induced cycle of length four, but no induced cycle of length five.

**Lemma 24.** *Let  $G$  be a non chordal  $\{P_5, C_5, K_{3,3}, K_4^-\}$ -free graph. Then, there exists an induced cycle  $C$  of length four in  $G$  such that  $N[C] = G$ .*

**Proof.** Since  $G$  is not chordal but  $\{P_5, C_5\}$ -free, it follows that  $G$  must contain an induced cycle of length four. Let  $C_0$  be such a cycle induced by the vertices  $v_1, v_2, v_3, v_4$ . If  $N[C_0] = G$ , we are done. Suppose there exists a vertex  $v$  at distance two of  $C_0$ . So we may assume, without loss of generality, that there is a vertex  $a$  adjacent to  $v_1$  and  $v$ . It follows from [Remark 2](#) that  $a$  must be non-adjacent to at least one of  $v_2, v_4$ . Without loss of generality, we may assume that  $a$  is non-adjacent to  $v_4$ . But then  $a$  must be adjacent to  $v_3$ , otherwise  $v, a, v_1, v_4, v_3$  induce a  $P_5$ , a contradiction. Thus, by [Remark 2](#),  $a$  is non-adjacent to  $v_2$ .

Now, consider the cycle  $C_1$  induced by the vertices  $a, v_1, v_2, v_3$ . If  $N[C_1] = G$ , we are done. Suppose there is a vertex  $w$  at distance two of  $C_1$ . Notice that  $v, a, v_1, v_4$  induce a  $P_4$ . Thus,  $w$  cannot be adjacent to any of  $v, v_4$  otherwise we obtain a  $P_5$  or a  $C_5$ , a contradiction. Hence, there exists a vertex  $b \neq v, v_4$  adjacent to  $w$  and to some vertex in  $C_1$ . If  $b$  is adjacent to exactly one vertex in  $C_1$  or to exactly two consecutive vertices in  $C_1$ , we clearly obtain a  $P_5$ , a contradiction. Thus, it follows from [Remark 2](#), that  $b$  is adjacent to two nonconsecutive vertices in  $C_1$ . We distinguish two cases:

- (a)  $b$  is adjacent to  $a$  and  $v_2$ . Then  $b$  must be adjacent to  $v_4$ , otherwise  $w, b, a, v_1, v_4$  induce a  $P_5$ , a contradiction. But now  $v_1, v_3, b, a, v_2, v_4$  induce a  $K_{3,3}$ , a contradiction.
- (b)  $b$  is adjacent to  $v_1$  and  $v_3$ . Then  $b$  must be adjacent to  $v$ , otherwise  $w, b, v_1, a, v$  induce a  $P_5$ , a contradiction. Now consider the cycle  $C$  induced by  $a, v_1, b, v_3$ . We claim that  $N[C] = G$ . Suppose there is a vertex  $z$  at distance two of  $C$ . Then, following the same reasoning as above,  $z$  cannot be adjacent to any of  $v_4, v, w, v_2$ , since otherwise we obtain a  $P_5$  or  $C_5$ , a contradiction. Thus, as before for vertex  $b$ , there exists a vertex  $c$  adjacent to  $z$  and to two non-adjacent vertices of  $C$ . If  $c$  is adjacent to  $v_1$  and  $v_3$ , then  $c$  must also be adjacent to  $v$ , otherwise  $z, c, v_3, a, v$  induce a  $P_5$ . But now  $v_1, v_3, v, a, b, c$  induce a  $K_{3,3}$ , a contradiction. Using the same arguments, we can show that if  $c$  is adjacent to  $a, b$ , then it must be adjacent to  $v_2$ , and again we obtain an induced  $K_{3,3}$ , a contradiction. Thus  $z$  does not exist and hence,  $G = N[C]$ .  $\square$

We will define now the following family of graphs. Start with a cycle  $C$  induced by the vertices  $a_1, b_1, a_2, b_2$ . Add two (possibly empty) stable sets  $S_a, S_b$ , such that every vertex in  $S_a$  is adjacent to  $a_1, a_2$  (but not to  $b_1, b_2$ ), every vertex in  $S_b$  is adjacent to  $b_1, b_2$  (but not to  $a_1, a_2$ ) and  $S_a$  is anticomplete to  $S_b$ . Furthermore, add two (possibly empty) sets  $K_a, K_b$  such that  $K_a$  (resp.  $K_b$ ) is complete to  $\{a_1\}$  (resp.  $\{b_1\}$ ) and anticomplete to  $\{a_2, b_1, b_2\}$  (resp.  $\{a_1, a_2, b_2\}$ ). Also, every vertex in  $K_a$  (resp.  $K_b$ ) is a simplicial vertex of degree at most three and  $K_a$  (resp.  $K_b$ ) is anticomplete to  $S_a \cup S_b \cup K_b$  (resp.  $S_a \cup S_b \cup K_a$ ). Finally, add a (possibly empty) set  $K_{ab}$  of vertices forming a clique of size at most two that is complete to  $\{a_1, b_1\}$  and anticomplete to the rest of the graph. Moreover, neither of  $a_1, b_1$  can belong to three cliques of size four and only  $a_1$  may belong to two cliques of size four not containing any vertices from  $K_{ab}$ . There are no other edges in the graph. Let us denote by  $\mathcal{W}_1$  the family of graphs described here before (see [Fig. 11](#) for an example).

Let  $B_1, B_2$  and  $B_3$  be the graphs shown in [Fig. 12](#). Finally, let  $\mathcal{W} = \mathcal{W}_1 \cup \{B_1, B_2, B_3\}$ .

**Lemma 25.** *Let  $G$  be a non chordal  $\{P_5, C_5, K_5, K_{3,3}, H_0, G_{p_2}, \overline{C_6}, K_4^-\}$ -free graph. Then  $G \in \mathcal{W}$ .*




Fig. 11. (a) An example of a graph from the family  $\mathcal{W}_1$ . (b) The corresponding contact  $B_0$ -VPG representation.




Fig. 12. (a) The graphs  $B_1, B_2$  and  $B_3$ . (b) A contact  $B_0$ -VPG representation of  $B_3$ .

**Proof.** Let  $G$  be a non chordal  $\{P_5, C_5, K_5, K_{3,3}, H_0, G_{p_2}, \overline{C_6}, K_4^-\}$ -free graph. It follows from Lemma 24 that there exists an induced cycle  $C$  of length four in  $G$  such that  $N[C] = G$ . Let  $C$  be induced by vertices  $a_1, b_1, a_2, b_2$ . Let  $S_a$  (resp.  $S_b$ ) be the set of vertices adjacent to  $a_1, a_2$  but not  $b_1, b_2$  (resp. to  $b_1, b_2$  but not  $a_1, a_2$ ). Notice that  $S_a$  (resp.  $S_b$ ) must be a stable set since  $G$  is  $K_4^-$ -free. Furthermore,  $S_a$  is anticomplete to  $S_b$ . Indeed, if a vertex  $v \in S_a$  is adjacent to some vertex  $w \in S_b$  then  $a_1, a_2, w, b_1, b_2, v$  induce a  $K_{3,3}$ , a contradiction.

Now, suppose there is a vertex  $v$  in  $G$  adjacent to only one vertex in  $C$ . Without loss of generality, we may assume that  $v$  is adjacent to  $a_1$ . Then, it is not possible to have a vertex  $w \neq v$  in  $G$  adjacent only to  $a_2$  in  $C$ , since the vertices  $v, a_1, b_1, a_2, w$  would induce a  $P_5$  (in case  $v$  and  $w$  are non-adjacent) or a  $C_5$  (in case  $v$  and  $w$  are adjacent). Therefore, if there is a vertex  $w \neq v$  adjacent to only one vertex in  $C$  and different from  $a_1$ , then we may assume, without loss of generality, that it is adjacent to  $b_1$ . Let  $K_a$  (resp.  $K_b$ ) be the set of vertices adjacent to only  $a_1$  (resp.  $b_1$ ). If there is a vertex  $v \in K_a$  adjacent to a vertex  $w \in K_b$ , then  $v, w, b_1, a_2, b_2$  induce a  $P_5$ , a contradiction. Hence  $K_a$  is anticomplete to  $K_b$ .

Let us now show that all the vertices in  $K_a$  are simplicial. Indeed, suppose that  $v \in K_a$  is not simplicial. Then, there exists  $w, u \in N(v)$  such that  $u, w$  are non-adjacent. It follows from the above that  $u, w \in K_a$ . But then,  $v, w, u, a_1$  induce a  $K_4^-$ , a contradiction. By symmetry, all vertices in  $K_b$  are simplicial as well. We will distinguish two cases.

First assume now that  $G$  is  $\overline{P_5}$ -free. Thus every vertex not in  $C$  is adjacent to exactly 1 vertex in  $C$ , since  $G$  is  $K_4^-$ -free. We claim that  $S_a$  is anticomplete to  $K_a$ . Indeed, if a vertex  $v \in S_a$  is adjacent to some vertex  $w \in K_a$ , then  $a_1, b_1, a_2, v, w$  induce a  $\overline{P_5}$ , a contradiction. Similarly,  $S_b$  is anticomplete to  $K_b$ . Next, suppose that some vertex  $v \in S_a$  is adjacent to some vertex  $w \in K_b$ . If  $S_b$  is non empty, then for any vertex  $u \in S_b$  we obtain a  $P_5$  induced by  $b_2, u, b_1, w, v$ , a contradiction. Thus,  $S_b$  is empty. Then, we may redefine our cycle  $C$  by taking the vertices  $a_1, b_1, a_2, v$ . Notice that this cycle also verifies  $N[C] = G$ . Now,  $w \in S_b$  (where  $S_b$  is now the set of vertices adjacent to  $b_1, v$  but not to  $a_1, a_2$ ) and  $b_2 \in S_a$ . We can proceed similarly if  $S_a$  is empty and there are adjacent vertices in  $S_b$  and  $K_a$ . Now, since  $S_b \neq \emptyset, S_a$  (resp.  $S_b$ ) is anticomplete to  $K_b$  (resp.  $K_a$ ). Since  $G$  is  $K_5$ -free, it follows that the degree of the simplicial vertices is at most three. Finally, since  $G$  is  $\{H_0, G_{p_2}\}$ -free, it follows that only  $a_1$  can belong to two cliques of size four and neither of  $a_1, b_1$  can belong to three cliques of size four. Hence,  $G \in \mathcal{W}_1$ .

Now, suppose that  $G$  contains a  $\overline{P}_5$  induced by the cycle  $C$  and a vertex  $v$  adjacent to  $a_1$  and  $b_1$ . First, assume there are no other vertices in  $G$  adjacent to two consecutive vertices in  $C$ . Notice that  $v$  cannot be adjacent to any vertex in  $S_a \cup S_b \cup K_a \cup K_b$ , since  $G$  is  $K_4^-$ -free. Moreover,  $S_a$  is anticomplete to  $K_a$ . Indeed, if  $w \in K_a$  is adjacent to  $u \in S_a$ , then  $w, u, a_2, b_1, v$  induce a  $P_5$ , a contradiction. The same applies to  $K_b$  and  $S_b$ . Finally, we may assume that  $K_a$  (resp.  $K_b$ ) is anticomplete to  $S_b$  (resp.  $S_a$ ) by using the same arguments as above and redefining the cycle  $C$  if necessary. Hence,  $G$  belongs to  $\mathcal{W}_1$ .

Next, assume there is another vertex in  $G$  (in addition to  $v$ ) adjacent to two consecutive vertices in  $C$ . Notice that  $a_1$  and  $b_2$  (resp.  $a_2$  and  $b_1$ ) cannot have a common vertex since  $G$  is  $P_5$ -free. If there is another vertex  $w$  adjacent to  $a_1$  and  $b_1$ , but there is no vertex adjacent to  $a_2$  and  $b_2$ , then  $w$  must be adjacent to  $v$ , otherwise  $a_1, b_1, v, w$  induce a  $K_4^-$ , a contradiction. Also,  $a_1$  (resp.  $b_1$ ) cannot belong to two cliques of size four whose vertices belong to  $K_a \cup \{a_1\}$  (resp.  $K_b \cup \{b_1\}$ ), since  $G$  is  $H_0$ -free. Thus,  $G$  belongs to  $\mathcal{W}_1$ , since  $G$  is  $K_5$ -free and thus no further vertex is adjacent to both  $a_1$  and  $b_1$ . Finally, suppose there is a vertex  $w$  adjacent to  $a_2$  and  $b_2$ . First notice that  $v$  and  $w$  are non-adjacent, otherwise  $v, w, a_1, b_1, a_2, b_2$  induce a  $\overline{C}_6$ , a contradiction. We claim that all the sets  $S_a, S_b, K_a$  and  $K_b$  must be empty. Indeed, if  $u \in S_a$ , then  $u$  is non-adjacent to  $w$ , since  $G$  is  $K_4^-$ -free. But then  $w, a_2, u, a_1, v$  induce a  $P_5$ , a contradiction. Thus,  $S_a = \emptyset$  and by symmetry we also conclude that  $S_b = \emptyset$ . Now suppose  $u \in K_a$ . Then the vertices  $u, a_1, b_1, a_2$  and  $w$  induce a  $P_5$  (if  $u, w$  are non-adjacent) or a  $C_5$  (if  $u, w$  are adjacent). Hence  $K_a = \emptyset$  and by symmetry  $K_b = \emptyset$ . If there are no more vertices,  $G$  is isomorphic to  $B_1$ . If there are more vertices in  $G$ , then by using the same arguments as before, these vertices have to be common neighbours of  $a_1$  and  $b_1$ , or  $a_2$  and  $b_2$ . But then  $G$  is necessarily isomorphic to either  $B_2$  or  $B_3$ , since  $G$  is  $\{K_5, \overline{C}_6, K_4^-\}$ -free (the same arguments as before apply again).

Finally assume that the  $\overline{P}_5$  contained in  $G$  is not induced by the cycle  $C$  together with some vertex  $v$  adjacent to two consecutive vertices in  $C$ . The only possibility is that the house is induced by  $a_1, b_1, a_2, u, w$ , with  $u \in S_a$  and  $w \in K_a$  (resp.  $a_1, b_1, b_2, u, w$ , with  $u \in S_b$  and  $w \in K_b$ ). But then, we may redefine our cycle  $C$  by taking the vertices  $a_1, b_1, a_2, u$  (resp.  $a_1, b_1, b_2, u$ ). Clearly this new cycle  $C$  also verifies that  $N[C] = G$ . Thus, we can apply the same arguments as before and show that  $G \in \mathcal{W}$ .  $\square$

**Lemma 26.** *Every graph in  $\mathcal{W}$  is contact  $B_0$ -VPG.*

**Proof.** Let  $G$  be a graph in  $\mathcal{W}_1$ . We construct a contact  $B_0$ -VPG representation of  $G$  as follows. First represent the main cycle  $C$  induced by  $a_1, b_1, a_2, b_2$ :  $P_{a_1}$  is a horizontal path lying on row  $x_i$ ;  $P_{a_2}$  is a horizontal path lying on row  $x_j$ ,  $j < i$ ;  $P_{b_1}$  is a vertical path lying on column  $y_k$ ;  $P_{b_2}$  is a vertical path lying on column  $y_\ell$ , with  $\ell > k + |S_a|$ ; furthermore, we make sure that  $b_1$  and  $b_2$  are middle-neighbours of  $a_1$  and  $a_2$  is a middle neighbour of  $b_1$  and  $b_2$ ; finally the paths  $P_{b_1}$  and  $P_{b_2}$  use column  $y_k$  respectively  $y_\ell$  down to row  $x_t$  with  $t + |S_b| < j$ . Now, each vertex in  $S_a$  can be represented by a vertical path on some column  $y_r$ , with  $k < r < \ell$ , and every vertex in  $S_b$  can be represented by a horizontal path on some row  $u$  with  $t < u < j$ . First assume that  $K_{ab} = \emptyset$ . Since  $P_{a_1}$  has both endpoints free, one can easily represent two cliques of size four, in case  $a_1$  belongs to such cliques and similarly, since  $P_{b_1}$  has one endpoint free, one can easily represent one clique of size four, in case  $b_1$  belongs to such a clique. All other vertices in  $K_a$  or  $K_b$  can clearly be represented by extending enough the paths  $P_{a_1}$  and  $P_{b_1}$ .

Now, assume that  $K_{ab} = \{v\}$ . Then, given a contact  $B_0$ -VPG representation of  $G - v$  as described before, we can easily obtain a contact  $B_0$ -VPG representation of  $G$  as follows: we add a path  $P_v$  lying on column  $y_k$  between some row  $x_q$  and row  $x_i$ , with  $i < q$ .

Next, assume that  $K_{ab} = \{v, v'\}$ . Thus,  $a_1$  belongs to at most one clique of size four in  $G - \{v, v'\}$  (the vertices of that clique belong to  $K_a$ , except for  $a_1$ ). We obtain a contact  $B_0$ -VPG representation as follows. Start with a contact  $B_0$ -VPG representation of  $G - v'$  as described above. Make sure that all vertices in  $K_a$  are represented by paths intersecting  $P_{a_1}$  to the right of column  $y_\ell$  (this is clearly always possible, since  $a_1$  belongs to at most one clique of size four whose vertices (except for  $a_1$ ) belong to  $K_a$ ). Finally, if necessary, reduce  $P_{a_1}$  such that its left endpoint corresponds to the grid point  $(x_i, y_k)$  (this is possible since  $P_{a_1}$  does not intersect any path to the left of that grid point anymore). Now add  $P_w$  as a horizontal path on row  $x_i$  with its right endpoint corresponding to the grid point  $(x_i, y_k)$ .

Finally, if  $G$  is one of the graphs  $B_1, B_2$  or  $B_3$ , then  $G$  is clearly contact  $B_0$ -VPG as can be seen in Fig. 12(b). Notice that  $B_1, B_2$  are induced subgraphs of  $B_3$ .  $\square$

From the lemmas above, we conclude the following.

**Corollary 27.** *Let  $G$  be a non chordal  $\{P_5, C_5, K_5, K_{3,3}, H, G_{p_2}, \overline{C}_6, K_4^-\}$ -free graph. Then  $G$  is contact  $B_0$ -VPG.*

Let us now focus on  $P_5$ -free graphs containing an induced cycle of length five.

**Lemma 28.** *Let  $G$  be a  $\{P_5, K_4^-\}$ -free graph. Let  $C$  be an induced cycle of length five in  $G$  such that no vertex is adjacent to exactly three non consecutive vertices in  $C$ . Then,  $N[C] = G$  and every vertex  $v \in N(C)$  is adjacent to exactly two non-consecutive vertices in  $C$ .*

**Proof.** Let  $C$  be induced by  $v_1, \dots, v_5$  and let  $v$  be a vertex in  $N(C)$ . It follows from Remark 2 that  $v$  cannot be adjacent to three consecutive vertices in  $C$ . If  $v$  is adjacent to exactly one vertex or to two consecutive vertices in  $C$ , then we clearly obtain a  $P_5$ , a contradiction. Thus,  $v$  has exactly two non consecutive neighbours in  $C$ .




Fig. 13. (a) An example of a graph in  $\mathcal{L}_2$ . (b) The corresponding contact  $B_0$ -VPG representation.




Fig. 14. The graphs  $G_1$ ,  $G_2$ ,  $G_3$  and  $G_4$ .

Now assume that there exists a vertex  $u$  which is at distance two of  $C$ . Thus, there is a vertex  $w \in N(C)$  adjacent to  $u$  and to two non consecutive vertices in  $C$ , say  $v_1, v_3$ . But then,  $v, w, v_1, u_5, v_4$  induce a  $P_5$ , a contradiction. Therefore  $N[C] = G$ .  $\square$

Let  $K_{3,3}^*$  be the graph obtained by subdividing exactly one edge in the graph  $K_{3,3}$ . We will now define several families of graphs. Start with a cycle  $C$  of length five induced by the vertices  $a, v, b, c, w$ . Add two (possibly empty) stable sets  $S_v, S_w$  such that  $S_v$  is complete to  $\{a, b\}$ ,  $S_w$  is complete to  $\{a, c\}$  and  $S_v$  is anticomplete to  $S_w$ . There are no other edges. Let us denote by  $\mathcal{L}_1$  the family of graphs described here before.

Let  $G \in \mathcal{L}_1$  and let  $G'$  be the graph obtained from  $G$  by adding a vertex  $u$  adjacent to  $a, b$  and  $c$ . Furthermore, add a (possible empty) set  $K_u$ , such that  $K_u$  is complete to  $\{u\}$  and anticomplete to  $V(C) \cup S_v \cup S_w$ . Also, every vertex in  $K_u$  is a simplicial vertex of degree at most three. Moreover,  $u$  can belong to only one clique of size four. There are no other edges. Let us denote by  $\mathcal{L}_2$  the family of graphs described here before (see Fig. 13(a) for an example).

Next, consider a graph  $G'$  in  $\mathcal{L}_2$  with  $S_v = S_w = \emptyset$  and  $u$  not belonging to any clique of size four. Add a vertex  $z$  adjacent to  $v, w$  and  $u$ . There are no other edges. Let us denote by  $\mathcal{L}_3$  the family of graphs obtained that way and let  $\mathcal{L} = \mathcal{L}_1 \cup \mathcal{L}_2 \cup \mathcal{L}_3$ .


Finally, let  $G_1, G_2, G_3$  and  $G_4$  be the graphs shown in Fig. 14.

**Lemma 29.** *Let  $G$  be a  $\{P_5, K_5, K_{3,3}^*, \overline{C_6}, G_1, G_2, G_3, G_4, K_4^-\}$ -free graph and assume  $G$  contains a cycle of length five. Then  $G \in \mathcal{L}$ .*

**Proof.** Let  $C$  be an induced cycle of length five with vertices  $a, v, b, c, w$ . Clearly, no vertex in  $N(C)$  is adjacent to exactly one vertex in  $C$  or to two consecutive vertices in  $C$ , since  $G$  is  $P_5$ -free. Consider first the vertices adjacent to two non-consecutive vertices in  $C$ . For any two vertices  $u, z$  that are adjacent to the same two non-consecutive vertices in  $C$ , we have that  $u$  and  $z$  are non-adjacent otherwise we obtain  $K_4^-$ , a contradiction. Suppose that there exist vertices  $u, z$  such that they have distinct neighbours in  $C$ , say  $u$  is adjacent to  $a$  and  $b$ , and  $z$  is adjacent to  $v$  and  $c$ . If  $u, z$  are adjacent, then together with the vertices of  $C$ , they induce a  $K_{3,3}^*$ , a contradiction. If  $u, z$  are non-adjacent, then  $u, a, v, z, c$  induce a  $P_5$ , a contradiction. Thus, we may assume now, without loss of generality, that every vertex adjacent to two nonconsecutive vertices in  $C$  is either adjacent to both  $a$  and  $b$  or adjacent to both  $a$  and  $c$ . Let  $S_v$  (resp.  $S_w$ ) be the set of vertices not in  $C$  adjacent to  $a, b$  (and not to  $v, c, w$ ) (resp.  $a, c$  (and not to  $b, v, w$ )). It follows from the above that  $S_v$  and  $S_w$  are stable sets. Finally, if there is a vertex  $u \in S_v$  adjacent to some vertex  $z \in S_w$ , then we obtain  $G_2$ , a contradiction. So  $S_v$  is anticomplete to  $S_w$ .

First assume that there exists no vertex in  $G$  that is adjacent to three non-consecutive vertices in  $C$ . It immediately follows from Lemma 28 that  $G = N[C]$  and that every vertex not in  $C$  is adjacent to two non-consecutive vertices in  $C$ . Thus,  $G \in \mathcal{L}_1$ .

Now, suppose that there exists a vertex  $u$  adjacent to three non-consecutive vertices in  $C$ , say  $a, b$  and  $c$ . We will first show that there cannot be another vertex adjacent to three non-consecutive vertices. If there is another vertex  $z$  adjacent



**Fig. 15.** A graph in  $\mathcal{L}_3$  and the corresponding contact  $B_0$ -VPG representation.

to  $a, b$  and  $c$ , then  $u$  and  $z$  must be adjacent otherwise  $u, z, c, b$  induce a  $K_4^-$ , a contradiction. But now the vertices  $a, u, z$  and  $b$  induce a  $K_4^-$ , again a contradiction. Now, suppose  $z$  is adjacent to  $v, b$  and  $w$ . Then,  $z$  and  $u$  are non-adjacent, since otherwise  $u, z, b, c$  induce a  $K_4^-$ , a contradiction. But now, the vertices of  $C$  together with  $u$  and  $z$  induce  $G_3$ , a contradiction as well. By symmetry, we conclude that  $z$  cannot be adjacent to  $v, w$  and  $c$ . Finally, if  $z$  is adjacent to  $a, v$  and  $c$ , the vertices  $a, v, z, u, b$  and  $c$  induce a  $\overline{C_6}$  if  $z$  and  $u$  are non-adjacent, a contradiction. But if  $z$  and  $u$  are adjacent, then  $u, z, b, c$  induce a  $K_4^-$ , again a contradiction. By symmetry,  $z$  cannot be adjacent to  $a, w$  and  $b$ . Hence, we conclude that  $u$  is the unique vertex adjacent to three non-consecutive vertices in  $C$ .

Now we will distinguish several cases, depending on which vertices  $u$  is adjacent to. First, assume that  $u$  is adjacent to  $a, b$  and  $c$ , and that  $S_v \cup S_w$  is non empty. Notice that  $u$  cannot be adjacent to any vertex in  $S_v \cup S_w$ , since  $G$  is  $K_4^-$ -free. It follows from Remark 2 and the fact that  $G$  is  $P_5$ -free that any vertex in  $G$  not belonging to  $V(C) \cup S_v \cup S_w \cup \{u\}$  has to be adjacent to  $u$  and anticomplete to  $V(C) \cup S_v \cup S_w$ . Let  $K_u = N(u) \setminus V(C)$  be the set of these vertices and consider  $z \in K_u$ . Then  $z$  is simplicial. Indeed, if  $z$  is not simplicial, it follows that there exist vertices  $z', z'' \in K_u \cap N(z)$  such that  $z', z''$  are non-adjacent. But then  $z, z', z'', u$  induce  $K_4^-$ , a contradiction. Furthermore, since  $G$  is  $K_5$ -free, it follows that every vertex  $z \in K_u$  has degree at most three. Finally, notice that  $u$  can only belong to at most one clique of size four, since  $G$  is  $G_1$ -free. Thus, we conclude that  $G \in \mathcal{L}_2$ .

Notice that if  $S_v = S_w = \emptyset$ , we can relabel the vertices in  $C$  such that  $u$  is adjacent to  $a, b$  and  $c$ , and we obtain a graph in  $\mathcal{L}_2$  as before. Thus, we may assume, without loss of generality, that there is a vertex  $z \in S_v$ . Now, we will consider different cases:

- If  $u$  is adjacent to  $v, b$  and  $w$ , or if  $u$  is adjacent to  $a, v$  and  $c$ , then we obtain  $G_2$  (notice that  $z$  and  $u$  cannot be adjacent since the graph is  $K_4^-$ -free), a contradiction.
- If  $u$  is adjacent to  $a, b$  and  $w$ , then  $S_w = \emptyset$ , otherwise  $a, v, b, c, u, w, t$ , where  $t \in S_w$ , induce  $G_2$  a contradiction. Now, we can relabel the vertices in  $C$  such that  $u$  is adjacent to  $a, b$  and  $c$ , without changing  $S_v$ , and we obtain a graph in  $\mathcal{L}_2$  as before.
- If  $u$  is adjacent to  $v, c$  and  $w$ , and  $z$  is non-adjacent to  $u$ , then  $z, a, v, u, c$  induce a  $P_5$ , a contradiction. So  $z$  and  $u$  must be adjacent. Notice again that  $S_w = \emptyset$ . Indeed, if  $t \in S_w$ , then  $t, a, v, b, u, c, w$  induce  $G_2$ , a contradiction. Moreover,  $|S_v| = 1$ : if  $z' \in S_v, z \neq z'$ , then  $z'$  must be adjacent to  $u$  as well, but now  $v, z, z', a, b, u$  induce a  $K_{3,3}$ , a contradiction. So we can relabel the vertices in  $C$  such that  $u$  is adjacent to  $a, b, c$ . With this new labelling,  $S_v = S_w = \emptyset$  and  $z$  is adjacent to  $v, w$  and  $u$ . Clearly, any vertex not belonging to  $V(C) \cup \{u, z\}$  has to be adjacent to  $u$ , since  $G$  is  $P_5$ -free. Let  $K_u$  be the set of these vertices. Using the same arguments than above, one can show that every vertex in  $K_u$  is simplicial and have degree at most three since the graph is  $K_5$ -free. Finally,  $u$  cannot belong to a clique of size four, since  $G$  is  $G_4$ -free. So we conclude that  $G \in \mathcal{L}_3$ .  $\square$

**Lemma 30.** Every graph in  $\mathcal{L}$  is contact  $B_0$ -VPG.

**Proof.** Let  $G \in \mathcal{L}_1$ . We construct a contact  $B_0$ -VPG representation of  $G$  as follows. Vertex  $b$  is represented by a path  $P_b$  lying on column  $y_j$  between rows  $x_k$  and  $x_t$ , with  $t > k + |S_v|$ ; vertex  $c$  is represented by a path  $P_c$  lying on column  $y_j$  between rows  $x_t$  and  $x_\ell$ , with  $\ell > t + |S_w|$ ; vertex  $a$  is represented by a path  $P_a$  lying on column  $y_i, i < j$ , between rows  $x_k$  and  $x_\ell$ ; vertex  $v$  is represented by a path  $P_v$  lying on row  $x_k$  between rows  $y_i$  and  $y_j$  and vertex  $w$  is represented by a path  $P_w$  lying on row  $x_\ell$  between rows  $y_i$  and  $y_j$ . Now each vertex in  $S_v$  is represented by a path between columns  $y_i$  and  $y_j$  lying on one of the  $|S_v|$  rows between  $x_k$  and  $x_t$ , and each vertex in  $S_w$  is represented by a path between columns  $y_i$  and  $y_j$  lying on one of the  $|S_w|$  rows between  $x_t$  and  $x_\ell$ .

If  $G \in \mathcal{L}_2$ , consider a representation of  $G - (K_u \cup \{u\})$  as described above. Now, it is possible to add  $P_u$  on row  $x_t$ , such that  $b$  and  $c$  are middle-neighbours of  $u$ , and  $u$  is a middle-neighbour of  $a$ . If  $u$  belongs to one clique of size four, then it is possible to represent this clique using the right endpoint of  $P_u$ . All the other vertices of  $K_u$  can easily be represented by eventually extending the path  $P_u$  to the right.

Finally, if  $G \in \mathcal{L}_3$ , consider the contact  $B_0$ -VPG representation of the graph shown in Fig. 15. Clearly, it is possible to add the paths representing the vertices of  $K_u$ , since  $u$  does not belong to any clique of size four.  $\square$

**Lemma 31.** The graphs  $K_{3,3}^*, \overline{C_6}, G_1, G_2, G_3, G_4$  are not contact  $B_0$ -VPG.

**Proof.** Consider the graph  $K_{3,3}$  with vertices  $a, c, e$  on one side of the bipartition and  $b, d, f$  on the other side. Assume that the edge  $ef$  is subdivided to obtain  $K_{3,3}^*$ . Consider the cycle induced by the vertices  $a, b, c, d$ . Following the same approach as in [Lemma 5](#), we may assume that  $P_a, P_c$  are horizontal paths,  $P_b, P_d$  are vertical paths and  $P_e$  is a horizontal path lying inside the rectangle, and  $P_f$  is a vertical path lying outside the rectangle. But now it is clearly impossible to add a path intersecting  $P_e$  and  $P_f$  without intersecting any other path. Thus,  $K_{3,3}^*$  is not  $B_0$ -VPG.

Next consider the graph  $\overline{C_6}$  with vertex set  $a, b, c, d, v, w$  such that  $a, b, c, d$  induce a cycle of length four,  $v$  is a common vertex of  $a$  and  $b$ ,  $w$  is a common neighbour of  $c$  and  $d$ , and  $v$  is adjacent to  $w$ . If  $\overline{C_6}$  is contact  $B_0$ -VPG, then we may assume that in a contact  $B_0$ -VPG representation, the paths  $P_a, P_c$  are horizontal and the paths  $P_b, P_d$  are vertical. Since  $b, c, v, w$  induce a cycle of length four, we conclude from the above that  $P_v$  has to be horizontal. But since  $a, d, v, w$  induce a cycle of length four as well, we also conclude that  $P_v$  has to be vertical, a contradiction. Hence,  $\overline{C_6}$  is not  $B_0$ -VPG.

Suppose now that the graph  $G_1$  is contact  $B_0$ -VPG. Without loss of generality, we may assume that  $P_u$  lies on some row  $x_i$ . Since  $u$  belongs to two cliques of size four, it follows from [Remark 3](#) that both endpoints of  $P_u$  are not free. Thus,  $a, b$  and  $c$  are middle neighbours of  $u$ , i.e.  $P_a, P_b, P_c$  are necessarily vertical paths. Thus,  $P_v, P_w$  must be horizontal paths, but this is impossible since no two paths can cross. We conclude that  $G_1$  is not contact  $B_0$ -VPG.

Using similar arguments, we conclude that if  $G_4$  is contact  $B_0$ -VPG, then  $b, c$  have to be middle neighbours of  $u$ ,  $u$  has to be a middle neighbour of  $a$  and  $P_v, P_w$  have to be horizontal paths. But now it is clearly impossible to add  $P_z$  such that it intersects  $P_v, P_w, P_u$  without crossing any path. Hence,  $G_4$  is not contact  $B_0$ -VPG.

Finally, consider the graphs  $G_2, G_3$  and suppose that they are contact  $B_0$ -VPG. First consider a contact  $B_0$ -VPG representation of  $G_2 - v$  (resp.  $G_3 - v$ ). Since  $t$  is adjacent to three non-consecutive vertices of a induced cycle of length five, we may assume, without loss of generality, that we have the following configuration:  $P_a, P_c, P_z$  are horizontal paths with  $P_a, P_z$  lying on a same row;  $P_b, P_w$  are vertical paths;  $P_t$  is a vertical path with one endpoint corresponding to the endpoints of  $P_a, P_z$  that intersect;  $t$  is a middle neighbour of  $c$ . But now it is clearly impossible to add a path representing vertex  $v$ , since it has to intersect  $P_a$  and  $P_b$ . Therefore,  $G_2, G_3$  are not contact  $B_0$ -VPG.  $\square$

We are now ready to prove the main result of this section.

**Theorem 32.** *Let  $G$  be a  $P_5$ -free graph. Let  $\mathcal{G} = \{K_5, H_0, G_{p_2}, K_{3,3}, K_{3,3}^*, \overline{C_6}, G_1, G_2, G_3, G_4, K_4^-\}$ . Then  $G$  is contact  $B_0$ -VPG if and only if  $G$  is  $\mathcal{G}$ -free.*

**Proof.** For the only if part, we use [Theorem 13](#), [Lemmas 5](#) and [31](#).

Suppose now that  $G$  is a  $P_5$ -free graph which is also  $\mathcal{G}$ -free. If  $G$  is chordal, the result follows from [Theorem 13](#), since  $G$  is  $\mathcal{F}$ -free (indeed, the graphs in  $\mathcal{F}$  different from  $H_0$  and  $G_{p_2}$  contain an induced  $P_5$ ). Now, assume that  $G$  is not chordal. If  $G$  is  $C_5$ -free, by [Corollary 27](#),  $G$  is contact  $B_0$ -VPG. Similarly, if  $G$  contains a  $C_5$ , by [Lemmas 29](#) and [30](#),  $G$  is also contact  $B_0$ -VPG.  $\square$

## Conclusions and future work

In this paper, we considered some special graph classes, namely chordal graphs, tree-cographs,  $P_4$ -tidy graphs and  $P_5$ -free graphs. We gave a characterisation by minimal forbidden induced subgraphs of those graphs from these families that are contact  $B_0$ -VPG. Moreover, we presented a polynomial-time algorithm for recognising chordal contact  $B_0$ -VPG graphs based on our characterisation. Notice that for the other graph classes considered here, the characterisation immediately yields a polynomial-time recognition algorithm.

In order to get a better understanding of the structure of general contact  $B_0$ -VPG graphs, one way could be to find further characterisations by forbidden induced subgraphs of contact  $B_0$ -VPG graphs within other interesting classes. Since classical graph problems are difficult in contact  $B_0$ -VPG graphs (see for instance [\[13\]](#)), these further insights in their structure may lead to good approximation algorithms for these problems.

## Acknowledgements

This work was partially supported by ANPCyT PICT-2015-2218 and UBACyT Grants 20020130100808BA and 20020160100095BA (Argentina).

## References

- [1] L. Alc3n, F. Bonomo, M.P. Mazzoleni, Vertex intersection graphs of paths on a grid: characterization within block graphs, *Graphs Combin.* 33 (4) (2017) 653–664.
- [2] A. Asinowski, E. Cohen, M.C. Golumbic, V. Limouzy, M. Lipshteyn, M. Stern, Vertex intersection graphs of paths on a grid, *J. Graph Algorithms Appl.* 16 (2) (2012) 129–150.
- [3] J.A. Bondy, U.S.R. Murty, *Graph Theory*, Springer, New York, 2007.
- [4] F. Bonomo, M.P. Mazzoleni, M.L. Rean, B. Ries, Characterising chordal contact  $B_0$ -VPG graphs, in: J. Lee, G. Rinaldi, A. Ridha Mahjoub (Eds.), *Proceedings of the International Symposium on Combinatorial Optimization 2018*, in: *Lecture Notes in Computer Science*, vol. 10856, 2018, pp. 89–100.

- [5] S. Chaplick, E. Cohen, J. Stacho, Recognizing some subclasses of vertex intersection graphs of 0-bend paths in a grid, in: P. Kolman, J. Kratochvíl (Eds.), Proceedings of the International Workshop on Graph-Theoretic Concepts in Computer Science 2011, in: Lecture Notes in Computer Science, vol. 6986, 2011, pp. 319–330.
- [6] S. Chaplick, V. Jelínek, J. Kratochvíl, T. Vyskocil, Bend-bounded path intersection graphs: Sausages, noodles, and waffles on a grill, in: M.C. Golumbic, M. Stern, A. Levy, G. Morgenstern (Eds.), Proceedings of the International Workshop on Graph-Theoretic Concepts in Computer Science 2012, in: Lecture Notes in Computer Science, vol. 7551, 2012, pp. 274–285.
- [7] S. Chaplick, T. Ueckerdt, Planar graphs as VPG-graphs, *J. Graph Algorithms Appl.* 17 (4) (2013) 475–494.
- [8] E. Cohen, M.C. Golumbic, B. Ries, Characterizations of cographs as intersection graphs of paths on a grid, *Discrete Appl. Math.* 178 (2014) 46–57.
- [9] E. Cohen, M.C. Golumbic, W.T. Trotter, R. Wang, Posets and VPG graphs, *Order* 33 (1) (2016) 39–49.
- [10] N. de Castro, F.J. Cobos, J.C. Dana, A. Márquez, M. Noy, Triangle-free planar graphs as segments intersection graphs, in: J. Kratochvíl (Ed.), Proceedings of the International Symposium on Graph Drawing and Network Visualization 1999, in: Lecture Notes in Computer Science, vol. 1731, 1999, pp. 341–350.
- [11] H. de Fraysseix, P. Ossona de Mendez, Representations by contact and intersection of segments, *Algorithmica* 47 (4) (2007) 453–463.
- [12] H. de Fraysseix, P. Ossona de Mendez, J. Pach, Representation of planar graphs by segments, *Intuitive Geom.* 63 (1991) 109–117.
- [13] Z. Deniz, E. Galby, A. Munaro, B. Ries, On contact graphs of paths on a grid, in: T. Biedl, A. Kerren (Eds.), Proceedings of the International Symposium on Graph Drawing and Network Visualization 2018, in: Lecture Notes in Computer Science, vol. 11282, 2018, pp. 317–330.
- [14] G. Ehrlich, S. Even, R. Tarjan, Intersection graphs of curves in the plane, *J. Combin. Theory Ser. B* 21 (1976) 8–20.
- [15] S. Felsner, K. Knauer, G.B. Mertzios, T. Ueckerdt, Intersection graphs of L-shapes and segments in the plane, *Discrete Appl. Math.* 206 (2016) 48–55.
- [16] P. Galinier, M. Habib, C. Paul, Chordal graphs and their clique graphs, in: M. Nagl (Ed.), Proceedings of the International Workshop on Graph-Theoretic Concepts in Computer Science 1995, in: Lecture Notes in Computer Science, vol. 1017, Springer, Berlin, Heidelberg, 1995, pp. 358–371.
- [17] V. Giakoumakis, F. Roussel, H. Thuillier, On  $P_4$ -tidy graphs, *Discrete Math. Theor. Comput. Sci.* 1 (1997) 17–41.
- [18] M.C. Golumbic, B. Ries, On the intersection graphs of orthogonal line segments in the plane: characterizations of some subclasses of chordal graphs, *Graphs Combin.* 29 (2013) 499–517.
- [19] P. Hliněný, Classes and recognition of curve contact graphs, *J. Combin. Theory Ser. B* 74 (1) (1998) 87–103.
- [20] P. Hliněný, Contact graphs of line segments are NP-complete, *Discrete Math.* 235 (1) (2011) 95–106.
- [21] C.T. Hoàng, Perfect graphs, (Ph.D. thesis), School of Computer Science, McGill University, Montreal, 1985.
- [22] B. Jamison, S. Olariu, A new class of brittle graphs, *Stud. Appl. Math.* 81 (1989) 89–92.
- [23] B. Jamison, S. Olariu, On a unique tree representation for  $P_4$ -extendible graphs, *Discrete Appl. Math.* 34 (1991) 151–164.
- [24] G. Tinhofer, Strong tree-cographs are Birkhoff graphs, *Discrete Appl. Math.* 22 (3) (1989) 275–288.