



Matching Cuts in Graphs of High Girth and H -Free Graphs

Carl Feghali¹ · Felicia Lucke² · Daniël Paulusma³ · Bernard Ries²

Received: 29 September 2023 / Accepted: 9 April 2025
© The Author(s) 2025

Abstract

The (PERFECT) MATCHING CUT problem is to decide if a connected graph has a (perfect) matching that is also an edge cut. The DISCONNECTED PERFECT MATCHING problem is to decide if a connected graph has a perfect matching that contains a matching cut. Both MATCHING CUT and DISCONNECTED PERFECT MATCHING are NP-complete for planar graphs of girth 5, whereas PERFECT MATCHING CUT is known to be NP-complete even for subcubic bipartite graphs of arbitrarily large fixed girth. We prove that MATCHING CUT and DISCONNECTED PERFECT MATCHING are also NP-complete for bipartite graphs of arbitrarily large fixed girth and bounded maximum degree. Our result for MATCHING CUT resolves a 20-year old open problem. We also show that the more general problem d -CUT, for every fixed $d \geq 1$, is NP-complete for bipartite graphs of arbitrarily large fixed girth and bounded maximum degree. Furthermore, we show that MATCHING CUT, PERFECT MATCHING CUT and DISCONNECTED PERFECT MATCHING are NP-complete for H -free graphs whenever H contains a connected component with two vertices of degree at least 3. Afterwards, we update the state-of-the-art summaries for H -free graphs and compare them with each other, and with a known and full classification of the MAXIMUM MATCHING CUT problem, which is to determine a largest matching cut of a graph G . Finally, by combining existing results, we obtain a complete complexity classification of PERFECT MATCHING CUT for \mathcal{H} -subgraph-free graphs where \mathcal{H} is any finite set of graphs.

This submission was selected as one of the best papers in ISAAC 2023.

✉ Bernard Ries
bernard.ries@unifr.ch

Carl Feghali
carl.feghali@ens-lyon.fr

Felicia Lucke
felicia.lucke@unifr.ch

Daniël Paulusma
daniel.paulusma@durham.ac.uk

¹ University of Lyon, EnsL, CNRS, LIP, 69342 Lyon Cedex 07, France

² Department of Informatics, University of Fribourg, Fribourg, Switzerland

³ Department of Computer Science, Durham University, Durham, UK

Keywords Matching cut · Perfect matching · Girth · H -free graph

1 Introduction

We consider classic graph problems for finding certain edge cuts, which have in common that their edges must form a matching. In order to explain this, let $G = (V, E)$ be a connected graph. A set $M \subseteq E$ is a *matching* of G if no two edges in M share an end-vertex; M is *perfect* if every vertex of G is incident to an edge of M . A set $M \subseteq E$ is an *edge cut* of G if V can be partitioned into two sets B and R , such that M consists of all the edges with one end-vertex in B and the other one in R . We say that M is a (*perfect*) *matching cut* of G if M is a (perfect) matching that is also an edge cut. We refer to Fig. 1 for some examples.

Graphs with matching cuts were introduced in 1970 by Graham [18] as *decomposable* graphs. Matching cuts have applications in number theory [18], graph drawing [34], graph homomorphisms [16], edge labelings [2] and ILFI networks [13]. Moreover, a connected graph with no vertex of degree 1 has a matching cut if and only if its line graph has a vertex cut that is an independent set (*stable cut set*). As such, (perfect) matching cuts are well studied in the literature.

Instead of considering perfect matchings that *are* edge cuts, we can also consider perfect matchings in graphs that *contain* edge cuts. Such perfect matchings are called *disconnected perfect matchings*; see Fig. 1 again. Note that every perfect matching cut is a disconnected perfect matching. However, there exist connected graphs, like the cycle C_6 on six vertices, that have a disconnected perfect matching (and thus a matching cut) but no perfect matching cut. There also exist connected graphs, like the path P_3 on three vertices, that have a matching cut, but no disconnected perfect matching (and thus no perfect matching cut either).

The problems MATCHING CUT, DISCONNECTED PERFECT MATCHING and PERFECT MATCHING are to decide if a connected graph has a matching cut, disconnected perfect matching or perfect matching cut, respectively. As explained below, all three problems are NP-complete, and have been extensively studied for special graph classes.

Our Focus. The *girth* of a graph that is not a forest is the number of edges of a shortest cycle in it; a forest has infinite girth. In 2003, Bonsma [6] asked if MATCHING CUT is still NP-complete for graphs of large girth and proved that this is indeed the case for planar graphs of girth 5. In the 2009 journal version of [6], Bonsma showed that every connected planar graph of girth at least 6 has a matching cut. Hence, the complexity status of MATCHING CUT for graphs of large girth remained unknown and was regularly posed as an open problem [9, 24, 27, 29].

Bouquet and Picouleau [8] proved that DISCONNECTED PERFECT MATCHING is NP-complete for planar graphs of girth $g = 5$ and left the cases where $g \geq 6$ open. In contrast, Le and Telle [27] proved that for every $g \geq 3$, PERFECT MATCHING CUT is NP-complete for subcubic bipartite graphs of girth at least g (a graph is *subcubic* if it has maximum degree at most 3). We focus on the two remaining open problems:

What is the complexity of MATCHING CUT and DISCONNECTED PERFECT MATCHING for graphs of large girth?



Fig. 1 The graph P_6 from [31] with a matching cut that is not contained in a disconnected perfect matching (left), a matching cut that is properly contained in a disconnected perfect matching (middle) and a perfect matching cut (right). In each figure, thick edges denote matching cut edges

A challenging task is to find gadgets with no edges that are subdivided twice; such gadgets always have a matching cut (take the two subdivision vertices on one side and all other vertices on the other) and cannot be used in any hardness reduction.

1.1 Other Relevant Known Results

We restrict ourselves to *hereditary* graph classes, that is, classes of graphs closed under vertex deletion. A class of graphs \mathcal{G} is hereditary if and only if the graphs in \mathcal{G} are $\mathcal{F}_{\mathcal{G}}$ -free for some unique set $\mathcal{F}_{\mathcal{G}}$, that is, they do not contain any graph from $\mathcal{F}_{\mathcal{G}}$ as an induced subgraph. For a *systematic* study, one may start with H -free graphs (so where $\mathcal{F}_{\mathcal{G}}$ has a single graph H).

Matching Cuts Chvátal [10] proved that MATCHING CUT is NP-complete even for $K_{1,4}$ -free graphs of maximum degree 4 (the graph $K_{1,r}$ is the $(r + 1)$ -vertex star); see [34] for an alternative hardness proof. In contrast, Chvátal [10] also proved that MATCHING CUT is polynomial-time solvable for graphs of maximum degree at most 3, whereas Bonsma [6] proved the same for $K_{1,3}$ -free graphs, thereby generalizing a known result of Moshi [32] for line graphs, and also for P_4 -free graphs; the latter result was extended to P_5 -free graphs in [14] and to P_6 -free graphs in [29]. Kratsch and Le [23] proved polynomial-time solvability for $(K_{1,4}, K_{1,4} + e)$ -free graphs. It is also known that if MATCHING CUT is polynomial-time solvable for H -free graphs for some graph H , then it is so for $(H + P_3)$ -free graphs [29]; for two vertex-disjoint graphs G_1 and G_2 , we write $G_1 + G_2 = (V(G_1) \cup V(G_2), E(G_1) \cup E(G_2))$.

Moshi [32] proved that MATCHING CUT is NP-complete even for bipartite graphs where the vertices in one set of the bipartition all have degree exactly 2. Consequently, MATCHING CUT is NP-complete for $(H_1^*, H_3^*, H_5^*, \dots)$ -free bipartite graphs, where $H_1^* = H^*$ denotes the graph that looks like the letter “H”, and for $i \geq 2$, the graph H_i^* is the graph obtained from $H^* = H_1^*$ by subdividing the middle edge of H_1^* exactly $i - 1$ times; see also Fig. 2.

Le and Randerath [26] proved that MATCHING CUT is NP-complete for $K_{1,5}$ -free bipartite graphs and thus for C_s -free graphs if s is odd. Recall that Bonsma [6] proved that MATCHING CUT is NP-complete for planar graphs of girth 5, and thus for (C_3, C_4) -free graphs. By using the graph transformation of Moshi [32], MATCHING CUT is NP-complete for C_s -free graphs also if s is even and at least 6 [29]. In [31] it was shown that MATCHING CUT is NP-complete even for $(3P_5, P_{15})$ -free graphs, strengthening a result of [14]. Afterwards, Le and Le [25] proved that MATCHING CUT is NP-complete even for $(3P_6, 2P_7, P_{14})$ -free graphs.

We refer to [7, 24, 29] for results for non-hereditary graph classes, to [3, 15, 17, 22, 23] for parameterized complexity results and exact algorithms, to [4, 17] for a generalization of MATCHING CUT to d -CUT (where instead of at most one neighbour




Fig. 2 The graphs $H^* = H_1^*$ (left) and H_i^* (right)

we allow each vertex to have at most d neighbours across the cut) and to [9] for a comprehensive overview.

Disconnected Perfect Matchings The DISCONNECTED PERFECT MATCHING problem was introduced by Bouquet and Picouleau [8]. They used a different name but we adapt the name of Le and Telle [27] to avoid confusion with PERFECT MATCHING CUT. Bouquet and Picouleau [8] showed that DISCONNECTED PERFECT MATCHING is, among others, polynomial-time solvable for $K_{1,3}$ -free graphs and P_5 -free graphs, but NP-complete for $K_{1,4}$ -free planar graphs, planar graphs of girth 5 and for bipartite graphs, and thus for C_s -free graphs for every odd s . Le and Le [25] proved that DISCONNECTED PERFECT MATCHING is NP-complete even for $(3P_6, 2P_7, P_{14})$ -free graphs, which improved the previous NP-completeness result of [31] for $(3P_7, P_{19})$ -free graphs.

Perfect Matching Cuts The PERFECT MATCHING CUT problem was first shown to be NP-complete by Heggernes and Telle [19]. Le and Telle [27] proved, besides NP-completeness for subcubic bipartite graphs of girth at least g (for any $g \geq 3$), that PERFECT MATCHING CUT is polynomial-time solvable for chordal graphs and for $S_{1,2,2}$ -free graphs; the graph $S_{1,2,2}$ is obtained by subdividing two of the edges of the claw $K_{1,3}$ exactly once. In [31], it was shown that PERFECT MATCHING CUT is polynomial-time solvable for P_6 -free graphs, and moreover for $(H + P_4)$ -free graphs if it is polynomial-time solvable for H -free graphs.

Even more recently, two new results were shown. Le and Le [25] proved that PERFECT MATCHING CUT is NP-complete even for $(3P_6, 2P_7, P_{14})$ -free graphs (using the same construction as for MATCHING CUT and DISCONNECTED PERFECT MATCHING), whereas Bonnet, Chakraborty and Duron [5] proved that PERFECT MATCHING CUT is NP-complete for 3-connected cubic bipartite planar graphs.

1.2 Our Results

As our main results, we solve, in Sect. 3, the aforementioned two open problems in the literature [6, 8, 9, 24, 27, 29] by showing that for every $g \geq 3$, MATCHING CUT is NP-complete for bipartite graphs of girth at least g and maximum degree at most 60 and DISCONNECTED PERFECT MATCHING is NP-complete for bipartite graphs of girth at least g and maximum degree at most 74. Note that our first result answers, in the negative, a question [33] from the *Open Problem Garden*:

“For every d does there exist a g such that every graph with average degree smaller than d and girth at least g has a matching-cut?”

As an immediate consequence of our second result, we have that DISCONNECTED PERFECT MATCHING is NP-complete for C_s -free graphs for even s (as mentioned above, previously this was only known for odd s [8]). To overcome the obstacle that connected graphs with a 2-subdivided edge have a matching cut, we use results from the theory of expander graphs. The proof of our result on MATCHING CUT is surprisingly short. Moreover, we can extend it in a straightforward way to show the following. For every $d \geq 2$ and $g \geq 3$, there exists a function $f(d)$ that only depends on d , such that even d -CUT is NP-complete for bipartite graphs of girth at least g and maximum degree at most $f(d)$; recall that 1-CUT is the MATCHING CUT problem.

In Sect. 3, we also highlight an implicit result of Le and Telle [27] for PERFECT MATCHING CUT. In their NP-hardness proof for subcubic bipartite graphs of high girth, they showed that a graph G has a *perfect* matching cut if and only if the graph obtained from G by subdividing some edge four times has a *perfect* matching cut. Since graphs with a 2-subdivided edge have a matching cut, this property shows a fundamental difference between matching cuts and perfect matching cuts.

For $i \geq 2$, recall from Fig. 2 that H_i^* is the graph obtained from $H^* = H_1^*$ by subdividing the middle edge of H_1^* exactly $i - 1$ times. Recall also that a result of Moshi [32] implies that MATCHING CUT is NP-complete for $(H_1^*, H_3^*, H_5^*, \dots)$ -free bipartite graphs. In Sect. 4, we extend this result by proving that for every $i \geq 1$, MATCHING CUT and DISCONNECTED PERFECT MATCHING are NP-complete for (H_1^*, \dots, H_i^*) -free graphs. We obtain these results by replacing the gadget of Moshi [32] with more advanced graph transformations. In Sect. 4, we also make explicit that the construction of Le and Telle [27] for subcubic bipartite graphs of girth at least g gives in fact NP-completeness of PERFECT MATCHING CUT for (H_1^*, \dots, H_i^*) -free subcubic bipartite graphs of girth at least g , for any $g \geq 3$. Hence, all three problems are NP-complete for H -free graphs whenever H has a connected component with two vertices of degree at least 3.

In Sect. 5, we combine our new results with the known results. We update the state-of-the-art summaries for the three problems on H -free graphs; basically, for all three problems, we only need to consider cases where H is a linear forest. Apart from comparing the (partial) classification of the three problems with each other, we also compare them with a recent complete classification of the MAXIMUM MATCHING CUT problem for H -free graphs [30]. This problem is to determine a matching cut in a graph G with a maximum number of edges (or output that no matching cut exists in G). Finally, we show how the results for PERFECT MATCHING CUT lead to a full classification of PERFECT MATCHING CUT on \mathcal{H} -subgraph-free graphs, for every finite set of graphs \mathcal{H} .

2 Preliminaries

We only consider finite, undirected graphs without multiple edges and self-loops. Let $G = (V, E)$ be a graph. For $u \in V$, the set $N(u) = \{v \in V \mid uv \in E\}$ is the

neighbourhood of u in G , where $|N(u)|$ is the *degree* of u . For an integer $p \geq 0$, G is p -regular if every $u \in V$ has degree p . Let $S \subseteq V$. The *neighbourhood* of S is the set $N(S) = \bigcup_{u \in S} N(u) \setminus S$. The graph $G[S]$ is the subgraph of G induced by $S \subseteq V$, that is, $G[S]$ is the graph obtained from G after deleting the vertices not in S . We write $G - S = G[V \setminus S]$. Let $u, v \in V$. The *distance* between u and v in G is the *length* (number of edges) of a shortest path between u and v in G . The subdivision of an edge $e = uv$ of G replaces e by a new vertex w and edges uw and wv .

We will now define some useful colouring terminology for matching cuts used in other papers as well (see e.g. [31]). In the remainder of this section, we let $G = (V, E)$ be a connected graph. A *red-blue colouring* of G colours every vertex of G either red or blue. If every vertex of some set $S \subseteq V$ has the same colour (red or blue), then S is said to be *monochromatic*. We also say that $G[S]$ is *monochromatic*. A red-blue colouring of G is *valid* if the following holds:

1. every blue vertex has at most one red neighbour;
2. every red vertex has at most one blue neighbour; and
3. both colours red and blue are used at least once.

If a red vertex u in G has a blue neighbour v , then u and v are said to be *matched*. See Fig. 1 for three examples of valid red-blue colourings of the P_6 .

For a valid red-blue colouring of G , we let R be the *red* set consisting of all vertices coloured red and B be the *blue* set consisting of all vertices coloured blue. Note that $V = R \cup B$. The *red interface* is the set $R' \subseteq R$ consisting of all vertices in R with a (unique) blue neighbour, and the *blue interface* is the set $B' \subseteq B$ consisting of all vertices in B with a (unique) red neighbour in R .

A red-blue colouring of G is *perfect* if it is valid and moreover $R' = R$ and $B' = B$; see Fig. 1 (middle) for an example of a perfect red-blue colouring (of the P_6). A red-blue colouring of G is *perfect-extendable* if it is valid and $G[R \setminus R']$ and $G[B \setminus B']$ both contain a perfect matching; see Fig. 1 (right) for an example of a perfect-extendable red-blue colouring (of the P_6). In other words, the matching defined by the edges with one end-vertex in R' and the other one in B' can be extended to a perfect matching in G or, equivalently, is contained in a perfect matching in G .

We now make the following straightforward observation (see also e.g. [31]).

Observation 1 *Let G be a connected graph. The following three statements hold:*

- (i) G has a matching cut if and only if G has a valid red-blue colouring;
- (ii) G has a disconnected perfect matching if and only if G has a perfect-extendable red-blue colouring;
- (iii) G has a perfect matching cut if and only if G has a perfect red-blue colouring.

Finally, we formally define the notion of a d -cut. For an integer $d \geq 1$ and a connected graph $G = (V, E)$, a set $M \subseteq E$ is a d -cut of G if V can be partitioned into two sets B and R , such that the following two conditions hold:

- (i) M consists of all the edges with one end-vertex in B and the other one in R ; and
- (ii) every vertex in B has at most d neighbours in R , and vice versa.

Recall that the corresponding d -CUT problem is to decide if a connected graph has a d -cut. Hence, 1-CUT and MATCHING CUT are the same problems.

3 Hardness for Arbitrary Given Girth

In this section, we will show that MATCHING CUT (Sect. 3.1) and DISCONNECTED PERFECT MATCHING (Sect. 3.2) remain NP-complete even for graphs of high girth and bounded maximum degree; recall that previously both problems were known to be NP-complete for (planar) graphs of girth at least g , only for $g \leq 5$ [6, 8]. We also briefly discuss how an implicit observation of Le and Telle [27] gives a simple, alternative proof for showing NP-completeness for PERFECT MATCHING CUT (Sect. 3.3) for graphs of high girth.

We need the following notions. The *edge expansion* $h(G)$ of a graph $G = (V, E)$ on n vertices is defined as

$$h(G) = \min_{1 \leq |S| \leq \frac{n}{2}} \frac{|\partial S|}{|S|},$$

where $\partial S := \{uv \in E \mid u \in S, v \in V \setminus S\}$ is the set of all edges of G with one end-vertex in S and one end-vertex outside S .

A connected graph G is said to be (*matching-*)*immune* if G admits no matching cut. We generalize this notion as follows. For an integer $d \geq 1$, we say that a connected graph G is *d-immune* if G admits no d -cut, so being 1-immune is the same as being immune. We make the following observation.

Observation 2 *For every integer $d \geq 1$, every connected graph G with $h(G) > d$ is d -immune.*

Proof Let $d \geq 1$, and let G be a connected graph with $h(G) > d$. For every set $S \subseteq V$ with $1 \leq |S| \leq \frac{n}{2}$, the average number of neighbours the vertices of S have in $V \setminus S$ is strictly larger than d . Hence, G cannot have a d -cut. In other words, G is d -immune. \square

3.1 Matching Cut and d-Cut

In order to prove our hardness results for MATCHING CUT and d -CUT for bipartite graphs of high girth and bounded maximum degree, we use known results on expander graphs and number theory.

Two integers a and b are *coprime*, if they do not have a common divisor greater than 1. The following result is well known.

Theorem 3 ([11]) *For two positive, coprime integers a and b , the sequence $a + bk$, for $k \in \mathbb{N}$ contains infinitely many primes.*

For an integer a and a prime p , the *Legendre symbol* is defined as $\left(\frac{a}{p}\right) \equiv a^{\frac{p-1}{2}} \pmod{p}$. It is well known (see, for example, [35]) that for any integer a and prime p , it holds that $\left(\frac{a}{p}\right) \equiv 0 \pmod{p}$, or $1 \pmod{p}$, or $(p-1) \pmod{p}$, and with slight abuse of notation one denotes these integers by 0, 1 and -1 , respectively.

We use Theorem 3 to prove the following lemma.

Lemma 4 *There are infinitely many primes q , such that*

- a) $q > 13$,
- b) $q \equiv 1 \pmod{4}$, and
- c) the Legendre symbol $\left(\frac{q}{13}\right) = -1$.

Proof We apply Theorem 3 for $a = 5$ and $b = 52$, which are coprime. This yields that the sequence $5 + 52k$, for $k \in \mathbb{N}$, contains infinitely many primes, and consequently, infinitely many primes satisfying a). Since for any $k \in \mathbb{N}$, we have $5 + 52k \equiv 1 \pmod{4}$, condition b) is also satisfied.

Finally, we will show that the Legendre symbol $\left(\frac{5+52k}{13}\right) = -1$. First we note that if $a \equiv b \pmod{p}$, then $\left(\frac{a}{p}\right) = \left(\frac{b}{p}\right)$ (see [35]). Hence, since $5 \equiv 5 + 52k \pmod{13}$, we have $\left(\frac{5}{13}\right) = \left(\frac{5+52k}{13}\right)$. We now deduce that $\left(\frac{5}{13}\right) \equiv 5^{\frac{13-1}{2}} \pmod{13} \equiv 12 \pmod{13} \equiv -1 \pmod{13}$. This proves condition c). \square

Lemma 5 uses known results from the theory of expander graphs. We need the fact that the graph in the statement of Lemma 5 has a perfect matching only in Sect. 3.2.

Lemma 5 *For every $g \geq 3$, there is an immune 14-regular bipartite graph with girth at least g that contains a perfect matching.*

Proof It follows from a construction of Lubotzky, Phillips and Sarnak [28] based on Cayley graphs that for every two primes p and q with the following four properties

- $q > p$,
- $p \equiv 1 \pmod{4}$,
- $q \equiv 1 \pmod{4}$, and
- $\left(\frac{p}{q}\right) = -1$,

there exists a $(p+1)$ -regular bipartite graph G with $\lambda_2(G) \leq 2\sqrt{p}$ and girth at least $4 \log_p q - \log_p 4$; here, $\lambda_2(G)$ denotes the second largest eigenvalue of the adjacency matrix of G .

We set $p = 13$, so $p \equiv 1 \pmod{4}$. We now combine the above with Lemma 4 to find that there exist infinitely many primes q , such that there exists a 14-regular bipartite graph G_q with $\lambda_2(G_q) \leq 2\sqrt{13}$ and girth at least $4 \log_{13} q - \log_{13} 4$. Moreover, Dodziuk [12] and, independently, Alon and Milman [1] showed that for every integer $\ell \geq 1$, every ℓ -regular graph G satisfies $h(G) \geq \frac{1}{2}(\ell - \lambda_2(G))$. This means that

$$h(G_q) \geq \frac{1}{2}(14 - \lambda_2(G_q)) \geq \frac{1}{2}(14 - 2\sqrt{13}) \approx 3.39 > 1.$$

Hence, by applying Observation 2, we find that G_q is immune.

By taking q sufficiently large, we conclude that for any $g \geq 3$, there exists an immune 14-regular bipartite graph G with girth at least g . Finally, as G is bipartite and regular, we find that G has a perfect matching (due to Hall's Marriage Theorem). \square

Remark 1 The arguments that we used to prove Lemma 5 do not allow us to set $p = 13$ to a smaller value. We need p to be prime, and moreover, it must hold that




Fig. 3 The graph F with designated vertices x and y at distance at least g (left) and the graph F' (right) from the proof of Theorem 6

$p \equiv 1 \pmod{4}$. Hence, the only alternative value for p that is smaller than 13 would be $p = 5$. However, $p = 5$ yields $h(G_q) \geq \frac{1}{2}(6 - 2\sqrt{5}) \approx 0.76$, so we cannot conclude from this that G_q is immune.

We use Lemma 5 in the proof of our first main result.

Theorem 6 *For every integer $g \geq 3$, MATCHING CUT is NP-complete for bipartite graphs of girth at least g and maximum degree at most 60.*

Proof Let $g \geq 3$. As the class of graphs of girth at least $g + 1$ is a subclass of the class of graphs of girth at least g , we may assume without loss of generality that g is divisible by 2 but not by 4, so $\frac{g}{2}$ is odd. We reduce from MATCHING CUT. Recall that MATCHING CUT is NP-complete for graphs of degree at most 4 [10]. Let G be a connected graph of maximum degree at most 4. From G we construct a graph G' with the required properties, but first we define an auxiliary graph F' .

By Lemma 5 there exists an immune 14-regular bipartite graph F with girth at least g . Note that F has constant size, so we can find F in constant time.

Let x and y be two vertices of distance $\frac{g}{2}$ in F . As $\frac{g}{2}$ is odd, x and y belong to opposite bipartition classes of F . We take two copies F_1 and F_2 of F , and we add an edge between the two copies x_1 and x_2 of x and an edge between the two copies y_1 and y_2 of y . This yields a graph F' . As F is bipartite and has girth at least g , we find that F' is bipartite and has girth at least g as well. The construction also gives us that x_1 and y_2 belong to the same bipartition class of F' . See also Fig. 6.

Now consider an edge uv in G . The F' -replacement of uv is obtained from the graphs G and F' by identifying u with x_1 and v with y_2 . We do an F' -replacement on every edge of G . This yields the graph G' . Since F' is bipartite such that the distance between x_1 and y_2 is even, we find that G' is bipartite as well. By construction, G' has girth at least g and maximum degree at most $4 \times (14 + 1) = 60$.

We claim that G has a matching cut if and only if G' has a matching cut. We prove this below, using Observation 1-(i) implicitly.

First suppose that G has a matching cut, so G has a valid red-blue colouring c . For every edge uv in G we do as follows. Let F' with copies F_1 and F_2 of F be the corresponding F' -replacement applied on uv . If u and v have the same colour, then we colour every vertex of $V(F') \setminus \{u, v\}$ with that colour. If u and v are coloured differently, say u is red and v is blue, then we colour every vertex in $V(F_1)$ red and every vertex in $V(F_2)$ blue. This yields a valid red-blue colouring of G' . Hence, G' has a matching cut.

Now suppose that G' has a matching cut, so G' has a valid red-blue colouring c' . As F is immune, every copy of it in G' is monochromatic. Thus, for two vertices

$x_1, y_2 \in V(F')$ in some graph F' in G' , it holds that x_1 and y_2 each have a neighbour in F' of the opposite colour if and only if x_1 and y_2 have different colours in G' . Hence, the restriction of c' to $V(G)$ is a valid red-blue colouring of G . Hence, G has a matching cut. \square

We now focus on the d - CUT problem for arbitrary $d \geq 1$, and we show how Theorem 6 can be generalized in a straightforward way. This requires us to replace Lemma 4 by the following lemma.

Lemma 7 *There are infinitely many primes p , such that*

- a) $p \geq 13$,
- b) $p \equiv 1 \pmod{4}$, and
- c) *there exists an infinite set Q_p such that the following holds for every $q \in Q_p$:*
 - i. $q > p$
 - ii. $q \equiv 1 \pmod{4}$, and
 - iii. *the Legendre symbol $\left(\frac{q}{p}\right) = -1$.*

Proof We first apply Theorem 3 to $a = 13$ and $b = 20$, which are coprime. In this way we find that the sequence $13 + 20k_1$, for $k_1 \in \mathbb{N}$, contains infinitely many primes $p \geq 13$. Hence, condition a) is satisfied. Since $p = 13 + 20k_1$ for some integer $k_1 \geq 0$, we find that $p \equiv 1 \pmod{4}$. Hence, condition b) is satisfied as well.

We now show condition c). We apply Theorem 3 to $a = 5$ and $b = 4p$. In this way we find that the sequence $5 + 4pk_2$, for $k_2 \in \mathbb{N}$, contains infinitely many primes q , so in particular an infinite set Q_p with $q > p$ for every $q \in Q_p$. This shows that condition c)i is satisfied. For every $q \in Q_p$ it holds that $q = 5 + 4pk_2$ for some $k_2 \geq 0$ and thus $q \equiv 1 \pmod{4}$. Hence, condition c)ii is satisfied as well.

Finally, we will show condition c)iii by proving that $\left(\frac{q}{p}\right) = -1$ holds for every $q \in Q_p$. First, we recall that if $a \equiv b \pmod{c}$, then $\left(\frac{a}{c}\right) = \left(\frac{b}{c}\right)$ (see [35]). We will also use the law of reciprocity, of which several equivalent versions exist in the literature. We use the following partial version (which can be found in [35]). For every two primes $a > 2$ and $b > 2$, it holds that

$$\left(\frac{a}{b}\right) = \left(\frac{b}{a}\right) \text{ if } a \equiv 1 \pmod{4} \text{ or } b \equiv 1 \pmod{4}$$

Combining both properties of the Legendre symbol we can now deduce that

$$\begin{aligned} \left(\frac{q}{p}\right) &= \left(\frac{5 + 4pk_2}{13 + 20k_1}\right) \\ &= \left(\frac{5 + 4k_2(13 + 20k_1)}{13 + 20k_1}\right) \\ &= \left(\frac{5}{13 + 20k_1}\right) \\ &= \left(\frac{13 + 20k_1}{5}\right) \end{aligned}$$

$$= \left(\frac{3}{5}\right) \\ \equiv -1 \pmod{3}.$$

Thus, condition c)iii is satisfied as well, and we completed the proof of the lemma. \square

We are now ready to generalize Theorem 6.

Theorem 8 *For every integer $d \geq 1$ and every integer $g \geq 3$, there is a function $f(d)$ that only depends on d , such that d -CUT is NP-complete for bipartite graphs of girth at least g and maximum degree at most $f(d)$.*

Proof Let $d \geq 1$. By Observation 2, every graph G with $h(G) > d$ is d -immune. Hence, we can construct a $(p + 1)$ -regular bipartite gadget F using the arguments from the proof of Lemma 5, where Lemma 7 plays the role of Lemma 4. That is, we can choose p such that

- (i) p is a prime congruent to 1 mod 4, and
- (ii) $\frac{1}{2}(p + 1 - \lambda_2(G)) \geq \frac{1}{2}(p + 1 - 2\sqrt{p}) > d$.

We now reduce from d -CUT. Gomes and Sau [17] proved that d -CUT is NP-hard for graphs of maximum degree at most $2d + 2$. Hence, we may assume that the instance G from d -CUT has maximum degree at most $2d + 2$. By applying the arguments of the proof of Theorem 6, we can use F and G to construct for every integer $g \geq 3$, a bipartite graph G' of girth at least g and maximum degree at most $f(d)$, such that G has a d -cut if and only if G' has a d -cut. \square

Remark 2 Since it is not possible to give the smallest prime p satisfying conditions (i) and (ii) in the proof of Theorem 8, we can merely state that the maximum degree of G' is bounded by some function $f(d)$ that only depends on d .

3.2 Disconnected Perfect Matching

We now show that DISCONNECTED PERFECT MATCHING is NP-complete for graphs of arbitrarily large fixed girth and bounded maximum degree. Our proof uses some new ideas, but we note that it is also possible to use a similar approach as in the proof of Theorem 6. However, this will lead to a slightly worse bound on the maximum degree, namely 75 instead of 74.

We first define some useful auxiliary graphs. Fix $g \geq 3$ such that g is divisible by 6 and $g/6$ is odd. Let F be a 14-regular immune bipartite graph with girth at least $2(g + 1)$ containing a perfect matching. We note that F exists by Lemma 5, and as F has constant size, F can be found in constant time.

Let s and t be two designated vertices in F at distance at least $g + 1$. We fix a perfect matching M of F . Let $x \in N_F(s)$ and $y \in N_F(t)$ be the (unique) neighbours of s and t in M . Then, since s and t are at distance at least $g + 1$, x and y are at distance at least $g - 1$. We add the edge xy and denote the resulting graph by $F(s, t)$; see also Fig. 4a.

We make the following observation.




Fig. 4 **a** Illustration of the graphs F (left), $F(s, t)$ (middle) and $H(s, t)$ (right). **b** Illustration of how the edges in the perfect matching of $H(s, t)$ (left) resp. $H(s, t) - \{s, t\}$ (right) are chosen (these edges are represented as red, thick edges)

Lemma 9 *The graph $F(s, t)$ is immune, bipartite and has girth at least g . Moreover, both $F(s, t)$ and $F(s, t) - \{s, t\}$ contain a perfect matching.*

Proof As F is immune, $F(s, t)$ is immune. By our choice of x and y , we find that $F(s, t)$ is bipartite and has girth at least g . The fixed perfect matching M of F is of course also a perfect matching of $F(s, t)$. The new edge xy ensures that $F(s, t) - \{s, t\}$ contains a perfect matching as well; indeed, we can take $(M \setminus \{sx, ty\}) \cup \{xy\}$. \square

We now take $k = \frac{g}{6}$ copies $F(s_1, t_1), \dots, F(s_k, t_k)$ of $F(s, t)$ and identify s_{i+1} with t_i for all $i \in \{1, \dots, k-1\}$. We set $s = s_1$ and $t = t_k$ and call the resulting graph $H(s, t)$; see also Fig. 4a.

In the following lemma we show some useful properties of $H(s, t)$ and $H(s, t) - \{s, t\}$.

Lemma 10 *The graph $H(s, t)$ is immune, bipartite, and has girth at least g . Moreover, $\text{dist}(s, t) \geq \frac{g}{2}$, and both $H(s, t)$ and $H(s, t) - \{s, t\}$ contain a perfect matching.*

Proof By taking two immune graphs and identifying two of their vertices (one from each graph), we obtain another immune graph. The graph $H(s, t)$ is obtained by repeatedly applying this operation on copies of $F(s, t)$, which is immune by Lemma 9. Hence, we conclude that $H(s, t)$ is immune.

By Lemma 9, every graph $F(s_i, t_i)$ for $i \in \{1, \dots, k\}$ is bipartite and has girth at least g . By constructing $H(s, t)$ from these graphs $F(s_i, t_i)$, we do not create any new cycles. Thus, $H(s, t)$ also is bipartite and has girth at least g . In every graph $F(s_i, t_i)$ we have $\text{dist}(s_i, t_i) = 3$, thus $\text{dist}(s, t) = 3 \times \frac{g}{6} = \frac{g}{2}$.

To see that $H(s, t)$ and $H(s, t) - \{s, t\}$ contain both a perfect matching, we will use the fact that both $F(s_i, t_i)$ and $F(s_i, t_i) - \{s_i, t_i\}$ contain a perfect matching by Lemma 9. For $H(s, t)$, we take a perfect matching of $F(s_1, t_1)$, $F(s_2, t_2) - \{s_2, t_2\}$, $F(s_3, t_3)$ and continue this alternation until $F(s_k, t_k)$, see Fig. 4b (left). For $H(s, t) - \{s, t\}$ we alternate as well but we start and end with $F(s_1, t_1) - \{s_1, t_1\}$ and $F(s_k, t_k) - \{s_k, t_k\}$, see Fig. 4b (right). \square

We are now ready to prove the following result.

Theorem 11 *For every integer $g \geq 3$, DISCONNECTED PERFECT MATCHING is NP-complete for bipartite graphs of girth at least g and maximum degree at most 74.*

Proof Let $g \geq 3$. We reduce from MATCHING CUT for bipartite graphs of girth at least g and maximum degree at most 60, which is NP-complete by Theorem 6. Similar to Theorem 6, we may assume without loss of generality that g is divisible by 12. Let G be a bipartite graph of girth at least g and maximum degree 60. We construct a graph G' by taking two copies G_1 and G_2 of G , where we connect every vertex $v \in V(G_1)$ and its copy $v' \in V(G_2)$ using the graph $H(v, v')$.

To see that G' has girth at least g , we consider first G_1 and G_2 , which both have girth at least g . Any cycle containing vertices from both copies has to pass twice through a graph $H(s, t)$. Thus, it will always have length at least g , and so G' has girth g . Every vertex inside $H(s, t)$ has degree at most 28, whereas s and t only have degree 14. The degree of a vertex $v \in V(G_1) \cup V(G_2)$ is the degree of the vertex in the original graph G plus the degree in the graph $H(v, v')$. Thus, the degree of v is at most 74.

We also claim that G' is bipartite. For a contradiction, assume that G' has an odd cycle C . As G_1 and G_2 are both bipartite, C must contain vertices from G_1 and G_2 . Note that C passes through an even number of graphs $H(v, v')$, where $v \in V(G_1)$ and $v' \in V(G_2)$. Hence, the number of edges in $E(G) \setminus (E(G_1) \cup E(G_2))$ is even. Since the graph $H(v, v')$ always connects a vertex v in G_1 and its copy v' in G_2 we can find an odd cycle C' in G_1 , consisting of the edges in $C \cap E(G_1)$ and the edges in $E(G_1)$ corresponding to the edges from $C \cap E(G_2)$, a contradiction.

Finally, we show that G admits a matching cut if and only if G' admits a disconnected perfect matching. Consider some vertex $v \in V(G_1)$ and its copy $v' \in V(G_2)$. Since v and v' are connected by the graph $H(v, v')$, which is immune, v and v' will always have the same colour in any valid red-blue colouring of G' . Thus, G_1 and G_2 will be coloured the same in any valid red-blue colouring. Now, if G admits a perfect-extendable red-blue colouring, then G admits a valid red-blue colouring, as it suffices to colour G the same as G_1 (or G_2).

Conversely, if G admits a valid red-blue colouring, then we obtain a perfect-extendable colouring of G' as follows. We colour G_1 and G_2 the same as G . Notice that we colour the immune graphs connecting two copies of the same vertex such that they are monochromatic. This gives us a valid red-blue colouring of G' , i.e. a matching cut M in G' . It remains to show that the matching cut is contained in a perfect matching of G' . Since the colourings of G_1 and G_2 are the same, we have that whenever a blue vertex $v \in V(G_1)$ is matched with a red vertex $u \in V(G_1)$, i.e. $vu \in M$, then their copies $v' \in V(G_2)$ and $u' \in V(G_2)$ are matched as well, i.e. $v'u' \in M$. By Lemma 10, we know that $H(v, v') - \{v, v'\}$ and $H(u, u') - \{u, u'\}$ contain both a perfect matching which we may add to M . For every $v \in V(G_1)$ with no neighbour of the other colour, we know that its copy $v' \in V(G_2)$ has no neighbour of the other colour either. Thus, we can use that $H(v, v')$ contains a perfect matching by Lemma 10 and add it to M . Repeatedly doing this yields a perfect matching of G' containing M . \square

3.3 Perfect Matching Cut

In any perfect red-blue colouring of a connected graph $G = (V, E)$, a vertex $v \in V$ of degree 2 has exactly one neighbour coloured the same as itself and exactly one neighbour coloured differently than itself. One can use this observation to prove the

following lemma. This lemma was implicit in [27]. Its proof is short, and we include it for completeness.

Lemma 12 ([27]) *Let $G = (V', E')$ be the graph obtained from a connected graph G by 4-subdividing an edge e of G . Now, G has a perfect matching cut if and only if G' has a perfect matching cut.*

Proof Let $e = uv$ and denote the resulting path in G' by $uu_1u_2u_3u_4v$. First suppose G has a perfect matching cut, so G has a perfect red-blue colouring c . If u and v are coloured alike, say u and v are red, then we colour u_1 red, u_2 blue, u_3 blue and u_4 red. Else we may assume that u is red and v is blue. In that case we colour u_1 blue, u_2 blue, u_3 red and u_4 red. In both cases we obtain a perfect red-blue colouring c' of G' . So G' has a perfect matching cut.

Now suppose that G' has a perfect matching cut, so G' has a perfect red-blue colouring c' . If u and v are coloured alike, say u and v are red, then u_1 must be red, u_2 blue, u_3 blue and u_4 red. Hence, the restriction of c' to V is a perfect red-blue colouring of G . Else we may assume that u is red and v is blue. In that case u_1 must be blue, u_2 blue, u_3 red and u_4 red. Again, the restriction of c' to V is a perfect red-blue colouring of G . Hence, in both cases we find that G has a perfect matching cut. \square

A simple proof for showing that PERFECT MATCHING CUT is NP-complete for graphs of girth at least g is to apply Lemma 12, say, g times on each edge of the input graph. Recall that the more involved gadget of Le and Telle [27] yields NP-completeness even for subcubic bipartite graphs of girth at least g .

4 Hardness for Forbidden Subdivided H-Graphs

We show that MATCHING CUT (Sect. 4.1), DISCONNECTED PERFECT MATCHING (Sect. 4.2) and PERFECT MATCHING CUT (Sect. 4.3) are NP-complete for (H_1^*, \dots, H_i^*) -free graphs, for every $i \geq 1$.

4.1 Matching Cut

Let uv be an edge of a graph G . We define an edge operation as displayed in Fig. 5, which when applied on uv will replace uv in G by the subgraph T_{uv}^i . Note that in the new graph, the only vertices from T_{uv}^i that may have neighbours outside T_{uv}^i are u and v .

Theorem 13 *For all $i \geq 1$, MATCHING CUT is NP-complete for (H_1^*, \dots, H_i^*) -free graphs.*

Proof Fix $i \geq 1$. Reduce from MATCHING CUT. Let $G = (V, E)$ be a connected graph. Replace every $uv \in E$ by the graph T_{uv}^i (see Fig. 5). This yields the graph $G' = (V', E')$.

We claim that G' is (H_1^*, \dots, H_i^*) -free. For a contradiction, assume that G' contains an induced $H_{i'}^*$ for some $1 \leq i' \leq i$. Then G' contains two vertices x and y that are




Fig. 5 The edge uv (left) which we replace by the subgraph T_{uv}^i (right)

centers of an induced claw, as well as an induced path from x to y of length i' . All vertices in $V' \setminus V$ are not centers of any induced claw. Hence, x and y belong to V . By construction, any shortest path between two vertices of V has length at least $i + 1$ in G' , a contradiction.

We claim that G' has a matching cut if and only if G has a matching cut. First suppose G' has a matching cut M' , so G' has a valid red-blue colouring c' . We prove a claim for G' :

Claim 13.1 For every edge $uv \in E(G)$ it holds that

- (a) either $c'(u) = c'(v)$, and then T_{uv}^i is monochromatic, or
- (b) $c'(u) \neq c'(v)$, and then c' colours u_1, \dots, u_{2i} with the same colour as u , while c' colours all vertices of $T_{uv}^i - \{u, u_1, \dots, u_{2i}\}$ with the same colour as v , and moreover, $uv_{2i}, vu_{2i} \in M'$.

Proof First assume $c'(u) = c'(v)$, say c' colours u and v red. As any clique of size at least 3 is monochromatic, all vertices in T_{uv}^i are coloured red, so T_{uv}^i is monochromatic.

Now assume $c'(u) \neq c'(v)$, say u is red and v blue. As before, we find that all vertices u_1, \dots, u_{2i} have the same colour as u , so are red, while all vertices v_1, \dots, v_{2i} have the same colour as v , so are blue. By definition, every edge xy with $c'(x) \neq c'(y)$ belongs to M' , so $uv_{2i}, vu_{2i} \in M'$.

We construct a subset $M \subseteq E$ in G as follows. We add an edge $uv \in E$ to M if and only if $c'(u) \neq c'(v)$ in G' . We now show that M is a matching in G . Let $u \in V$. For a contradiction, suppose that M contains edges uv and uw for $v \neq w$. Then $c'(u) \neq c'(v)$ and $c'(u) \neq c'(w)$. By Claim 13.1, we find that M' matches u in G' to vertices in T_{uv}^i and T_{uw}^i , contradicting our assumption that M' is a matching (cut). Hence, M is a matching.

Now let c be the restriction of c' to V . If c colours every vertex of G with one colour, say red, then c' would also colour every vertex of G' red by Claim 13.1, contradicting the validity of c' . Hence, c uses both colours. Moreover, for every $uv \in E$, the following holds: if $c(u) \neq c(v)$, then $c'(u) \neq c'(v)$ and thus $uv \in M$. Hence, as M is a matching, c is valid, and thus M is a matching cut of G .




Fig. 6 The edge uv (left) which we replace by the subgraph G_{uv}^i (right)

Conversely, assume that G admits a matching cut, so V has a valid red-blue colouring c . We construct a red-blue colouring c' of V' as follows.

- For every edge $uv \in E$ with $c(u) = c(v)$, we let $c'(x) = c(u)$ for every $x \in V(T_{uv}^i)$.
- For every edge $uv \in E$ with $c(u) \neq c(v)$, we let $c'(u) = c'(u_1) = \dots = c'(u_{2i}) = c(u)$ and $c'(v) = c'(v_1) = \dots = c'(v_{2i}) = c(v)$.

As c is valid, c uses both colours and thus by construction, c' uses both colours. Let $u \in V$. Again as c is valid, $c(u) \neq c(v)$ holds for at most one neighbour v of u in G . Hence, by construction, u belongs to at most one non-monochromatic gadget T_{uv}^i . Thus, c' colours in G' at most one neighbour of u with a different colour than u . Let $u \in V' \setminus V$. By construction, we find again that c' colours at most one neighbour of u with a different colour than u . Hence, c' is valid, and so G' has a matching cut. This completes the proof of Theorem 13. \square

4.2 Disconnected Perfect Matching

We show our next result in a similar way as Theorem 13. First we define the edge operation displayed in Fig. 6. This operation replaces an edge uv in a graph G by the subgraph G_{uv}^i for some integer $i \geq 1$. Note that in the resulting graph, the only vertices from G_{uv}^i that may have neighbours outside G_{uv}^i are u and v .

Theorem 14 *For every $i \geq 1$, DISCONNECTED PERFECT MATCHING is NP-complete for (H_1^*, \dots, H_i^*) -free graphs.*

Proof Fix an integer $i \geq 1$. As the class of (H_1^*, \dots, H_i^*) -free graphs is contained in the class of $(H_1^*, \dots, H_{i-1}^*)$ -free graphs if $i \geq 2$, we may assume without loss of generality that i is even (we need this assumption at a later place in our proof). We reduce from DISCONNECTED PERFECT MATCHING itself. Let $G = (V, E)$ be a connected graph.

Let uv be an edge of G . We recall the edge operation displayed in Fig. 6, which when applied on uv replaces uv by the subgraph G_{uv}^i . Recall that in the resulting graph, u and v are the only vertices from G_{uv}^i that may have neighbours outside G_{uv}^i .

In G we now replace every edge $uv \in E$ by the graph G_{uv}^i . Let $G' = (V', E')$ be the resulting graph.

We claim that G' is (H_1^*, \dots, H_i^*) -free. For a contradiction, assume that G' contains an induced $H_{i'}^*$ for some $1 \leq i' \leq i$. Then G' contains two vertices x and y that are centers of an induced claw, as well as an induced path from x to y of length i' . The only vertices in $V' \setminus V$ that are the center of an induced claw are the vertices v_{2i+2} (see Fig. 6). Suppose for a contradiction that $x = v_{2i+2}$. Since the graph $G[\{u, u', v_{2i+2}, v_{2i+1}, v_{i+1}\}]$ contains an induced C_4 , the induced path from x to y may not contain u . Thus, either $y = v$ or the path from x to y passes through v . Since a shortest path from v_{2i+2} to v has length $i + 1$, we get a contradiction. Hence, x and y belong to V . By construction, any shortest path between two vertices of V has length at least $i + 3$ in G' , a contradiction.

We claim that G' has a disconnected perfect matching if and only if G has a disconnected perfect matching. First assume that G' has a disconnected perfect matching M' , so V' has a perfect-extendable red-blue colouring c' . We prove the following claim for G' :

Claim 14.1 For every edge $uv \in E(G)$, it holds that

- (a) either $c'(u) = c'(v)$, and then G_{uv}^i is monochromatic, or
- (b) $c'(u) \neq c'(v)$, and then c' colours u' and all neighbours of u in $G - v$ with the same colour as u , while c' colours all vertices of $G_{uv}^i - \{u', u\}$ with the same colour as v , and moreover, $uv_{i+1} \in M'$ and either $vv' \in M'$ or $vv_{i+2} \in M'$.

Proof First assume $c'(u) = c'(v)$, say c' colours u and v red. As any clique of size at least 3 is monochromatic, all vertices v', v_1, \dots, v_{2i+2} are coloured the same as v , so they are red. Now u' has two red neighbours, and thus u' must be red as well. Hence, G_{uv}^i is monochromatic.

Now assume $c'(u) \neq c'(v)$, say u is red and v blue. As before, we find that v', v_1, \dots, v_{2i+2} are all coloured the same as v , so they are blue. Now u , which is red, has one blue neighbour, namely v_{i+1} , and thus, u' must be red. So u and u' have the same colour. By definition, every non-monochromatic edge belongs to M' . Hence, $uv_{i+1} \in M'$. Recall that M' is perfect and observe that $|V(G_{uv}^i)|$ is even. As the vertices of $G_{uv}^i - \{u, v\}$ have no neighbours outside $G_{uv}^i - \{u, v\}$, these two facts imply that either $vv' \in M'$ or $vv_{i+2} \in M'$. If $c'(u) \neq c'(w)$ for some neighbour w of u in $G - v$ then, as we just argued, M' matches u with a vertex of G_{uw}^i , a contradiction, as M' is a matching in G' . This proves Claim 14.1.

We now construct a subset $M \subseteq E$ in G by the following rule: add an edge $uv \in E$ to M if and only if in G' , either $c'(u) = c'(v)$ and M' matches both u and v with vertices from G_{uv}^i , or $c'(u) \neq c'(v)$.

We now show that M is a perfect matching in G . Let $u \in V$. First suppose u belongs in G' to a non-monochromatic graph G_{uv}^i . By Claim 14.1, we have $c'(u) \neq c'(v)$ and $c'(u) = c'(w)$ for every neighbour w of u in $G - v$, and moreover, M' matches both u and v with vertices from G_{uv}^i . Hence, M matches u to v . As $c'(u) = c'(w)$ for every neighbour w of u in $G - v$, each G_{uw}^i is monochromatic. However, as M' already matched u to some vertex from G_{uv}^i , we find that M' (being a matching) does not match u to any vertices of any G_{uw}^i with $w \neq v$. Hence, our rule does not add any edges uw with $w \neq v$ to M . So M matches u only to v .




Fig. 7 The different colourings and matchings in the graph G_{uv}^i

Now suppose that u only belongs to monochromatic graphs G_{uv}^i . By Claim 14.1, we have $c'(u) = c'(v)$ for every neighbour v of u in G . As M' is a perfect matching, u has exactly one neighbour v in G , such that M' matches u with a vertex of G_{uv}^i . As $|V(G_{uv}^i)|$ is even and the vertices of $G_{uv}^i - \{u, v\}$ have no neighbours outside $G_{uv}^i - \{u, v\}$, this means that M' also matches v with a vertex from G_{uv}^i . Hence, M matches u to v and to no other vertex of G . We conclude that M is indeed a perfect matching in G .

Finally, let c be the restriction of c' to V . If c colours every vertex of G with one colour, say red, then c' would also colour every vertex of G' red by Claim 14.1, contradicting the perfect-extendability (and thus validity) of c' . Hence, c uses both colours. For every $uv \in E$, the following holds: if $c(u) \neq c(v)$, then $c'(u) \neq c'(v)$ and thus $uv \in M$. Hence, as M is a (perfect) matching, c is perfect-extendable. Thus M is a disconnected perfect matching of G .

Now, assume that G has a disconnected perfect matching M , so V has a perfect-extendable red-blue colouring c . We construct a red-blue colouring c' of V' and a matching M' in G' as follows:

- For every edge $uv \in E$ with $c(u) = c(v)$, we let $c'(x) = c(u)$ for every $x \in V(G_{uv}^i)$.
 - If $uv \in M$, then we add vv' , v_1v_{i+2} , \dots , $v_{i+1}v_{2i+2}$, uu' to M' ; see also Fig. 7a.
 - If $uv \notin M$, then we add $v'v_1$, v_2v_3 , \dots , $v_{2i}v_{2i+1}$, $v_{2i+2}u'$ to M' (recall that i is even, so this is possible); see also Fig. 7b.
- For every edge $uv \in E$ with $c(u) \neq c(v)$, we let $c'(u) = c'(u') = c(u)$ and $c'(x) = c(v)$ for every $x \in V(G_{uv}^i) \setminus \{u, u'\}$. We add uv_{i+1} , $u'v_{2i+2}$ to M' as well as vv' , v_1v_{i+2} , \dots , v_iv_{2i+1} ; see also Fig. 7c.

As c is valid, c uses both colours and thus by construction, c' uses both colours. Let $u \in V$. Again as c is valid, $c(u) \neq c(v)$ holds for at most one neighbour v of u in G . Hence, by construction, u belongs to at most one non-monochromatic gadget G_{uv}^i . Thus, c' colours in G' at most one neighbour of u with a different colour than u . Let $u \in V' \setminus V$. By construction, we find again that c' colours at most one neighbour of u with a different colour than u . Hence, c' is valid. Again by construction, M' is a perfect matching containing all edges xy of G' with $c'(x) \neq c'(y)$, and thus, M' is a disconnected perfect matching of G' . This completes the proof of Theorem 14. \square

4.3 Perfect Matching Cut

We apply Lemma 12 (implicit in [27]) sufficiently times on every edge of a subcubic bipartite graph of girth at least g and combine this with the NP-completeness of PERFECT MATCHING CUT for the class of subcubic bipartite graph of girth at least g [27]. Note that applying Lemma 12 does not change the degree of a graph. Hence, Le and Telle essentially proved the following:

Theorem 15 ([27]) *For every $i \geq 1$ and $g \geq 3$, PERFECT MATCHING CUT is NP-complete for (H_1^*, \dots, H_i^*) -free subcubic bipartite graphs of girth at least g .*

5 Consequences and Open Problems

We give some consequences of our new results on MATCHING CUT, DISCONNECTED PERFECT MATCHING and PERFECT MATCHING CUT for H -free graphs and H -subgraph-free graphs.

5.1 H-Free Graphs

We give three up-to-date classifications for H -free graphs by combining the results from [6, 8, 10, 25–27, 29, 31, 32] (see Sect. 1.1) with our new results. That is, we took the three state-of-the-art theorems in [31] and added both the result for H_i^* -free graphs and the result for $(3P_6, 2P_7, P_{14})$ -free graphs from [25]. For DISCONNECTED PERFECT MATCHING we also added the new result for C_s -free graphs for even s , as implied by our girth result. We also compare the three partial classification with a recent, full classification of the optimization problem MAXIMUM MATCHING CUT [30]. We write $G' \subseteq_i G$ if G' is an induced subgraph of G and $G' \supseteq_i G$ if G is an induced subgraph of G' .

Theorem 16 *For a graph H , MATCHING CUT on H -free graphs is*

- polynomial-time solvable if $H \subseteq_i sP_3 + K_{1,3}$ or $sP_3 + P_6$ for some $s \geq 0$, and
- NP-complete if $H \supseteq_i K_{1,4}, P_{14}, 3P_5, 2P_7, C_r$ for some $r \geq 3$ or H_j^* for some $j \geq 1$.

Theorem 17 *For a graph H , DISCONNECTED PERFECT MATCHING on H -free graphs is*

- polynomial-time solvable if $H \subseteq_i K_{1,3}$ or P_5 , and
- NP-complete if $H \supseteq_i K_{1,4}, P_{14}, 3P_6, 2P_7, C_r$ for some $r \geq 3$ or H_j^* for some $j \geq 1$.

Theorem 18 *For a graph H , PERFECT MATCHING CUT on H -free graphs is*

- polynomial-time solvable if $H \subseteq_i sP_4 + S_{1,2,2}$ or $sP_4 + P_6$ for some $s \geq 0$, and
- NP-complete if $H \supseteq_i K_{1,4}, P_{14}, 3P_6, 2P_7, C_r$ for some $r \geq 3$ or H_j^* for some $j \geq 1$.

Theorem 19 *For a graph H , MAXIMUM MATCHING CUT on H -free graphs is*

- *polynomial-time solvable if $H \subseteq_i sP_2 + P_6$ for some $s \geq 0$, and*
- *NP-hard if $H \supseteq_i K_{1,3}, 2P_3, C_r$ for some $r \geq 3$.*

A *subdivided claw* is a graph obtained from the claw $K_{1,3}$ by subdividing each of its edges zero or more times. Let \mathcal{S} be the class of graphs, each connected component of which is either a path or a *subdivided claw*. From Theorem 16, it follows that MATCHING CUT is NP-complete for H -free graphs if H has a cycle, a vertex of degree at least 4, or a connected component with two vertices of degree 3. Hence, the remaining open cases for MATCHING CUT on H -free graphs are all restricted to cases where H is a graph from \mathcal{S} . The same remark holds for DISCONNECTED PERFECT MATCHING due to Theorem 17, and for PERFECT MATCHING CUT due to Theorem 18.

5.2 H -subgraph-free Graphs

For a graph H , a graph G is *H -subgraph-free* if G does not contain H as a subgraph. Every H -subgraph-free graph is H -free, whereas the reverse direction only holds if H is a complete graph. For a set \mathcal{H} of graphs, a graph G is *\mathcal{H} -subgraph-free* if G is H -subgraph-free for every $H \in \mathcal{H}$.

For an integer p , a *p -subdivision* of an edge uv in a graph replaces uv by a path from u to v of length $p + 1$. The *p -subdivision* of a graph G is obtained from G by p -subdividing each edge of G . For a graph class \mathcal{G} , we let \mathcal{G}^p consist of all the p -subdivisions of the graphs in \mathcal{G} . A graph problem Π is NP-complete *under edge subdivision of subcubic graphs* if there is an integer $q \geq 1$ such that the following holds: if Π is NP-complete for the class \mathcal{G} of subcubic graphs, then Π is NP-complete for \mathcal{G}^{pq} for every $p \geq 1$. Now, Π is a *C123-problem* if (C1) Π is polynomial-time solvable for graphs of bounded treewidth; (C2) Π is NP-complete for subcubic graphs; and (C3) Π is NP-complete under edge subdivision of subcubic graphs.

In [20], it was shown that for every finite set of graphs \mathcal{H} , any C123-problem Π on \mathcal{H} -subgraph-free graphs is polynomial-time solvable if \mathcal{H} contains a graph from \mathcal{S} and NP-complete otherwise. Le and Telle [27] observed that PERFECT MATCHING CUT satisfies C1 and proved C2 and C3 (see Lemma 12). Applying the above meta-theorem from [20] yields:

Theorem 20 *For any finite set of graphs \mathcal{H} , PERFECT MATCHING CUT on \mathcal{H} -subgraph-free graphs is polynomial-time solvable if \mathcal{H} contains a graph from \mathcal{S} and NP-complete otherwise.*

5.3 Open Problems

Apart from completing the classifications of Theorems 16–18, we also pose the following open problem.

5.3.1 Open Problem 1

Classify the computational complexity of MATCHING CUT and DISCONNECTED PERFECT MATCHING for \mathcal{H} -subgraph-free graphs.

We note that classifications of MATCHING CUT and DISCONNECTED PERFECT MATCHING are unknown even for H -subgraph-free graphs (so when we forbid only a single graph H as a subgraph). So far, only a partial classification for MATCHING CUT restricted to \mathcal{H} -subgraph-free graphs has been shown [21]. In particular, the transformations for MATCHING CUT and DISCONNECTED PERFECT MATCHING from Sect. 4 do not decrease the girth and yield graphs with many cycles of varying length as subgraphs. Hence, new techniques are needed.

Finally, with our current technique (Lemma 5) we cannot obtain a better bound on the maximum degree of the graphs of arbitrarily large fixed girth in the proofs of Theorems 6 and 11.

5.3.2 Open Problem 2

Can the two maximum degree bounds in Theorems 6 and 11 be improved?

Acknowledgements We thank Hoang-Oanh Le for a significant simplification of our original proof of Theorem 6, which we simplified a bit further. We thank Van Bang Le for observing the bound on the maximum degree in Theorem 6, solving an Open Problem Garden question.

Author Contributions All four authors contributed to the paper.

Funding Open access funding provided by University of Fribourg

Data Availability Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Declarations

Conflict of interest The authors have no Conflict of interest as defined by Springer, or other interests that might be perceived to influence the results and/or discussion reported in this paper.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Alon, N., Milman, V.D.: λ_1 , isoperimetric inequalities for graphs, and superconcentrators. J. Comb. Theory Ser. B **38**, 73–88 (1985)
2. Araújo, J., Cohen, N., Giroidre, F., Havet, F.: Good edge-labelling of graphs. Discret. Appl. Math. **160**, 2502–2513 (2012)

3. Aravind, N.R., Kalyanasundaram, S., Kare, A.S.: Vertex partitioning problems on graphs with bounded tree width. *Discret. Appl. Math.* **319**, 254–270 (2022)
4. Aravind, N.R., Saxena, R.: An FPT algorithm for Matching Cut and d -Cut. In: *Proc. IWOCA 2021*, LNCS, vol. 12757, pp. 531–543. (2021)
5. Bonnet, E., Chakraborty, D., Duron, J.: Cutting barnette graphs perfectly is hard. In: *Proc. WG 2023*, LNCS, to appear
6. Bonsma, P.S.: The complexity of the matching-cut problem for planar graphs and other graph classes. *J. Graph Theory* **62**, 109–126 (2009). ((conference version: WG 2003))
7. Borowiecki, M., Jesse-Józefczyk, K.: Matching cutsets in graphs of diameter 2. *Theoret. Comput. Sci.* **407**, 574–582 (2008)
8. Bouquet, V., Picouleau, C.: The complexity of the Perfect Matching-Cut problem. [arxiv:2011.03318](https://arxiv.org/abs/2011.03318) (2020)
9. Chen, C.-Y., Hsieh, S.-Y., Le, H.-O., Le, V.B., Peng, S.-L.: Matching cut in graphs with large minimum degree. *Algorithmica* **83**, 1238–1255 (2021)
10. Chvátal, V.: Recognizing decomposable graphs. *J. Graph Theory* **8**, 51–53 (1984)
11. Dirichlet, P.L.: Beweis des Satzes, dass jede unbegrenzte arithmetische Progression, deren erstes Glied und Differenz ganze Zahlen ohne gemeinschaftlichen Faktor sind, unendlich viele Primzahlen enthält. *Abhandlungen der Königlich Preussischen Akademie der Wissenschaften zu Berlin*, (1837)
12. Dodziuk, J.: Difference equations, isoperimetric inequality and transience of certain random walks. *Trans. Am. Math. Soc.* **284**, 787–794 (1984)
13. Farley, A.M., Proskurowski, A.: Networks immune to isolated line failures. *Networks* **12**, 393–403 (1982)
14. Feghali, C.: A note on Matching-Cut in P_t -free graphs. *Inf. Process. Lett.* **179**, 106294 (2023)
15. Golovach, P.A., Komusiewicz, C., Kratsch, D., Le, V.B.: Refined notions of parameterized enumeration kernels with applications to matching cut enumeration. *J. Comput. Syst. Sci.* **123**, 76–102 (2022)
16. Golovach, P.A., Paulusma, D., Song, J.: Computing vertex-surjective homomorphisms to partially reflexive trees. *Theoret. Comput. Sci.* **457**, 86–100 (2012)
17. Gomes, G., Sau, I.: Finding cuts of bounded degree: complexity, FPT and exact algorithms, and kernelization. *Algorithmica* **83**, 1677–1706 (2021)
18. Graham, R.L.: On primitive graphs and optimal vertex assignments. *Ann. N. Y. Acad. Sci.* **175**, 170–186 (1970)
19. Heggenes, P., Telle, J.A.: Partitioning graphs into generalized dominating sets. *Nordic J. Comput.* **5**, 128–142 (1998)
20. Johnson, M., Martin, B., Oostveen, J.J., Pandey, S., Paulusma, D., Smith, S., van Leeuwen, E.J.: Complexity framework for forbidden subgraphs I: The framework. [arxiv:2211.12887](https://arxiv.org/abs/2211.12887) (2022)
21. Johnson, M., Martin, B., Pandey, S., Paulusma, D., Smith, S. and Van Leeuwen, E.J.: Complexity framework for forbidden subgraphs III: When problems are tractable on subcubic graphs. In: *Proc. MFCS 2023, LIPIcs*, Vol 272, pp. 57:1–57. (2023)
22. Komusiewicz, C., Kratsch, D., Le, V.B.: Matching cut: kernelization, single-exponential time FPT, and exact exponential algorithms. *Discret. Appl. Math.* **283**, 44–58 (2020)
23. Kratsch, D., Le, V.B.: Algorithms solving the Matching Cut problem. *Theoret. Comput. Sci.* **609**, 328–335 (2016)
24. Le, H.-O., Le, V.B.: A complexity dichotomy for Matching Cut in (bipartite) graphs of fixed diameter. *Theoret. Comput. Sci.* **770**, 69–78 (2019)
25. Le, H.-O., Le, V. B.: Complexity results for matching cut problems in graphs without long induced paths. In: *Proc. WG, LNCS*, to appear (2023)
26. Le, V.B., Randerath, B.: On stable cutsets in line graphs. *Theoret. Comput. Sci.* **301**, 463–475 (2003)
27. Le, V.B., Telle, J.A.: The perfect matching cut problem revisited. *Proc. WG 2021 LNCS* **12911**, 182–194 (2021)
28. Lubotzky, A., Phillips, R., Sarnak, P.: Ramanujan graphs. *Combinatorica* **8**, 261–277 (1988)
29. Lucke, F., Paulusma, D., Ries, B.: On the complexity of Matching Cut for graphs of bounded radius and H -free graphs. *Theor. Comput. Sci.* **936**, 33–42 (2022)
30. Lucke, F., Paulusma, D., Ries, B.: Dichotomies for maximum matching cut: H -Freeness, bounded diameter, bounded radius. *Proc. MFCS 2023, LIPIcs*, **272**, 64:1–64:15 (2023)
31. Lucke, F., Paulusma, D., Ries, B.: Finding matching cuts in H -free graphs. *Algorithmica*, to appear
32. Moshé, A.M.: Matching cutsets in graphs. *J. Graph Theory* **13**, 527–536 (1989)

33. Open problem garden. http://www.openproblemgarden.org/op/matching_cut_and_girth. Accessed 22 June 2023
34. Patrignani, M.: The complexity of the Matching-Cut problem. Proc. WG 2001 LNCS **2204**, 284–295 (2001)
35. Shapiro, H.N.: Introduction to the Theory of Number. Dover Publications, New York (2008)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.